

NEW TAXA AND NEW COMBINATIONS IN JUNGERMANNIALES (HEPATICAE)

НОВЫЕ ТАКСОНЫ И НОВЫЕ КОМБИНАЦИИ ПЕЧЕНОЧНИКОВ ИЗ ПОРЯДКА JUNGERMANNIALES (НЕРАТИКАЕ)

N.A.KONSTANTINOVA¹ & A.A.VILNET¹

Н.А.КОНСТАНТИНОВА¹, А.А. ВИЛЬНЕТ¹

Abstract

The names of one new family (*Hygrobiellaceae* (Jørg.) Konstant. & Vilnet) and 7 new genera (*Pseudolophozia* Konstant. & Vilnet, *Pseudotritomaria* Konstant. & Vilnet, *Schljakovia* Konstant. & Vilnet, *Schljakovianthus* Konstant. & Vilnet, *Lophoziopsis* Konstant. & Vilnet, *Biantheridion* (Grolle) Konstant. & Vilnet, *Heterogemma* (Jørg.) Konstant. & Vilnet) are validated. Correspondingly, 21 new combinations are made in these genera.

Резюме

Обнародованы названия одного нового семейства (*Hygrobiellaceae* (Jørg.) Konstant. & Vilnet) и 7 новых родов (*Pseudolophozia* Konstant. & Vilnet, *Pseudotritomaria* Konstant. & Vilnet, *Schljakovia* Konstant. & Vilnet, *Schljakovianthus* Konstant. & Vilnet, *Lophoziopsis* Konstant. & Vilnet, *Biantheridion* (Grolle) Konstant. & Vilnet, *Heterogemma* (Jørg.) Konstant. & Vilnet), а также 21 новая комбинация в этих родах.

KEYWORDS: liverworts, taxonomy, nomenclature, new taxa

The last ten years are characterized by a breathtaking speed of development of molecular study of hepaticas. As a result, the ideas about phylogeny of hepaticas have changed drastically (Davis et al., 2004; Yatsentyuk et al., 2004; H-Nygren et al., 2006; Forrest et al., 2006; Hentschel et al., 2007; De Roo et al., 2007; Crandall-Stotler et al., 2008; Vilnet et al., 2008, 2009). Many genera traditionally accepted as a “good” ones were found to be polyphyletic, which forced to search for new taxonomic solutions for their placement. Our study of Jungermanniales (Vilnet et al., 2008, 2009, unpublished data) is one those, that demonstrate a high level of incongruence between traditional taxonomy and novel ideas of their relationships. This require a certain rearrangement of taxa, a number of new combinations and description of several new

genera and one new family for such recently found groups. These names are needed for the simultaneously published checklist of liverworts of Russia (Konstantinova, Bakalin et al., 2009).

NEW FAMILY

Hygrobiellaceae (Jørg.) Konstant. & Vilnet, *stat. nov.* – Based on *Cephaloziaceae* Mig. subfam. *Hygrobielloideae* (Jørg.) R.M.Schust. ex Grolle, J. Bryol. 7: 210. 1972.

The family includes one genus, *Hygrobiella* Spruce.

NEW GENERA

Pseudolophozia Konstant. & Vilnet, *gen. nov.* – *Genus hoc generi Lophoziae s. str. affine sed foliis trilobatis frequenter alatis, cellulis caulinum saepiuscule homogeneis in sectione transversali, gemmis rubello-fuscis vel ferrugineis et ITS1-2 nrDNA et trnL-F cpDNA sequentiae differt.*

¹ – Polar-Alpine Botanical Garden, Kola Sci. Center of Russian academy of Sciences, Kirovsk-6, Murmansk Province 184256 Russia – Россия 184256, Кировск-6 Мурманской области, Полярно-альпийский ботанический сад-институт КНЦ РАН; e-mail: nadya50@mail.ru & anya_v@list.ru

Typus: Pseudolophozia sudetica (Nees ex Huebener) Konstant. & Vilnet. (*Jungermannia sudetica* Nees ex Huebener).

Species: Pseudolophozia sudetica (Nees ex Huebener) Konstant. & Vilnet, *comb. nov.* – *Jungermannia sudetica* Nees ex Huebener, Hepaticol. Germ. 142 Mannheim. 1834.

Pseudolophozia sudetica var. *anomala* (Schljakov) Konstant. & Vilnet, *comb. nov.* – *Lophozia alpestris* (F. Weber) Evans var. *anomala* Schljakov, Novosti Syst. Nizъ. Rast. 11: 352. 1974.

Pseudolophozia debiliformis (R.M. Schust. & Damsh.) Konstant. & Vilnet, *comb. nov.* – *Lophozia debiliformis* R.M. Schust. & Damsh., Phytologia 63(5): 326. 1987.

Pseudolophozia debiliformis var. *concolor* (R.M. Schust. & Damsh.) Konstant. & Vilnet, *comb. nov.* – *Lophozia debiliformis* var. *concolor* R.M. Schust. & Damsh., Phytologia 63: 326. 1987.

Pseudotritomaria Konstant. & Vilnet, *gen. nov.* – *Planta minima, 1.0-1.4(-1.6) mm lata, foliis subsymmetricis, trilobatis, parietibus cellularum in angulis perincrassatis.* Genus hoc genera *Tritomaria* et *Saccobasis* appropinquat sed ab eis marginibus foliorum perichaetalium undulato-crispatis, spinuloso-dentatis vel spinuloso-ciliatis et ITS1-2 nrDNA et trnL-F cpDNA sequentia differt.

Typus: Pseudotritomaria heterophylla (R.M. Schust.) Konstant. & Vilnet (*Tritomaria heterophylla* R.M. Schust.).

Pseudotritomaria heterophylla (R.M. Schust.) Konstant. & Vilnet, *comb. nov.* – *Tritomaria heterophylla* R.M. Schust., Canad. J. Bot. 36: 172. 1958.

Schljakovia Konstant. & Vilnet, *gen. nov.* – Genus hoc generi Orthocauli simile sed foliis profunde bilobatis, caulinis gemmiferis haud modificatis et ITS1-2 nrDNA et trnL-F cpDNA sequentia recedit.

Typus: Schljakovia kunzeana Konstant. & Vilnet (*Jungermannia kunzeana* Huebener).

Schljakovia kunzeana Konstant. & Vilnet, *comb. nov.* – *Jungermannia kunzeana* Huebener Hepaticol. Germ.: 115. 1834.

Schljakovianthus Konstant. & Vilnet, *gen. nov.* – Genus hoc generi Orthocauli simile sed

foliis quadrilobatis ad 0.6 laminae partitis, marginibus sinuum distincte revolutis, cuticula per-papillosa, gemmis nullis et ITS1-2 nrDNA et trnL-F cpDNA sequentia differt.

Typus: Schljakovianthus quadrilobus (Lindb.) Konstant. & Vilnet (*Jungermannia quadriloba* Lindb.).

Schljakovianthus quadrilobus (Lindb.) Konstant. & Vilnet, *comb. nov.* – *Jungermannia quadriloba* Lindb., Meddeland Soc. Fauna Fl. Fenn. 9: 162, 1889 (nom. subnud.) et in Arnell & Lindb., Kongl. Svenska Vetenskapsakad. Handl. 23 (5): 55. 1889.

Schljakovianthus quadrilobus var. *glareosa* (Jiurg.) Konstant. & Vilnet, *comb. nov.* – *Jungermannia quadriloba* var. *glareosa* Jiurg., Bergens Mus. Erbok 18. s. p. 1895.

Schljakovianthus quadrilobus var. *collenchymatica* (R.M. Schust.) Konstant. & Vilnet, *comb. nov.* – *Lophozia quadriloba* (Lindb.) A. Evans var. *collenchymatica* R.M. Schust. Hepat. Anthocer. N. Amer. 2: 281. 1969.

Lophoziopsis Konstant. & Vilnet, *gen. nov.* – Genus hoc generi *Lophoziae* s. str. simile sed ab eo gemmis rubris ad purpurascensibus vel rubello-fuscis, cellulis leptodermaticis, bracteis femineis saepe dentatis et ITS1-2 nrDNA et trnL-F cpDNA sequentia recedit.

Typus Lophoziopsis excisa (Dicks.) Konstant. & Vilnet (*Jungermannia excisa* Dicks.). *Lophoziopsis excisa* (Dicks.) Konstant. & Vilnet, *comb. nov.* – *Jungermannia excisa* Dicks., Pl. Crypt. Brit. Fasc. 3: 11. 1793.

Lophoziopsis excisa var. *elegans* (R.M. Schust.) Konstant. & Vilnet, *comb. nov.* – *Lophozia excisa* (Dicks.) Dumort. var. *elegans* R.M. Schust., Hepat. Anthocer. N. Amer. 2: 522. 1969.

Lophoziopsis excisa var. *infuscata* (R.M. Schust. & Damsh.) Konstant. & Vilnet, *comb. nov.* – *Lophozia excisa* (Dicks.) Dumort. var. *infuscata* R.M. Schust. & Damsh., Meddel. Gr?nland 199: 94. 1974.

Lophoziopsis excisa var. *succulenta* (R.M. Schust. & Damsh.) Konstant. & Vilnet, *comb. nov.* – *Lophozia excisa* (Dicks.) Dumort. var. *succulenta* R.M. Schust. & Damsh., Meddel. Gr?nland 199: 96. 1974.

Lophoziopsis longidens (Lindb.) Konstant. & Vilnet, *comb. nov.* – *Jungermannia longidens*

Lindb., Helsigfors Dagblad 1876 (323): 2. 1876.
et Bot. Not. 27. 1877.

Lophoziopsis pellucida (R.M.Schust.) Konstant. & Vilnet, *comb. nov.* – *Lophozia pellucida* R.M.Schust., Canad. J. Bot. 39: 978. 1961.

Lophoziopsis personii (H.Buch & S.W.Arnell) Konstant. & Vilnet, *comb. nov.* – *Lophozia personii* H.Buch & S.W.Arnell in Buch, Bot. Not. 1944: 382, figs. 1-2. 1944.

Lophoziopsis polaris (R.M.Schust.) Konstant. & Vilnet, *comb. nov.* – *Lophozia alpestris* (Schleicher ex F. Weber) A.Evans ssp. *polaris* R.M. Schust., Hepat. Anthocer. N. Amer. 2: 614, New-York, 1969.

Lophoziopsis polaris var. *sphagnorum* (R.M.Schust.) Konstant. & Vilnet, *comb. nov.* – *Lophozia alpestris* (Schleicher ex F. Weber) A.Evans ssp. *polaris* R.M. Schust. f. *sphagnorum* R.M.Schust., Hepat. Anthocer. N. Amer. 2: 619, fig. 218: 15-19. 1969.

Lophoziopsis propagulifera (Gottsché) Konstant. & Vilnet, *comb. nov.* – *Jungermannia propagulifera* Gottsché, Int. Polarforsch., Deutsch. Exped. 2: 451, 1890.

Lophoziopsis rubrigemma (R.M.Schust.) Konstant. & Vilnet, *comb. nov.* – *Lophozia rubrigemma* R.M.Schust., Hepat. Anthocer. N. Amer. 2: 621. 1969.

Biantheridion (Grolle) Konstant. & Vilnet, *stat. nov.* – *Jamesoniella* sect. *Biantheridion* Grolle, Trans. Brit. Bryol. Soc. 4, 4: 662. 1964

Biantheridion undulifolium (Nees) Konstant. & Vilnet, *comb. nov.* – *Jungermannia schraderi* var. *undulifolia* Nees, Naturgesch. Eur. Leberm. 1: 306, 1833.

Heterogemma (Jørg.) Konstant. & Vilnet, *stat. nov.* – *Lophozia* sect. *Heterogemma* Jørg., Bergens Mus. Skr. 16: 1959, 1934.

Heterogemma laxa (Lindb.) Konstant. & Vilnet, *comb. nov.* – *Jungermannia laxa* Lindb., Acta Soc. Sci. Fenn. 10: 539. 1875.

Heterogemma capitata (Hook.) Konstant. & Vilnet, *comb. nov.* – *Jungermannia capitata* Hook., Brit. Jungermann.: Tab. 80. 1816.

ACKNOWLEDGEMENTS

We thank M.S. Ignatov, J. Váňa, L. Söderström, A. Hagborg for numerous comments and suggestions. Dr. R. Ochyra is cordially thanked for trans-

lation of Latin diagnoses. This work was partly supported by RFBR 09-04-00281, 06-05-64137.

LITERATURE CITED

- CRANDAL-STOTLER, B.J., R.E. STOTLER & D.G. LONG 2008. Morphology and classification of the Marchantiophyta. – Goffinet, B. & A.J.Shaw (eds.) *Bryophyte Biology*, 2 ed. Cambridge: 1-54.
- DAVIS, E.C. 2004. A molecular phylogeny of leafy liverworts (Jungermanniidae: Marchantiophyta). – *Monogr. Syst. Bot. Missouri Bot. Garden* **98**: 61-86.
- DE ROO, R.T., T.A. HEDDERSON & L. SÖDERSTRÖM 2007. Molecular insights into the phylogeny of the leafy liverwort family Lophoziacae Cavers. – *Taxon* **56**: 301-314.
- FORREST, L.L., E.C.DAVIS, D.G.LONG, B.J. CRANDAL-STOTLER, A. CLARK & M.L. HOLLINGSWORTH 2006. Unravelling the evolutionary history of the liverworts (Marchantiophyta): multiple taxa, genomes and analyses. – *Bryologist* **109**: 303-334.
- HENTSCHEL, J., J.A. PATON, H. SCHNEIDER & J. HEINRICHS 2007. Acceptance of *Lioclada* Nees and *Solenostoma* Mitt., the systematic position of *Eremotus* Pearson and notes on *Jungermannia* L. s.l. (Jungermanniidae) based on chloroplast DNA sequence data. – *Pl. Syst. Evol.* **268**: 147-157.
- HE-NYGREN X., A. JUSLÉN, D., GLENNY & S. PIIPPO 2006. Illuminating the evolutionary history of liverworts (Marchantiophyta) – towards a natural classification. – *Cladistics* **22**: 1-31.
- KONSTANTINOVA, N.A., V.A. BAKALIN ET AL. 2009. Checklist of liverworts (Marchaniophyta) of Russia. – *Arc-tica* **18**: 1-66.
- [VILNET, A.A., N.A. KONSTANTINOVA & A.V. TROITSKY] ВИЛЬНЕТ А.А., Н.А. КОНСТАНТИНОВА, А.В. ТРОИЦКИЙ 2007. К молекулярной филогении семейства Gymnomitriaceae H.Klinggr. (Hepaticae). – [On molecular phylogeny of Gymnomitriaceae H.Klinggr. (Hepaticae)] Мат. международн. конф. «Вычислительная филогенетика и геносистематика «ВФГС»» КМК, Москва [Proceedings of Conference “Computational phylogenetics and molecular systematics “COMS’2007”, KMK, Moscow]: 24-26.
- VILNET, A.A., N.A. KONSTANTINOVA & A.V. TROITSKY 2008. Phylogeny and systematics of the genus *Lophozia* s. str. (Dumort.) Dumort. (Hepaticae) and related taxa from nuclear ITS1-2 and chloroplast *trnL*-F sequences. – *Mol. Phylogen. Evol.* **47**: 403-418.
- [VILNET, A.A., N.A. KONSTANTINOVA & A.V. TROITSKY] ВИЛЬНЕТ А.А., Н.А. КОНСТАНТИНОВА, А.В. ТРОИЦКИЙ 2009. Геносистематика и новый взгляд на филогению и систему печеночников. – [Genosystematics and new insight on phylogeny and taxonomy of liverworts] Молек. биол. [Molec. Biol.] **43** (5): 1-12.
- YATSENTYUK, S.P., N.A. KONSTANTINOVA, M.S. IGNATOV, J. HYVÖNEN & A.V. TROITSKY 2004. On phylogeny of Lophoziacae and related families (Hepaticae, Jungermanniales) based on *trnL-trnF* intron-spacer sequences of chloroplast DNA. – *Monographs in Systematic Botany from the Missouri Botanical Garden* **98**: 150-167.