

The spider genus *Clubiona* Latreille, 1804 (Arachnida Aranei Clubionidae) in  
the USSR fauna: a critical review with taxonomical remarks.

Пауки рода *Clubiona* Latreille, 1804 (Arachnida Aranei Clubionidae) фауны  
СССР: критический обзор с таксономические замечания.

K.G.Mikhailov

К.Г.Михайлов

Zoological Museum of the Moscow State University, Herzen str., 6, Moscow К-9 103009 Russia.  
Зоологический музей Московского университета, ул. Герцена, 6, Москва К-9, 103009 Россия.

KEY WORDS: *Clubiona*, new species, catalogue, taxonomical remarks, USSR fauna.

КЛЮЧЕВЫЕ СЛОВА: *Clubiona*, новые виды, каталог, таксономические замечания, фауна СССР.

**ABSTRACT:** A full revised faunistical list of the spider genus *Clubiona* of ex-USSR including 71 species is given. *C.transbaicalica* sp.n. (♂, Buryatia), *C.pseudosaxatilis* sp.n. (♂♀, E-Kazakhstan, Tuva), *C.subborealis* sp.n. (♂♀, S-Siberia, Mongolia), *C.komissarovi* sp.n. (♀, Primorye), *Crybinus* sp.n. (♂♀, S-Kazakhstan, Kirghizia), *C.saurica* sp.n. (♀, E-Kazakhstan), *C.bashkirica* sp.n. (♂♀, Bashkiria) are described. A catalogue of citations of all USSR species as well as the materials for new regions are given. The entire genus is divided into 3 subgenera, one of them, *Clubiona* s.str. is subdivided into 12 species groups. The world distribution of all the cited species and infrageneric groupings is characterized. The female of *C.wunderlichi* sp.n. from Mongolia is described. *C.maracandica* Kroneberg, 1875 (♀) is redescribed. New synonymy is established: *C.lyubarskii* Michailov, 1991 = *C.kunashirensis* Michailov, 1990, syn.n., *C.subinteriecta* Strand, 1907 = *C.propinqua* L.Koch, 1879, syn.n.

**РЕЗЮМЕ:** В статье дан полный отривизованный фаунистический список пауков рода *Clubiona* СССР, включающий 71 вид, в том числе *C.transbaicalica* sp.n. (♂, Бурятия), *C.pseudosaxatilis* sp.n. (♂♀, Восточный Казахстан, Тыва), *C.subborealis* sp.n. (♂♀, юг Сибири, Монголия), *C.komissarovi* sp.n. (♀, Приморье), *Crybinus* sp.n. (♂♀, юг Казахстана, Киргизия), *C.saurica* sp.n. (♀, Восточный Казахстан), *C.bashkirica* sp.n. (♂♀, Башкирия). Приведен каталог литературных указаний по распространению в СССР всех форм, а также материалы по новым для данного вида регионам. Все виды рода разделены на три подрода; подрод *Clubiona* s.str. подразделен на 12 групп видов. Охарактеризовано мировое распространение всех упомянутых в тексте видов и внутриродовых категорий. Дано описание самки *C.wunderlichi* sp.n. из Монголии. Переописана *C.maracandica* Kroneberg, 1875 (♀). Установлена новая синонимия: *C.lyubarskii*

Michailov, 1991 = *C.kunashirensis* Michailov, 1990, syn.n., *C.subinteriecta* Strand, 1907 = *C.propinqua* L.Koch, 1879, syn.n.

The aim of this paper is to give a general faunistical list (including catalogue) of USSR *Clubiona*. Some partial faunistical reports [Mikhailov, 1990ab, 1991] as well as the general list prepared in 1989 (without catalogue) [Mikhailov, 1992] have been published. During the last two years a lot of new materials have been studied, especially after visiting the Zoological Institute of the USSR Academy of Sciences in Leningrad (since 1992: of the Russian Academy of Sciences in St.-Petersburg) and the Institute of Zoology, Polish Academy of Sciences, Warsaw, both in autumn 1991. I am deeply indebted to the curators of these arachnid collections, Dr. V.I.Ovtsharenko and Prof. Dr. J.Proszynski, respectively, for their kind assistance, as well as to all persons who supply me by *Clubiona* collections.

The majority of the materials treated below were collected by Mr. A.M.Basarukin (AMB), Ms. V.A.Bragina (VAB), Dr. S.N.Danilov (SND), Dr. P.M.Dunin (PMD), Mr. V.E.Efimik (VEE); Dr. K.Yu.Eskov (KYE), Dr. S.L.Esyunin (SLE); Mr. G.N.Ganin (GNG), Mr. D.K.Kurenschikov (DKK), Dr. D.V.Logunov (DVL), Mr. O.V.Lyakhov (OVL), Dr. Yu.M.Marusik (YMM), Ms. R.Meldazhite (RM), Dr. K.G.Mikhailov (KGM), Mr. S.V.Ovchinnikov (SVO), Mr. F.Z.Popov (FZP), Dr. N.M.Poryadina (NMP), Mr. A.B.Ryvkin (ABR), Dr. L.G.Savelieva (LGS), the late Dr. V.K.Soldatov (VKS), the late Prof. S.A.Spassky (SAS), Dr. I.D.Sukacheva (IDS), the late Prof. V.P.Tyshchenko (VPT), Mr. N.A.Utkin (NAU), Dr. E.M.Zhukovets (EMZ), Mr. S.L.Zonshtain (SLZ). The above names are referred to below only by the respective abbreviations.

The material studied is kept in the following museums and collections: Zoological Museum of the Moscow State University (ZMUM), Zoological In-

stitute of the Russian Academy of Sciences, St.-Petersburg (ZIP), Biological Institute, Siberian Division of the Russian Academy of Sciences, Novosibirsk (BIN), Chair of Invertebrate Zoology, Perm State University (PSU), Institute of Biology, Buryatian Academy of Sciences, Ulan-Ude (UU), Institute of Water and Ecological Problems, Khabarovsk (Kh), Institute of Zoology, Polish Academy of Sciences, Warsaw (IZPAN), Naturhistorisches Museum Wien (NHMW), Zoological Museum, University of Helsinki, Finland (ZMH). The above mentioned abbreviations are used in "Material".

In the "Catalogue", the following sequence and abbreviations of geographical units are accepted:

Russia: MRM - Murmansk Area, KRL - Karelia, LNN - Leningrad Area, PSK - Pskov Area, NVG - Novgorod Area, KLN - Kalinin (= Tver) Area, KLG - Kaliningrad Area, KLZh - Kaluga Area, YaRS - Yaroslavl Area, MSK - Moscow Area, VLD - Vladimir Area, TLS - Tula Area, RZN - Ryazan Area, BRN - Bryansk Area, ORL - Orel Area, KUR - Kursk Area, LPTs - Lipetsk Area, TMB - Tambov Area, BLG - Belgorod Area, VRN - Voronezh Area, GRK - Gorkiy (= Nizhniy Novgorod) Area, MRS - Mariyskaya (= Mari-El) Republic, ChVSh - Chuvashia, MRD - Mordovia, ULN - Ulyanovsk (= Simbirsk) Area, PNZ - Penza Area, TTR - Tartarian ASSR, KBSH - Kuybyshev (= Samara) Area, VLG - Volgograd Area, AST - Astrakhan Area, RST - Rostov Area, KLM - Kalmykia, KRS - Krasnodar Province, STV - Stavropol Province, KBA - Kabardino-Balkaria, SOA - N-Osetia, ChIA - Checheno-Ingushetia, DAG - Dagestan, ARKh - Arkhangelsk Area, KOM - Komi Republic, VOL - Vologda Area, KST - Kostroma Area, KRV - Kirov (= Vyatka) Area, UDM - Udmurtia, PRM - Perm Area, SVR - Sverdlovsk (= Ekaterinburg) Area, BShK - Bashkiria, ChLB - Chelyabinsk Area, KRG - Kurgan Area, TMN - Tyumen Area, OMS - Omsk Area, NVS - Novosibirsk Area, TMS - Tomsk Area, KMR - Kemerovo Area, ALT - Altai Province, KRN - Krasnoyarsk Province, YaKT - Yakutia, MGD - Magadan Area, KMCh - Kamchatka Area, IRK - Irkutsk Area, BUR - Buryatia, ChIT - Chita Area, AMR - Amur Area, KhAB - Khabarovsk Province, PRI - Primorye Province, SAKh - Sakhalin Area.

#### Estonia. Latvia. Lithuania.

Byelorussia: GRD - Grodno Area, MIN - Minsk Area, VIT - Vitebsk Area, BRS - Brest Area, GOM - Gomel Area.

Ukraine: VOL - Volyn Area, LVV - Lvov Area, TRN - Ternopol Area, ZAK - Zakarpatskaya Area, IFR - Ivano-Frankovsk Area, ChRTs - Chernovtsi Area, RVN - Rovno Area, ZhTM - Zhitomir Area, KhML - Khmelnytskyi Area, SUM - Sumy Area, KVS - Kiev Area, ChRK - Cherkassy Area, PLT - Poltava Area, KhRK - Kharkov Area, KRG - Kirovograd Area,

DNP - Dnepropetrovsk Area, ODS - Odessa Area, KhRS - Kherson Area, ZPR - Zaporozhye Area, DNTs - Donetsk Area, VRSh - Voroshilovgrad Area, Krim (=Crimea).

#### Moldavia.

Georgia. Armenia. Azerbaijan.

Kazakhstan: URL - Uralsk Area, KST - Kustanay Area, KKCh - Kokchetav Area, PVL - Pavlodar Area, TsLN - Tselinograd (= Akmolinsk) Area, EKA - East-Kazakhstan Area, AAO - Alma-Ata Area.

Uzbekistan: SMR - Samarkand Area, SRD - Syrdarya Area, TShK - Tashkent Area.

Turkmenistan. Kirghizia. Tadzhikistan.

When a locality is situated at the border between two areas, the slash is used: BUR/ChIT: Muy Ketlle. The literature data proved by the collections are marked by an asterisk (\*). The data with only citations of former authors are added by the part "-cit": Thorell, 1875-cit.

The Catalogue was prepared in 1982-1991, so modern changes in geographical names (especially of 1991-1992) are not taken into consideration, with some exceptions (Georgia instead of Georgian SSR, etc.). The paper of Belimov [1975] on Yakutian spiders is ignored by me, since no Latin (only Russian!) names were included there.

In the text, the following abbreviations are used: F - femur, Pt - patella, Ti - tibia, Mt - metatarsus, d. - dorsally, v. - ventrally, l. - laterally, rl. - retrolaterally. All the measurements are given in mm.

#### Subgenus *Paracubiona* (Lohmander, 1945).

Type species: *Clubiona corticalis* (Walckenaer, 1802).

DISTRIBUTION. Palaearctic, Indo-Malayan, Australian regions. Europe - 1 species, Nepal - 1 species (not described: H.Ono, pers. comm.), Japan - Honshu (*C.kurosawai* Ono, 1986, *C.yaginumai* Hayashi, 1989), Ryukyu (*C.cirrosa* Ono, 1987), China (*C.lyriformis* Song et M.Zhu, 1991, *C.mordalis* Song et M.Zhu, 1991), Burma (*C.concinna* (Thorell, 1887)), etc.

*Clubiona (Paracubiona) corticalis* (Walckenaer, 1802)

MATERIAL. Azerbaijan: Nakhichevan: 2♀ (ZMUM), Shakhbzur distr., Kemyur, 1500 m, *Carpinus*, *Quercus* forest, 9.07.1988, PMD; 2♀ (ZMUM), Bichenek, 1900 m, *Carpinus*, *Quercus* forest, 13.07.1988, PMD; Armenia: 1♀ (ZIP, SAS collection), Khach-Tekh, 07.1926.

CATALOGUE. Ukraine: ZAK [Polozhentsev, Akimtseva, 1980]. Azerbaijan [Dunin\*, 1989; Mikhailov, 1990b].

Wrong determinations: Russia: MSK [Pokrovskiy, 1925; Mikhailov, 1983b-cit]. TTR [Utochkin, 1977].

DISTRIBUTION. USSR - Carpathians, Caucasus. S-, C-Europe, Scandinavia: S-Sweden.

Subgenus *Japoniona* Michailov, 1990.

Type species: *Clubiona japonica* L.Koch, 1878.

DISTRIBUTION. E.Asia - 4 species; *C.jucunda* (Karsch, 1879) needs a detailed redescription as forming a separate species group. The redescription of the male [Paik, 1990b: Figs. 30-32] is quite insufficient.

*Clubiona (Japoniona) japonica* L.Koch, 1878.

MATERIAL. Russia, Primorye: 1♂, Mikhailovka distr., Tarasovka, 27.07.1973, Kulikova; 1♀ (ZIP), environs of Kangauz (Anisimovka), 13.07.1968, FZP; 4♂ 2♀, Popova Island, Alexeeva Bay, 2-6.09.1991, KGM.

CATALOGUE. Russia: KhAB: Bolshekhekhtyrskiy State Reserve [Mikhailov, 1990a]. PRI: Terney, Kedrovaya Pad State Reserve, Khasan distr. [Mikhailov, 1990a].

DISTRIBUTION. USSR - continental S-Far East. All Japan, S-Korea, China (to Zhejiang).

*Clubiona (Japoniona) vigil* Karsch, 1879.

CATALOGUE. Russia: SAKh: Kunashir [Mikhailov, 1990a].

DISTRIBUTION. USSR - Kunashir. All Japan, S-Korea, China (?).

*Clubiona (Japoniona) flavipes* (Saito, 1939).

MATERIAL. 2♀, Russia, Sakhalin, Kholmsk distr., Slepikovskogo Cape, 6-8.07.1990, AMB.

CATALOGUE. Russia: SAKh: S-Sakhalin, Kunashir [Mikhailov, 1990a].

DISTRIBUTION. USSR - S-Sakhalin, Kunashir. All Japan.

*Clubiona (Japoniona) jucunda* (Karsch, 1879).

1879 *Liocranum jucundum* Karsch: 92.

1957 *C.jucunda* [sic!]: Yaginuma: pl.IV, fig.8 (♂).

1987 *C.propinqua*: Song: 325-326, fig.280A,B (♀ only!).

1990b C.j.: Paik: 71-73, figs.28-39 (♂, ♀).

MATERIAL. 1♀, Russia, Kunashir, Mendeleeva Volcano, 1-3.09.1987, AMB.

DISTRIBUTION. New for the USSR fauna. All Japan, S-Korea; N-Korea (original data; IZPAN).

Subgenus *Clubiona* sensu stricto.

Type species: *Clubiona pallidula* (Clerck, 1757).

DISTRIBUTION. Cosmopolitan (?). Data from S-America are doubtful. 15 species-groups in the Holarctic fauna, 3 of them (*maritima*, *abboti*, and *zilla*) are absent in USSR.

The *trivialis* - group.

DISTRIBUTION. Holarctic, Indo-Malayan region. USSR - 8 species. Europe; E-Asia beyond the limits of USSR: *C.insulana* Ono, 1989 (Ryukyu). N-America - 4 species. S-China: *C.baimensis* Song et M.Zhu, 1991, *C.duoconcava* Zhand et Hu, 1991, *C.subrostrata* Zhang et Hu, 1991.

*Clubiona (Clubiona) trivialis* C.L.Koch, 1843.

MATERIAL. Russia, Komi Republic: 1♂, Vorkuta distr., 6 km NW of Sivomaskinskiy, tundra, 0.5 km N of northern forest line, 9.07.1981, A.V.Tanasevitch; Tyumen Area: 1♀ (PSU), Surgut distr., Yuganskiy State Reserve, Pech-Pan-Yakh, a swamp, 07.1988, V.G.Novokshonov; Sakhalin: 1♀, 3-5 km E of Novoalexandrovsk, 9.07.1985, AMB; 1♀, Aniva distr., Pik Chekhova Mt., 1000 m, 3.07.1986, AMB. Latvia: 1♂, Tukums distr., environs of Berzciems, bank of Engures Lake, 12.08.1987, KGM.

CATALOGUE. Russia: MRM [Palmgren, 1943]. KRL [Palmgren, 1943; Uzenbaev, 1985, 1986, 1987]. LNN [Palmgren, 1943]. KLN: Ostashkov Uezd [Fedotov, 1912c], Belskiy Uezd [Pokrovskiy, 1930]. MSK [Freiberg, 1897; Mikhailov, 1983b-cit]. KOM: Pechoro-Ilychskiy State Reserve [Pakhorukov, 1980]. KST [Veselova, Mikhailov, 1987\*; Mikhailov, 1989]. UDM [Zubko, Roshchinenko, 1981]. N-Ural [Pakhorukov, 1979]. PRM: Basegi State Reserve [Esynin, 1991]. SVR [Pakhorukov, 1977]. KRN: Mirnoe, Evenkiya [Eskov, 1988\*]. MGD: environs of Magadan [Mikhailov, 1990a]. Estonia [Schenkel-Haas, 1924; Kauri, 1938; Vilbaste, 1957, 1972, 1973, 1974, 1980, 1987]. Lithuania [Vaitskute, 1960, 1963]. Byelorussia [Litvinova et al., 1980]. Ukraine: ZAK [Balogh, 1940].

Doubtful data. Russia: KLM [Minoranskiy, Ponomarev, 1984]. Ukraine: S part [Astakhova, 1978], Krim [Spassky, 1927].

DISTRIBUTION. USSR - all European part (except extreme south), C- and NE-Siberia, Sakhalin. S-Europe: Italy (Toscana), C-, N-Europe.

*Clubiona (Clubiona) diversa* O.Pickard-Cambridge, 1862.

MATERIAL. Russia: 1♀, Leningrad Area, Stariy Peterhof, 7.05.1983, YMM; 1♀, Krasnodar Province, Caucasian State Reserve, ca. 12 km SE of Guzeripl, Abago Plateau, subalpine belt, 1880-1900 m, 15.06.1991, KGM; Novosibirsk Area: 1♀, Toguchin distr., Kotorovo, meadow, litter, 19.07.1987, Ya.Barushkina: 1♀, Novosibirsk distr., Koltsovo, margin of *Betula* forest, in grass, 29.05.1988, leg. et det. DVL; Tuva: 6♂, ca. 5 km W of Kyzyl, agrobiological station, 650-700 m, meadow, in grass, 23.07.1989, DVL; 2♂ 1♀ Tes-Khem distr., 10 km NW of Khol-Oozhu, Belenghit, *Larix* stand, 1800 m, 9-11.07.1989, DVL; 1♂, Piy-Khem distr., ca. 5 km NW of Sesarlig, ca. 25 km NW of Kyzyl, *Larix* - *Salix* forest, along a stream, 1100-1200 m, 24.07.1989, DVL; 1♀ (BIN 47), Ulug-Khem distr., 8 km S of Torgalygh, stepped hills, 1100-1250 m, 9.05.1990, DVL; Krasnoyarsk Province, Khakassia: 1♀, Askiz distr., 25-27 km NE of Askiz, weathered rock near Abakan River, mesophytous meadow, 15.08.1990, DVL; 2♂ 2♀ (BIN 62) Shira distr., ca. 3-5 km E of Shira, Itkul Lake, mesophytous meadow, in grass, 21-22.07.1990, DVL; 2♀ (BIN 63) 6♀, Altayskiy distr., ca. 40 km SE of Belyi Yar, ca. 15-17 km

E of Novorossiyskoe, ca. 400 m, *Festuca sulcata* steppe, 23-24.06.1990, DVL; Chita Area, Kyra distr., Sokhondo State Reserve: 1♂, Ingoda River Valley, *Alnus* stand, 1350-1400 m, 8-9.06.1991, leg. et det. DVL; 1♀, Tsanginanduy River Valley, "tundrosteppe", 1350-1400 m, 10.06.1991, leg. et det. DVL. 1♂, Ukraine, Zakarpatskaya Area, Perechin distr., Lumshora, 900 m, 11.08.1981, leg. et det. YMM. Kazakhstan: 1♂ (ZIP), Akmolinsk Area, Kokshetau, *Salix*, 20.06.1957, VPT; 2♂ 4♀ Alma-Ata, Medeo, 2000 m, S slope, litter, 27.08.1983, YMM; 1♀, East-Kazakhstan Area, Zaysan distr., Saur Mt.Ridge, Karaungur River (Kenderlyk Basin), montane meadow, ca. 1800 m, 19.06.1990, KYE.

Mongolia: 1♂, Eastern Aymak, Sumber somon, Bulin River, 10 km E of Kharkhont Frontier Post, 31.07.1986, K.Ulykpan.

CATALOGUE. Russia: MSK [Greze, 1914; Mikhailov, 1983b-cit]. BRN [Kharitonov, 1936]. KUR [Pichka, 1984ab]. TTR [Zheltukhina, Utochkin, 1984]. STV: Teberda State Reserve [Mikhailov, 1990b]. SOA [Mikhailov, 1990b]. UDM [Zubko, Roshchinenco, 1981]. PRM [Azheganova, Glukhov, 1981]: Basegi State Reserve [Esyunin, 1991]. ChLB: Troitskiy Reserve [Azheganova, 1951; Utochkin, Golovashkina, 1977; Pakhorukov, 1985], Ilmen State Reserve [Polyanin, Pakhorukov, 1988]. KRN: Evenkiya [Eskov, 1988\*]. MGD: Kolyma upper flow, environs of Magadan [Mikhailov, 1990a]. KhAB: Okhotsk distr. [Mikhailov, 1990a]. Estonia [Vilbaste, 1957]. Lithuania [Zhukauskene, 1966]. Byelorussia [Litvinova et al., 1980; Litvinava et al., 1981]: VIT: Berezina State Reserve [Shlyakhtyonok, 1986; Pristavko, Zhukovets, 1987]. Azerbaijan: Pirkuli State Reserve [Mikhailov, 1990b].

DISTRIBUTION. USSR - all European part, Caucasus, Siberia. S-Europe: Balkans, Italy, C-, N-Europe, Mongolia, S-Korea, Japan.

#### *Clubiona (Clubiona) subtilis* L.Koch, 1866.

MATERIAL. Russia, Novosibirsk Area: different localities; Tuva: 1♂ 1♀, Erzin distr., ca. 3-5 km S of Erzin, Tes-Khem River floodplain, 1100 m, litter, 14.08.1989, DVL; Chita Area, Kyra distr., Sokhondo State Reserve: 1♀, Agutsa River valley, swampy meadow, 1100 m, 16.06.1991, leg. et det. DVL. Kirghizia: 1♂ 1♀, Frunze, Karagachevaya Roshcha, 15.05.1979, SLZ; 2♀, Frunze, Baytyk Valley, 08.1978, SLZ.

Mongolia: 1♀, Selenginskiy Aymak, Shamor, Orkhon River floodplain, 31.07.1982, KGM.

CATALOGUE. Russia: KRL [Uzenbaev, 1985, 1986, 1987]. TTR [Zheltukhina, Utochkin, 1984]. KRV [Garkusha, 1980]. SVR [Pakhorukov, 1977]. ChLB: Troitskiy Reserve [Azheganova, 1951], Ilmen State Reserve [Polyanin, Pakhorukov, 1988]. AMR: Arkhara, Khinghan State Reserve [Mikhailov, 1990a]. KhAB: Bolshevikheksyrskiy State Reserve [Mikhailov, 1990a]. Estonia [Vilbaste, 1959, 1964, 1969, 1972, 1973, 1980, 1987]. Byelorussia: MIN [Zhukovets, 1989]. Ukraine: KhRK [Kirilenko, Legotay, 1981].

DISTRIBUTION. USSR - C-, N-European part, W-, S-Siberia, Kirghizia. S-Europe (Italy), C-, N-Europe, N-Mongolia.

REMARK. *C.subtilis* sensu Paik, 1990 constitutes a separate species not conspecific with the European *subtilis*.

#### *Clubiona (Clubiona) juvenis* Simon, 1878.

MATERIAL. Lithuania: 1♂ 2♀, Shilutee, Vente Ornithological Station, 1984, RM; 1♂, Zhuvintas State Reserve, Sphagneto-Phragmitetum, 16.06.1986, leg. RM, det. EMZ. Ukraine: 3♀ (ZIP), environs of Berdyansk, 15.05.1938, leg. V.Nikolaev, det. SAS.

CATALOGUE. Russia: KRS: Slavyanka distr. [Mikhailov, 1990b]. Estonia [Vilbaste, 1987]. Ukraine: steppes near Black Sea [Mikityuk, 1982]. Uzbekistan: SRD [Mikhailov, 1990b].

DISTRIBUTION. USSR - W-, S-European part, Middle Asia. C-Europe, C-China (Kansu).

#### *Clubiona (Clubiona) basarukini* Michailov, 1990.

MATERIAL. Russia, Buryatia: 1♀ (UU) Ulan-Ude, Vakhmistrovo, pine forest, 31.05.1983, SND; 2♀ (UU), Bichura distr., Okino-Klyuchi, pine forest, 31.05.1983, SND; 2♀ (ZMUM), ca. 15 km W of Ulan-Ude, Mostovoy, pine forest, 11.08.1983, SND.

Mongolia: 1♀ sbad (ZMUM) Selenga Aimak, Shamor, pine forest, 1.08.1982, KGM.

CATALOGUE. S-Sakhalin [Mikhailov, 1990a].

DISTRIBUTION. USSR - Transbaikalia, S-Sakhalin. N-Mongolia.

#### *Clubiona (Clubiona) rostrata* Paik, 1985.

MATERIAL. Russia, Khabarovsk Prov.: 1♀ (Kh), Nanayskiy distr., Slavyanka, Siberian pine-broadleaved forest, 29.05.1990, DKK; Primorye: 1♂ (ZIP), Vladivostok, 04-06.1903, N.Palchevskiy; 1♀ (ZIP), Shkotovo distr., environs of Kangauz (Anisimovka), coniferous-broadleaved forest, 13.07.1968, FZP; 1♂ (ZIP), Kangauz, in a room, 6.07.1968, FZP.

CATALOGUE. Russia: KhAB: Bolshevikheksyskiy State Reserve [Mikhailov, 1990a]. PRI [Mikhailov, 1990a]. SAKh: S-Sakhalin [Mikhailov, 1990a].

DISTRIBUTION. USSR - S-Far East (continental and island). All Japan, S-Korea, N-Korea (original data; IZPAN), China: Zhejiang.

#### *Clubiona (Clubiona) transbaicalica* Michailov, sp.n.

Fig. 1a,b.

MATERIAL. Russia, Buryatia: 1♂ (holotype ZMUM Ta-4801), Selenga distr., Taezhniy, 17.08.1984, BPZ; 1♂ (paratype, ZMUM Ta-4802), Ulan-Ude, Vakhmistrovo, pine forest, 31.05.1983, SND.

DIAGNOSIS. By the structure of the palp the male is more close to *C.rostrata*, differing by both the relatively little developed tibial apophysis and the shorter embolus; the latter's apical part never reaches the basal half of the bulbus.

DESCRIPTION. MALE (2). Total length 5.30-6.10. Carapace length 2.35-2.75, width 1.65-2.00, their ratio 1.38-1.42. Carapace, chelicerae, and legs reddish-brown. Leg measurements:

	I	II	III	IV
Femur	1.70-2.03	1.80-2.13	1.38-1.95	2.03-2.65
Patella	0.90-1.15	0.90-1.13	0.75-0.90	0.88-1.08
Tibia	1.68-2.13	1.78-2.28	1.13-1.48	1.65-2.08
Metatarsus	1.20-1.53	1.33-1.63	1.38-1.75	2.13-2.68
Tarsus	0.55-0.80	0.65-0.83	0.50-0.60	0.68-0.78

Leg armature: F I d.1.1.2 (1.1.3), II d.1.1.3 (1.1.2), III-IV d.1.1.3, Pt III-IV rl.1, Ti I-II v.2.2, III d.2.2, v.1.1, IV d.2.2, v.2.1.1 (1.1.1, 1.2.1), Mt I-II v.2, III d.2.1.2, l.1.2, v.2.2, IV d.2.1.2, l.2.2, v.2.1.2.

Palp as in Fig. 1a,b. Lengths of cymbium, tibia, patella, and femur are 0.75-0.95, 0.25-0.35, 0.33-0.40, 0.63-0.93, respectively.

Abdomen length 2.90-3.55, width 1.75-1.85, their ratio 1.66-1.92; dark brown.

FEMALE unknown.

REMARK. The paratype is considerably less than the holotype.

DISTRIBUTION. USSR - Transbaikalia.

*Clubiona (Clubiona) amurensis* Michailov, 1990.

MATERIAL. 1♀ (ZIP), Russia, S-Primorye, midstream of Kedrovka River, 16.05.1968, FZP.

CATALOGUE. Russia, AMR: Khinghan State Reserve [Mikhailov, 1990a].

DISTRIBUTION. USSR - continental S-Far East. Japan: Hokkaido.

REMARK. The male is described by Matsuda [1991: 22-24, Figs. 1, 3-4].

The *marmorata* - group.

DISTRIBUTION. Palaearctic, 2 species.

*Clubiona (Clubiona) marmorata* L.Koch, 1866.

MATERIAL. 1♀, Moldavia, Strasheny distr., Panasheshty, 13.08.1985, Maevskaya.

CATALOGUE. Byelorussia [Litvinova et al., 1980\*]: BRS




Fig.1. *Clubiona (Clubiona) transbaicalica* sp.n., male palp: a) ventral view; b) external view. Scale = 0.14.  
Рис.1. *Clubiona (Clubiona) transbaicalica* sp.n., пальпа самца: а) вентрально; б) снаружи. Масштаб 0.14.

[Litvinova, 1978\*]. Ukraine: IFR + LVV [Legotay, Tarasyuk, 1964]. Krim [Spassky, 1927; Kharitonov, 1937; Bukovskiy, 1940].

DISTRIBUTION. USSR - S-, W-European part. S-Europe (Balkans, Italy), C-Europe to Denmark.

*Clubiona (Clubiona) hyrcanica* Michailov, 1990.

CATALOGUE. Azerbaijan: Hyrcan State Reserve [Mikhailov, 1990b].

DISTRIBUTION. Type locality only.

The *brevipes* - group.

DISTRIBUTION. Palaearctics, 2 species.

*Clubiona (Clubiona) brevipes* Blackall, 1841.

MATERIAL. Russia: 1♀ (ZMUM) Stavropol Prov., 40 km W of Pyatigorsk, Suvorovskaya, in a "nest" of *Hyphantria cunea*, 11.06.1983, E.A.Prokofyeva.

CATALOGUE. Russia: MSK [Freiberg, 1894; Mikhailov, 1983b-cit]. Byelorussia: VIT: Berezinskiy State Reserve [Shlyakhtyonok, 1986]. Ukraine: ZAK [Legotay, 1980; Bulyginskaya, Kalinkin, 1985: *Clubiona*; Kalinkin, 1985]. Georgia: Mariamdvare State Reserve [Mikhailov, 1990b]. Azerbaijan: Kutkashen distr. [Mikhailov, 1990b].

Wrong identifications. Russia: SAKh: N-Kurile Isls. [Saito, 1932, 1933].

DISTRIBUTION. USSR - W-European part, Caucasus. S-, C-Europe; N-Europe: S-Sweden.

*Clubiona (Clubiona) alexeevi* Michailov, 1990.

MATERIAL. Russia: 1♂ (ZIP), Kabardino-Balkaria, Nalchik, 06-07.1928, Lobodina. Erroneously determined by S.A.Spassky as *C.brevipes*.

CATALOGUE. Russia: SOA [Mikhailov, 1990b].

DISTRIBUTION. N-Caucasus, C part.

The *comta* - group.

DISTRIBUTION. Palaearctic, ca. 10 species. Subgroup *genevensis*: *C.genevensis* L.Koch, *C.vegeta* Simon, 1878, *C.diniensis* Simon, 1878, *C.leucaspis* Simon, 1932, *C.wunderlichi* sp.n., and Macaronesian *C.decora* Blackwall, 1859, *C.minor* Wunderlich, 1987, *C.pseudominor* Wunderlich, 1987.

*Clubiona (Clubiona) comta* C.L.Koch, 1839.

MATERIAL. Russia, Krasnodar Prov.: 1♀ (ZMUM), Goryachi Klyuch, Defanovka, *Quercus*, *Carpinus* forest, litter, 18.05.1983, S.I.Golovach.

CATALOGUE. Estonia [Vilbaste, 1974, 1987]. Latvia [Sternbergs, 1981a]. Byelorussia [Litvinova et al., 1980; Litvinova et al., 1981]. VIT [Pereleshina, 1930\*]. Ukraine: Carpathians [Legotay, 1973]. ZAK [Legotay, 1958, 1989]. IFR + LVV [Legotay,

Tarasyuk, 1964].

Doubtful data. Russia: MSK [Greze, 1911; Mikhailov, 1983b-cit; Terrestrial invertebrates, 1989].

DISTRIBUTION. USSR - W-European part, N-Caucasus. S-, C-, N-Europe, Algeria.

*Clubiona (Clubiona) genevensis* L.Koch, 1866.

MATERIAL. Russia: 1♀ (PSU), Bashkiria, Bashkirian State Reserve, montane steppe, 13.06.1988, leg. VEE, det. SLE; 2♂, Krasnoyarsk Prov., Khakassia, Altayskiy distr., ca. 40 km SE of Belly Yar, ca. 15-17 km E of Novorossiyskoe, ca. 400 m, *Festuca sulcata* steppe, 23-24.06.1990, DVL. 1♂, Kazakhstan, Zaisan distr., Saur Mt.Ridge, Akkolka River Valley (Kenderlyk Basin), dry stony steppe, 5-17.06.1990, KYE.

CATALOGUE. Latvia [Prieditis, Sternbergs, 1981: *C.decora*; Sternbergs, 1983: *C.decora*]. Armenia [Mikhailov, 1990b]. Azerbaijan [Mikhailov, 1990b]. Kazakhstan [Fet, 1983\*]. Turkmenistan: SW Kopetdagh Mts. [Fet, 1983\*; Mikhailov, Fet, 1986]. Kirghizia [Zonstein, 1984\*]. Tadzhikistan [Andreeva, 1975, 1976: *C.vegeta*].

Doubtful data. Russia: LPTs: Galichya Gora State Reserve [Panteleeva, 1982].

DISTRIBUTION. USSR - W-European part, Caucasus, Kazakhstan, Middle Asia, Khakassia. S-, C-Europe; N-Europe: S-Sweden, Algeria, Morocco, Palestine.

*Clubiona (Clubiona) wunderlichi* Michailov, sp.n.

Fig. 2 a,b.

MATERIAL. Mongolia: 1♀ (holotype ZMUM Ta-4803), Sukhe-Bator Aimak, Tumentsoqt, *Stipa grandis*, 7.07.1983, K.Ulykpan.

DIAGNOSIS. The species is more close to *C.genevensis*, differing by both the angular curve of the copulatory tubes in posterior angles of the epigyne and the atria situated mesally than spermathecae. The copulatory tube opens to the atrio-spermathecal part laterally, not inferiorly, as in both *C.genevensis* and *C.vegeta* [cp. Helsdingen, 1979, Figs.7,9].

DESCRIPTION. FEMALE. Total length 7.00. Carapace length 3.65, width 2.30, their ratio 1.59. Carapace and legs pale, chelicerae reddish-coloured. Palp atypically swollen. Leg measurements:

	I	II	III	IV
Femur	1.50	1.70	1.43	2.13
Patella	1.00	1.03	0.83	1.00
Tibia	1.23	1.38	0.85	1.48
Metatarsus	0.98	1.05	0.98	1.75
Tarsus	0.58	0.60	0.45	0.65

Leg armature: F I-II d.1.1.2, III-IV d.1.1.3, Pt III-IV rl.1, Ti I v.0, II v.1.1, III d.2, IV d.2.2, v.1.1, Mt I v.0, II v.1, III d.2.2, 1.2, v.2.2, IV d.2.1.2, 1.2.2, v.1.2. Such a pattern "aberrant" for *Clubiona* is due to the small size of the spider: a lot of spines are reduced.

Abdomen: length 3.50, width 2.25, their ratio 1.56; pale. Epigyne as in Fig.2 a,b.

MALE unknown.

DISTRIBUTION. The single locality in Mongolian steppes.

DERIVATIO NOMINIS. The new species is dedicated to the famous German arachnologist, Jorg Wunderlich.

The *lutescens* - group.

DISTRIBUTION. Holarctic. Palaearctic - 9-10 species. Outside USSR - *C.saxatilis* L.Koch, 1867 (= *C.dvoraki* Miller, 1943), *C.bilaris* Simon, 1878, maybe *C.andreinii* Caporiacco, 1936 (described by a single male from Perudgia, Italy).

*Clubiona (Clubiona) lutescens* Westring, 1851.

*C.neglecta* - Greze, 1911; Pereleshina, 1928a.

MATERIAL. Russia, Ryazan Area: Okskiy State Reserve (leg. et det. KYE, 1981); Komi Republic: 2♀, ca. 60 km SE of Syktyvkar, Kortkeros distr., Dan, Lokchim River floodplain, 17.22.06.1988, KGM; Perm Area (different localities, PSU, det. SLE); Kurgan Area: 1♀, Uval, Tobol River floodplain, 8.06.1989, NAU; Tyumen Area: 1♂, Yarkovo distr., 70 km SW of Tobolsk, Mazurovo, 1.07.1982, NMP; Novosibirsk Area: different localities; Kemerovo Area: 1♂ 1♀, 60-70 km NE of Kemerovo, Zolotoy Kitoy River floodplain, *Picea*, *Betula*, *Salix* forest, 20.06.1990, DVL; Krasnoyarsk Prov., Khakassia: 1♀ (BIN 56), 2♀, Askiz distr., Birikchul, Askiz River floodplain, *Salix*, under bark, 1100 m, 18.07.1990, DVL; 1♀, Altay distr., ca. 40 km SE of Belyi Yar, Berezovskoe Lake, 350-400 m, salt meadow with *Caragana*, 22-23.06.1990, DVL; 1♂ 3♀, id., ca. 17 km E of Novorossiyskoe, Yenisey River floodplain, poplar forest, litter, 24.06.1990, DVL; Yakutia: 1♂, Sangar, wet *Betula* stand, 23-28.06.1989, leg. KYE, det. YMM; Buryatia: 1♂ 1♀ (BIN 2), Selenga River delta, Murzino, 10.07.1985, BPZ; 1♀ (UU), Selenginsk, *Betula* stand, 27.07.1983, leg. et det. SND; Khabarovsk Prov.: 1♂ (Kh) 1♀, Ulcha distr., environs of Zhrebtsovo, *Calamagrostis* in *Salix* stand, 26.06.1990, DKK. Lithuania: 1♀, Zhuvintas State Reserve, Pineto-Ledetum, 26.06.1986,

leg. RM, det. EMZ. Kazakhstan: 1♂ 5♀, Zaysan distr., Saur Mts., Karaungur River (Kenderlyk River Basin), plain poplar forest, 18-22.06.1990, KYE.

CATALOGUE. Russia: KRL [Palmgren, 1943; Tselarius, Shorokhov, 1985; Uzenbaev, 1986]. LNN [Palmgren, 1943]. KLN: Bologoe [Kharitonov, 1936\*]. KLZh [Vagner, 1895]. MSK [Vagner, 1892; Greze, 1911\*: CL+ C.n.; 1915b; Pereleshina, 1928a\*: CL+ C.n.; Beer, 1968; Sharova, 1970; Mikhailov, 1983ab]. KUR [Pichka, 1984ab]. LPTs: Galichya Gora State Reserve [Panteleeva, 1982]. BLG: Valuyki [Kulczynski, 1913], "Les na Vorskle" State Reserve [Ovsyannikov, 1937]. VRN [Pichka, 1965]. ChVSh [Holzmayer, 1934]. PNZ [Pyatin, 1988]. TTR [Garkusha, 1977; Utochkin, 1977; Zheltukhina, Utochkin, 1984]. KBSh: Zhiguli State Reserve [Krasnobaev, Ovtsharenko, 1986]. RST [Spassky, 1914, 1919, 1925, 1940]. KRS [Mikhailov, 1990b]. STV: Teberda State Reserve [Mikhailov, 1990b]. SOA [Mikhailov, 1990b]. ChIA [Minoranskiy, Ponomarev, Slyusarev, Gramotenko, 1984]. DGS [Mikhailov, 1990b]. KOM: Pechorollichsky State Reserve [Utochkin, Pakhorukov, 1976; Pakhorukov, 1980]. KST [Veselova, 1986\*; Veselova, Mikhailov, 1986, 1987; Mikhailov, 1989]. UDM [Zubko, Roshchinenco, 1981]. PRM: Basegi State Reserve [Esynin, 1991]. ChLB: Troitskiy Reserve [Pakhorukov, 1985], Ilmen State Reserve [Pakhorukov, Polyanin, 1987; Polyanin, Pakhorukov, 1988]. TMN: S-Yamal [Greze, 1909b; Bakhvalov, Korshunov 1976-cit], between Tobolsk and Obdorsk (Salekhard) [Bergroth, 1881], Tobolsk [Spassky, Lavrov, 1928], Tobol floodplain [Shlykov, 1978], Tura Plain [Shlykov, 1975]. TMS [Ermolajev, 1934]. KRN: Antsiferovo (Yenisey River, 59° 10' N) [Holm, 1973], Mirnoe [Eskov, 1988], mouth of Podkamennaya Tunguska River [Eskov, 1988\*]. AMR: Kundur [Mikhailov, 1990a]. Estonia [Vilbaste, 1965, 1974, 1987]. Latvia [Sternbergs, 1979, 1981ab; Prieditis, Sternbergs, 1981]. Byelorussia [Litvinova et al., 1980]. Belovezhskaya Pushcha State Reserve [Chebotareva, 1987]. MIN [Zhukovets, 1989\*]. VIT [Pereleshina, 1930\*]. Ukraine: LVV [Wajgiel, 1868, 1874; Nowicki, 1870]. TRN [Gnelitsa, 1989]. ZAK [Legotay, 1958, 1980, 1989; Polozhentsev, Akimtseva, 1980]. IFR [Nowicki, 1874]. ChRTs [Rosca, 1930, 1936]. KhML: Slavuta [Fedotov, 1912a], Proskurov [Pereleshina, 1930\*]. KVS [Pereleshina, 1930\*]. DNP: "Ekaterinoslav" (= Dnepropetrovsk) [Thorell,




Fig.2. *Clubiona (Clubiona) wunderlichi* sp.n., female: a) epigyne; b) vulva. Scale = 0.14.

Рис.2. *Clubiona (Clubiona) wunderlichi* sp.n., самка: а) эпигина; б) вульва. Масштаб 0.14.

1875]. KhRS [Talanov, Nazarenko, 1989]. VRSh [Polchaninova, 1992]. Moldavia [Talitskiy, Legotay, 1982; Matsyuk, Yanyuk, Verlan, 1984]. Georgia [Mikhailov, 1990b]. Armenia [Mikhailov, 1990b]. Azerbaijan [Dunin, 1989; Mikhailov, 1990b]. Kazakhstan: KKCh: Borovoe [Spassky, Lavrov, 1928]. EKA [Savelieva\*, 1970, 1979].

Doubtful data. Russia: SAKh: S-Sakhalin [Saito, 1934].

**DISTRIBUTION.** USSR - European part, Caucasus, Siberia. S-Europe: Balkans, Italy, C-, N-Europe, Japan (a single reliable locality - Saga: Bösenberg, Strand, 1906), N-America: British Columbia (Canada), Washington (USA).

*Clubiona (Clubiona) alpicola* Kulczynski, 1882.

*C.frutetorum* - Dunin, 1984, 1989; Fet, 1983; Koch, 1866 (part); Mikhailov, Fet, 1986; Thorell, 1875 (part).

**MATERIAL.** Ukraine: 1♂ (ZMUM) Krim, Simferopol, in a garden, 5.05.1981, VAB; 1♀ (ZIP) Sevastopol, Pliginsky.

**CATALOGUE.** Ukraine: ZAK [Balogh, 1940]. IFR + LVV [Legotay, Tarasyuk, 1964]. "Caucasus" [L.Koch, 1866: *C.frutetorum*; Thorell, 1875-cit: *C.frutetorum*]. Georgia: Lagodekhi State Reserve [Mikhailov, 1990b]. Armenia [Mikhailov, 1990b]. Azerbaijan [Dunin, 1984, 1989: *C.frutetorum*; Mikhailov, 1990b]. Turkmenistan: SW-Kopetdagh Mts. [Mikhailov, Fet, 1986: *C.frutetorum*; Fet, 1983: *C.frutetorum*, 1985; Mikhailov, 1990b].

**DISTRIBUTION.** USSR - Carpathians, Krim, Caucasus, Kopetdagh Mts. C-Europe, Bulgaria, Rumania. The species may be found also in Turkey.

*Clubiona (Clubiona) frutetorum* L.Koch, 1866.

**MATERIAL.** Russia, Leningrad Area, Luga: 9♀ (ZIP 3799), 8-14.07.1926, leg. E.Pokrovskaya, det.D.E.Kharitonov; 1♀ (ZIP 2916), on a Syringa, 31.07.1929, leg.E.Pokrovskaya, det. V.E.Pereleshina; 1♀ (ZIP), Mariyskaya Republic, Sovetskiy distr., Fokino, 15.07.1961; 1♀, Komi Republic, Kortkeros distr., ca. 60 km SE of Syktyvkar, Dan, forest margin, *Betula* stand, 21.06.1988, KGM. 1♀, Ukraine, Cherkassy Area, Kanev State Reserve, 11.09.1981, KGM.

**CATALOGUE.** Russia: KRL [Palmgren, 1943; Uzenbaev, 1986]. KLN: Ostashkov [Fedotov, 1912c], Belskiy Uezd [Pokrovskiy, 1930]. MSK [Vagner, 1892; Freiberg, 1897; Greze, 1914; Pereleshina, 1928a\*; Beer, 1968; Mikhailov, 1983b]. BLG: Valuyki [Kulczynski, 1913], "Les na Vorskle" State Reserve [Marusik, 1984]. PNZ [Pyatin, 1988]. TTR [Zheltukhina, Utochkin, 1984]. RST [Spassky, 1914, 1919]. PRM [Kharitonov, 1926a; Azheganova, Glukhov, 1981]. ChLB: Troitskiy Reserve [Azheganova, 1951; Pakhorukov, 1985]. KRN: Stolby State Reserve [Sternbergs, 1977]. Estonia [Vilbaste, 1969, 1972, 1973, 1974, 1980, 1987]. Latvia [Prieditis, Sternbergs, 1981; Sternbergs, 1981a, 1983]. Byelorussia [Litvinova et al., 1980]. Belovezhskaya Pushcha State Reserve [Kokhmanyuk, Yaroshchuk, 1981; Chebotareva, 1987]. GOM: Rogachev [Pereleshina, 1928b\*]. Ukraine: ZAK [Polozhentsev, Akimtseva, 1980; Legotay, 1989]. KhRK [Kirilenko, Legotay, 1981]. KhRS: Chernomorskiy State Reserve [Polchaninova, 1988\*]. DNTs: Amvrosievka [Spassky, 1914, 1919]. Moldavia [Babidorich, 1980; Talitskiy, Legotay,

1982]. Kazakhstan: EKA [Savelieva, 1970\*, 1979\*].

Doubtful identifications. Black Sea Coast of the Caucasus [Spassky, 1937] - may be actually *C.alpicola*.

Wrong identifications. "Caucasus" [L.Koch, 1866; Thorell, 1875-cit]. Azerbaijan [Dunin, 1984, 1989]. Turkmenistan: SW Kopetdagh Mts. [Mikhailov, Fet, 1986; Fet, 1983]. All these data belong to *C.alpicola*.

Other wrong identifications. Russia: AMR [Azheganova, Stenchenko, 1977]. SAKh: N-Kurile Isls. [Saito, 1932, 1933].

**DISTRIBUTION.** USSR - European part, S of W- and C (?)-Siberia. All Europe.

*Clubiona (Clubiona) riparia* L.Koch, 1866.

*C.pallidula* - Eskov, 1988.

**MATERIAL.** Russia, Bashkiria: 1♀ (PSU), Shulghan-Tash State Reserve, bank of Belya River, stones, 4.08.1989, VEE; Tuva: 5♀, Tandinskiy distr., E bank of Chagytay Lake, 1000 m, *Carex* swamp, 29.06.1989, DVL; 1♂ 2♀, Erzin distr., 3-5 km S of Erzin, Tes-Khem River floodplain, 1100 m, 14.08.1989, DVL; 1♂ 2♀, Erzin distr., Tere-Khol Lake, a swamp, 29.05.1989, DLV; 1♂, Erzin distr., 20-25 km NW of Erzin, Dus-Khol Lake, 800-900 m, under stones, in a grass, 31.05.1989, DVL; Yakutia: 1♀, left bank of Aldan River, environs of Krestkhaldzhey, 6.09.1990, IDS; Magadan Area: 1♀ (ZIP), Anadyr; Kamchatka: 1♀ 3 juv. (syntypes of *C.picata*, IZPAN), "nad rz. Kamchatka", leg. B.Dybowski; 1♀, 2 juv. (IZPAN, *C.picata* from Kulczynski's collection), "nad rz. Kamchatka", leg. B.Dybowski; Chita Area, Kyra distr., Sokhondo State Reserve: 1♂ (BIN 95), Ernichniy Stream, valley *Betula*, *Larix* forest, 1400-1450 m, 2-3.06.1991, DVL; 1♀, Ingoda River floodplain, 1350m, 9.06.1991, DVL; 2♂, 3-5 km E of Kyra, swampy meadow in Kyra River floodplain, 800-850 m, 30.06.1991, DVL; Amur Area: 1♀ (ZIP), Dep River (left tributary of Zeya River), 07-08.1911, P.Rippas; Primorye: 1♀ sbad, lower flow of Kedrovka River, 20.06.1968, FZP; N-Kuriles: 6♀, Paramushir Isl., 26-31.07.1989, AMB; 2♀, Shumshu Isl., environs of Bolshoe Lake, 18-22.08.1989, AMB.

Mongolia: 1♀, 80 km NE of Ulan-Bator, S-Khangai Mts., S exposition, Tereldsh Kisher, leg. et det. S.Heimer.

**CATALOGUE.** Russia: KRN: Evenkiya [Eskov, 1988\*]. YaKT: Yana River mouth [Kulczynski, 1908\*]. MGD: Kolyma River upper flow, environs of Magadan [Mikhailov, 1990a]. KMCh [Kulczynski, 1885\*, 1926; Strand, 1906-cit; Schenkel, 1930; Syshevskaja, 1935\*; Mikhailov, 1990b]. AMR: Khingan State Reserve [Mikhailov, 1990a]. KhAB: S areas [Mikhailov, 1990a]. PRI: Khasan distr. [Mikhailov, 1990a]. SAKh: Sakhalin, Kunashir [Mikhailov, 1990a], Iturup [Peelle, Saito, 1932].

**DISTRIBUTION.** Boreal species. USSR - S-Ural, C-, E-Siberia, all Far East. Mongolia, Japan (Hokkaido), N-America from Alaska to Maryland and NewMexico, USA.

*Clubiona (Clubiona) terrestris* Westring, 1851.

**MATERIAL.** Byelorussia: 1♂ 2♀, Brest Area, Kamenets distr., Belovezhskaya Pushcha State Reserve, 7.09-15.10.1982, N.G.Dyachenko.

**CATALOGUE.** Ukraine: IFR + LVV [Legotay, Tarasyuk,

1964]. ZAK [Legotay, 1958, 1989]. Carpathians [Legotay, 1973]. Moldavia [Babidorich, 1980, 1983; Talitskiy, Legotay, 1982; Matsyuk, Yanyuk, Verlan, 1984].

Wrong identifications. Russia: MSK [Pokrovskiy, 1925; Mikhailov, 1983b-cit]. ChLB: Ilmen State Reserve [Pakhorkov, Polyanin, 1987; Polyanin, Pakhorukov, 1988]. KRN: Boguchany distr. [Izmailova, Verzhutskiy, 1981]. AMR [Azheganova, Stenchenko, 1977]. Ukraine: S-steppes [Astakhova, 1978].

DISTRIBUTION. USSR - W-European part. All Europe.

*Clubiona (Clubiona) golovatchi* Michailov, 1990.

CATALOGUE. Russia: KRS: Sochi, Krasnodar, Slavyanka distr. [Mikhailov, 1990b]. STV: Zheleznyodolsk [Mikhailov, 1990b]. SOA [Mikhailov, 1990b]. DGS [Mikhailov, 1990b]. Georgia: Lagodekhi State Reserve, Adjaria [Mikhailov, 1990b]. Armenia [Mikhailov, 1990b]. Azerbaijan [Mikhailov, 1990b].

DISTRIBUTION. USSR - Caucasus Major and Minor.

*Clubiona (Clubiona) pseudosaxatilis* Michailov, sp.n.

Figs.3 a-c, 4 a,b.

MATERIAL. Holotype: 1♂ (ZMUM Ta-4804), Russia, Tuva, Mongun-Taiginskij distr., 5-8 km SE of Mugur-Aksy, Kargy River floodplain, *Salix*, *Populus*, *Larix* forest, liter, 1700-1800 m, 20.05.1990, DVL, OVL. Paratypes: Russia, Tuva: 5♂ (ZMUM) 1♂ (ZIP), as holotype; 1♀ Barlyk River floodplain, stony stream with *Larix*, 6.06.1990, OVL. Kazakhstan, E-Kazakhstan Area: 2♀, Ust-Kamenogorsk, hills, bank of a pond, blackberries, 1-15.06.1989, LGS; 1♀ (ZMUM Ta-4805), Zaisan distr., Saur Mts, Karaungur River (Kenderlyk River Basin), valley poplar forest, 18-22.06.1990, KYE.

DIAGNOSIS. The male palp differs from the congeners' ones by a combination of the *lutescens*-like tibial apophysis and the *saxatilis*-like embolus beginning at the medial part of the bulb [cp. Miller, 1943, Taf.III, Fig.8; Thaler, 1981, Abb.41]. By the structure of genitalia the female differs by the combination of the *saxatilis*-like copulatory openings situated in the posterior part (not in the posterior margin) of the epigynal plate and the *lutescens*-like non-narrowed, a little rounded copulatory tubes. From *C.alpicola*, *C.frutetorum*, *C.terrestris*, and *C.bilaris* it differs by the relatively small size of the atrium; the latter's diameter is equal to the diameter of the copulatory tubes.

DESCRIPTION. MALE (7). Carapace: length  $2.58 \pm 0.36$ , width  $1.91 \pm 0.28$ , their ratio  $1.35 \pm 0.11$ ;




Fig.3. *Clubiona (Clubiona) pseudosaxatilis* sp.n., male palp: a) ventral view; b) extero-ventral view; c) protector-like structure, internal view. Scale = 0.14.

Рис.3. *Clubiona (Clubiona) pseudosaxatilis* sp.n., пальпа самца: а) вентрально; б) наружно-вентрально; в) протекторовидная структура, изнутри. Масштаб 0.14.




Fig.4. *Clubiona (Clubiona) pseudosaxatilis* sp.n., female: a) epigyne; b) vulva. Scale = 0.14.  
Рис.4. *Clubiona (Clubiona) pseudosaxatilis* sp.n., самка: а) эпигина; б) вульва. Масштаб 0.14.

straw-coloured. Chelicerae chestnut-coloured, legs straw-coloured. Leg measurements:

	I	II	III	IV
Femur	♂ 1.95±0.28 ♀ 2.05±0.66	2.06±0.23 2.06±0.85	1.71±0.22 1.89±0.48	2.39±0.32 2.54±0.96
Patella	♂ 1.05±0.18 ♀ 1.16±0.32	1.09±0.15 1.14±0.47	0.88±0.10 1.00±0.25	1.04±0.12 1.13±0.45
Tibia	♂ 1.84±0.27 ♀ 1.81±0.52	1.96±0.24 1.87±0.77	1.33±0.18 1.36±0.39	1.98±0.12 2.04±0.72
Metatarsus	♂ 1.32±0.24 ♀ 1.29±0.38	1.40±0.16 1.32±0.51	1.50±0.19 1.51±0.51	2.49±0.32 2.53±1.10
Tarsus	♂ 0.81±0.19 ♀ 0.83±0.20	0.84±0.08 0.81±0.24	0.60±0.07 0.65±0.15	0.75±0.05 0.83±0.22

Leg armature: F I d.1.1.2, II d.1.1.2 (1.1.3), III-IV d.1.1.3, Pt III-IV rl.1, Ti I v.2.2 (2.3), II v.2.2, III d.2.2, v.1.1, IV d.2.2, v.1.1.1, Mt I v.2 (1), II v.2, III d.2.1.2 (2.2.2), I.1.2, v.2.2, IV d.2.1.2, I.2.2, v.2.1.2.

Palp as in Fig. 3a-c. Lengths of cymbium, tibia, patella, and femur are 0.88±0.10, 0.36±0.06, 0.40±0.05, 0.80±0.10, respectively.

Abdomen: length 3.31±0.48, width 1.79±0.25, their ratio 1.85±0.21; pale brown.

FEMALE (4). Carapace: length 3.15±0.83, width 2.29±0.68, their ratio 1.38±0.07; straw-coloured or pale brown. Chelicerae brown or chestnut-coloured, legs straw coloured. Leg measurements see above. Leg armature as in male, but F II d.1.1.2 (1.1.1), Ti I v.2.2, II v.2.2 (1.2), III d.2.2, v.1.1 (1.2), IV d.2.2, v.1.1.1 (1.2.1), Mt I v.2, III d.2.1.2, I.1.2 (2.2), v.2.2.

Abdomen: length 4.26±1.76, width 2.53±1.07, their ratio 1.69±0.09; brown. Epigyne as in Fig. 4a,b.

REMARK. Female specimen from Saur Mts. of a less size than others.

DISTRIBUTION. E-Kazakhstan, Tuva.

ECOLOGY. Apparently the species is associated with valley forests, rather leaved than coniferous.

The *similis* - group.

DISTRIBUTION. Palaearctic, 6 species. Outside USSR - *C.aducta* Simon, 1932 (Pyrenees).

*Clubiona (Clubiona) similis* L.Koch, 1866.

MATERIAL. Russia, Bryansk Area: 1♂, Bezhetsk distr., Gorodishchi, a meadow, 28.04.1984, N.A.Kalinichev; Voronezh Area: 1♀, Kantemirovka distr., Mitrofanovka, near a stream, 17.07.1979, A.A.Zyuzin; Astrakhan Area: 1♀, environs of Cherniy Yar, 25.02-25.03.1912, V.Lisovich, B.Shumakov; Novosibirsk Area: 1♂, environs of Akademgorodok, 23-31.05.1986, leg. W.Schawaller, det. F.Renner.

CATALOGUE. Russia: KRL [Palmgren, 1943]. LNN [Palmgren, 1943]. KLZh [Vagner, 1895]. MSK [Vagner, 1892; Mikhailov, 1983b-cit]. BLG: "Les na Vorskle" State Reserve [Ovsyannikov, 1937]. TTR [Utochkin, 1977]. KBSH: Zhibuli State Reserve [Krasnobaev, Ovtsharenko, 1986\*]. RST [Spassky, 1914, 1919]. KRS: Krasnodar [Mikhailov, 1990b]. STV: Novopavlovsk, Teberda State Reserve [Mikhailov, 1990b]. ChIA [Minoranskiy, Ponomarev, Slyusarev, Gramotenko, 1984]. UDM [Zubko, Roshchinenco, 1981]. PRM [Azheganova, Glukhov, 1981]. ChLB: Troitskiy Reserve [Azheganova, 1951; Utochkin, Golovashkina, 1977]. TMN: Tobol floodplain [Shlykov, 1978]. Estonia [Vilbaste, 1987]. Latvia [Sternbergs, 1981a]. Ukraine: ZAK [Kolosvary, 1939]. Moldavia [Talitskiy, Legotay, 1982]. Georgia [Mkheidze, 1964]. Azerbaijan [Mikhailov, 1990b].

Doubtful identifications. Russia: BUR/ChIT: Muy Kettle [Verzhutskiy, Bessolitsyna, Seryshev, 1985]. AMR [Azheganova, Stenchenko, 1977]. Kazakhstan: AAO: Zailiyskiy Alatau Mts.

[Folkina, 1978].

DISTRIBUTION. USSR - European part, Caucasus, S of W-Siberia. All Europe, Mongolia.

*Clubiona (Clubiona) neglecta* O.Pickard-Cambridge, 1862.

*C.montana* - A.Becker, 1888; Nowicki, 1874; Thorell, 1875.

MATERIAL. Russia: Novosibirsk Area: different localities; Tuva: 3♂ 1♀, environs of Kyzyl, floodplain forest, 7.06.1989, DVL; 1♂ 1♀, 5 km W of Kyzyl, 650-700 m, *Lasiagrostis splendens*, 25.05.1989, DVL; 1♂ (BIN 48), id., 15-22.05.1990, DVL; 1♂ 4♀ (ZMUM), 1♀ (BIN 51), id., 1.07.1990, DVL; 1♀, id., meadow, 23.07.1989, DVL; 2♀ (BIN 55), id., 1.07.1990, DVL; 2♂ 6♀, Tandinskij distr., environs of Chagytay Lake, steppe, 1100 m, *Artemisia*, 28.06.1989, DVL; 1♀, Erzin distr., 3-5 km S of Erzin, Tes-Khem River floodplain, 1100 m, 14.08.1989, DVL; 2♂ 21♀, 15 km W of Erzin, Dus-Khol Lake, 800-900 m, 31.05, 13.08.1989, DVL; 1♂, 30 km W of Erzin, *Lasiagrostis splendens* near a lake, 1100-1150 m, 27.05.1989, DVL; Krasnoyarsk Prov., Khakassia: 2♀, Askiz distr., 25-27 km NE of Askiz, meadow on a rock, 15.08.1990, DVL; 1♂ 1♀, Altay distr., ca 40 km SE of Belyi Yar, Berezovskoe Lake, 350-400 m, salt meadow with *Caragana*, 22-23.06.1990, DVL; Chita Area: 1♀, Kyra, 19.06.1991, leg. et det. DVL; 1♀ (BIN 92), Sokhondo State Reserve, 3-5 km SW of Buninda, steppe on slopes, 1200-1300 m, 14-15.06.1991, leg. et det. DVL; Ukraine, Krim: 1♂, Simferopol distr., Krasnolesye, 11.07.1981, VAB; Kazakhstan: Kora, 08.1937, V.N.Shnitnikov. Kirghizia: 1♀, Frunze, Chon-Aryk, leg. Sarkisova; 1♀, environs of Frunze, 4.07.1980, SLZ; 1♂ (ZIP), Osh, 1-2.11.1933, V.N.Shnitnikov; 1♀, Issyk-Kul Lake, Cholpon-Ata, 6.08.1977, SLZ; 2♂, Issyk-Kul Lake, Kuturga, 27.07.1977, SLZ; 2♂ 6♀, Kungey-Alatoo Mt.Ridge, Chon-Uryukty, 2000-2500m, *Picea* forest belt, 21-22.06.1983, SVO; 3♀, Terskey-Alatoo Mt.Ridge, Dzhelandy, 22.08.1980, SLZ.

CATALOGUE. Russia: LPTs: Galichya Gora State Reserve [Panteleeva, 1982]. BLG: Valuyki [Kulczynski, 1913]. TTR [Utochkin, 1977]. KBSH: Zhiguli State Reserve [Krasnobaev, Ovtsharenko, 1986\*]. VLG: Sarepta [Thorell, 1875; A.Becker, 1888]. KLM [Minoranskiy, Ponomarev, 1984]. Semideserts of European part [Ponomarev, 1981]. KRS: Mikhailovskiy Pass [Spassky, 1937], Slavyanka distr. [Mikhailov, 1990b]. STV: Teberda State Reserve [Mikhailov, 1990b]. SOA [Mikhailov, 1990b]. DGS [Mikhailov, 1990b]. UDM [Zubko, Roshchinenko, 1981]. BShK: Bashkirian State Reserve [Pakhorkov, Efimik, 1988\*]. ChLB: Ilmen State Reserve [Pakhorkov, Polyanin, 1987; Polyanin, Pakhorkov, 1988]. NVS: Chany Lake [Spassky, Lavrov, 1928\*]. IRK: Nizhnee Priangarye [Dmitrienko, 1987]. BUR [Odenvall, 1901; Izmailova, 1989]. BUR/ChLT: Muy Kettle [Verzhutskiy, Bessolitsyna, Seryshev, 1985]. ChLT [Izmailova, 1989]. Estonia [Vilbaste, 1973, 1974, 1980, 1987]. Latvia [Sternbergs, 1981a]. Byelorussia: MIN [Zhukovets, 1989]. VIT: Berezina State Reserve [Shlyakhtyonok, 1986\*; Pristavko, Zhukovets, 1987\*]. Ukraine: ChRTs [Nowicki, 1874: C.m.]. KhRK + PLT [Astakhova, 1974]. Moldavia [Talitskiy, Legotay, 1982]. "Caucasus" [Koch, 1866: C.m.; Thorell, 1875: C.m.]. Georgia [Mkhidze, 1964]. Batumi [Mikhailov, 1990b]. Armenia [Mikhailov, 1990b]. Azerbaijan [Verzhbitskiy, 1902; Dunin, 1989; Mikhailov, 1990b].

Kazakhstan: EKA [Savelieva, 1979\*]. AAO [Spassky, Shnitnikov, 1937; Tarabaev, 1979].

Doubtful identifications. Russia: AMR [Azheganova, Stenchenko, 1977].

Wrong identifications. Russia: MSK [Greze, 1911; Pereleshina, 1928a; Mikhailov, 1983b-cit] - revised as *Clutescens*.

DISTRIBUTION. USSR - European part, Caucasus, mountainous Middle Asia, S of W- and C-Siberia. All Europe, Morocco, Afghanistan, China; Shansi, S-Korea.

*Clubiona (Clubiona) pseudosimilis* Michailov, 1990.

MATERIAL. Russia, Krasnodar Prov., Caucasian State Reserve, ca. 12 km SE of Ghuzeripl, Abago Plateau: 1♂ 8♀, upper forest line, 1750-1850 m, aspen-birch forest, 12-17.06.1991, KGM; 3♀, ca. 1780 m, hay multiherbaceous grassland, 12.06.1991, KGM; 2♂, sparse *Acer* stand near upper forest line, ca. 1780 m, 11.06.1991, KGM; 1♀, ca. 1900 m, subalpine multiherbaceous grassland near upper forest line, 15.06.1991, KGM.

CATALOGUE. Russia: STV: Teberda State Reserve [Mikhailov, 1990b]. SOA [Mikhailov, 1990b]. Georgia [Mikhailov, 1990b]. Armenia: Sevan [Mikhailov, 1990b]. Azerbaijan: Zakataly State Reserve [Mikhailov, 1980b].

DISTRIBUTION. USSR - Caucasus Major and Minor, middle and high mountain belts.

*Clubiona (Clubiona) germanica* Thorell, 1871.

MATERIAL. Russia: 1♂ (ZIP 96), Pskov Area, Porkhov Uezd, coniferous forest, 29.05.1904, leg. Zinov'yev, det. V.E.Pereleshina; Arkhangelsk Area: 1♀, Onega distr., environs of Ust-Kozha, wet forest, 1984, DVL; 2♀, Kozhozero, environs of Letniy Konets, 22.08.1984, DVL; 1♀ (ZMUM Ta-3254), Kaluga Area, Tarusa, 15.07.1924, S.V.Pokrovskiy; 1♂ (PSU), Chelyabinsk Area, Satka distr., mixed forest, 18.07.1984, leg. N.M.Pakhorkov, det. A.B.Polyanin; 1♀, Tyumen Area, Tarko-Sale, *Betula* forest, 23.07.1984, NMP; 1♂ (BIN), 1♂, Novosibirsk Area, Kolyvan distr., Boyarka, *Betula* forest, 23.07.1986, leg. et det. DVL; 1♀, Krasnoyarsk Prov., Kulingna River, 4.10.1989, ABR; 2♂ 9♀ (UU), Buryatia, Selenginsk, *Betula* forest, 24.08.1983, leg. et det. SND. 4♂ 3♀ (ZMUM), Kirghizia, Frunze, 1932, D.Dementiev;

CATALOGUE. Russia: MRM [Fedotov, 1912b; Palmgren, 1943]. KRL [Palmgren, 1943; Uzenbaev, 1986]. LNN [Kharitonov, 1928\*; Palmgren, 1943]. KLN: Ostashkov [Fedotov, 1912c], Belskiy Uezd [Pokrovskiy, 1930]. YaRS [Fedotov, 1915]. MSK [Greze, 1914; Pokrovskiy, 1925\*; Pereleshina, 1928a\*; Mikhailov, 1983ab]. LPTs: Galichya Gora State Reserve [Panteleeva, 1982]. VRN [Pichka, 1965, 1967]. ChVSh [Holzmayer, 1934]. KBSH: Zhiguli State Reserve [Krasnobaev, 1992]. TTR [Utochkin, 1977]. BLG: Sarepta [Thorell, 1875; A.Becker, 1888]. RST [Spassky, 1914, 1919; Minoranskiy, Gramotenko, Ponomarev, 1977]. KOM: Pechoro-Ilychskiy State Reserve [Pakhorkov, 1980]. PRM [Kharitonov, 1926a; Azheganova, Glukhov, 1981]. Basegi State Reserve [Esyunin, 1991]. SVR [Kharitonov, 1923; Pakhorkov, 1977]. BShK [Olshvang, Malozemov, 1987; Pakhorkov, Efimik, 1988\*]. TMN: Tura plain [Shlykov, 1975]. Between Tomsk and Krasnoyarsk [Koch, 1879; Holm, 1973].

KRN: Mirnoe [Eskov, 1988\*], Anninskoe (Yenisey River, 63° 30' N) [Koch, 1879; Holm, 1973]. BUR: Nizhne-Angarskiy distr. [Izmailova, 1989]. Estonia [Vilbaste, 1969, 1972, 1973, 1974, 1980, 1987]. Latvia [Prieditis, Sternbergs, 1981, 1982; Sternbergs, 1981a, 1982, 1983; Anchianova, Sternbergs, 1984]. Byelorussia [Litvinova et al., 1980]. Belovezhskaya Pushcha State Reserve [Kokhmanyuk, Yaroshchuk, 1981; Chebotareva, 1987]. VIT [Pereleshina, 1930\*]. Ukraine: LVV [Nowicki, 1874]. ZAK [Polozhentsev, Akimtseva, 1980]. ChRTs [Nowicki, 1874]. KhRS [Talanov, Nazarenko, 1989]. VRSh: Provalye [Greze, 1909a]. Georgia: Kolkhida State Reserve [Mikhailov, 1990b]. Kazakhstan: EKA [Savelieva\*, 1970, 1972b, 1979]. Semirechye [Spassky, Snnitnikov, 1937].

Doubtful identifications. Russia: AMR [Azheganova, Stenchenko, 1977].

DISTRIBUTION. USSR - European part, Caucasus, W-, C-Siberia. S-Europe: Balkans, Italy, C-, N-Europe.

#### *Clubiona (Clubiona) congentilis* Kulczynski, 1913.

1971 *Clubiona germanica*: Miller: Tab.XI, Fig.7.

MATERIAL. Russia, Moscow Area, environs of Pushchino: 6♀, Oka River floodplain, 31.05-1.06.1981, KGM; 5♀, Pushchino, 14.06-5.07.1982, V.E.Gokhman; Belgorod Area: 1♂ 1♀ (syntypes, NHMW-13999), "Valuiki, leg. Velitchkovsky, 6.08.1899"; Ros-tov Area: 3♂ 3♀ (ZIP), environs of Novocherkassk, Persianovka, 25.05.1911, 28.05.1913, SAS, N.Karpova. Kazakhstan: N-Kazakhstan Area: 1♀ (BIN), Sokolovka distr., Bolshaya Malyshka, 12-18.06.1986, DVL; Pavlodar Area: 1♂ 1♀ (P), 40 km S of Pavlodar, environs of Yamyshev, forest plantation, 12-13.09.1989, OVL; Akmolinsk Area: 1♀ (ZIP), Kokshetau, Terisakan River bank, 20.06.1957, VPT; E-Kazakhstan Area: different localities.

CATALOGUE. Russia: BLG: Valuyki [Kulczynski, 1913\*]. Kazakhstan: EKA [Savelieva\*, 1972ab, 1979].

DISTRIBUTION. Steppe and forest-steppe from Czechoslovakia to E-Kazakhstan. New for Czechoslovakian fauna.

REMARK. Only one of two epigyne pictures of *C. germanica* by Miller [1971] actually belongs to *C. congentilis*. Unfortunately the locality remains unknown.

#### The *japonicola* - group.

DISTRIBUTION. Palaearctic, Indo-Malayan region, 2 species. Outside USSR - *C. yasudai* Ono, 1991 (Hokkaido).

#### *Clubiona (Clubiona) japonicola* Bösenberg et Strand, 1906.

CATALOGUE. Russia: AMR: Khingan State Reserve [Mikhailov, 1990a]. PRI: Khanka, Khasan distr. [Mikhailov, 1990a].

DISTRIBUTION. USSR - S-Far East. All Japan, S-Korea, N-Korea (original data; IZPAN), all China, Taiwan, Philippines.

#### The *obesa* - group.

DISTRIBUTION. Numerous species in E-Asia and N-America. Japan: *C.akagiensis* Hayashi, 1985, *C.chikunii* Hayashi, 1986 (subgroup *akagiensis*), *C.uenoi* Ono, 1986, *C.inaensis* Hayashi, 1989, *C.yoshidai* Hayashi, 1989. Japan & China: *C.corrugata* Bösenberg et Strand, 1906. China: *C.pbragnitoides* Schenkel, 1963, *C.fuzhouensis* Gong, 1985, *C.manshanensis* M.Zhu et An, 1988, *C.aciformis* Zhang et Hu, 1991.

#### *Clubiona (Clubiona) furcata* Emerton, 1919.

MATERIAL. Russia: 1♀, Yakutia, environs of Khandyga, 1.09.1990, IDS; 1♀ (BIN 94), 1♀, Chita Area, Kyra distr., Sokhondo State Reserve, confluence of Larionova Stream and Agutsa River, valley forest of *Picea*, *Betula*, *Larix*, ca. 1350 m, crowns of *Betula* and *Larix*, 14.06.1991, DVL.

CATALOGUE. Russia: MGD: Kolyma River upper flow [Mikhailov, 1990a, 1992; Marusik, 1992].

DISTRIBUTION. True boreal species. USSR - mountains of S-Siberia and N-Far East. N-America to Utah, USA.

#### *Clubiona (Clubiona) praematura* Emerton, 1909.

CATALOGUE. Russia: YaKT: Kolyma River lower flow [Mikhailov, 1990a]. MGD: Chukotka [Mikhailov, 1990a].

DISTRIBUTION. USSR - N-Far East. N-America to Maine, USA.

#### *Clubiona (Clubiona) kurilensis* Bösenberg et Strand, 1906.

MATERIAL. Russia: Amur Area: 1♂ 1♀, Tambovka distr., environs of Kozmodemyanova, meadow, litter, 06.1990, GNG; Primorye: 1♀ (ZIP), lower flow of Kedrovka River, swampy meadow, 20.06.1968, FZP.

CATALOGUE. Russia: AMR: Khingan State Reserve [Mikhailov, 1990a]. KhAB: Nanayskiy distr., Jewish AO, Khabarovsk, Bolshekhekhtsyrskiy State Reserve [Mikhailov, 1990a]. PRI: Khanka, Khasan distr. [Mikhailov, 1990a]. SAKh: S-Sakhalin [Saito, 1934; Mikhailov, 1990a], Kunashir [Mikhailov, 1990a], Iturup [Bösenberg, Strand, 1906; Strand, 1909], N Kurile Isles. [Saito, 1932, 1933].

DISTRIBUTION. USSR - S-Far East (both continental and island). All Japan, S-Korea, all China.

#### *Clubiona (Clubiona) logunovi* Michailov, 1990.

MATERIAL. Russia: 1♂, Primorye, Sikhote-Alin State Reserve, coniferous forest, 24.06.1984, M.N.Gromyko.

CATALOGUE. Russia: KhAB: Bolshekhekhtsyrskiy Reserve [Mikhailov, 1990a].

DISTRIBUTION. USSR - S-Far East (continental).

*Clubiona (Clubiona) marusiki* Michailov, 1990.

CATALOGUE. Russia: AMR: Arkhara [Mikhailov, 1990a].  
DISTRIBUTION. The single type locality.

*Clubiona (Clubiona) paiki* Michailov, 1991.

CATALOGUE. Russia: KhAB: Bolshevikheksyrskiy Reserve [Mikhailov, 1991].

DISTRIBUTION. The single type locality.

The *latericia*-subgroup.

DISTRIBUTION. Holarctic, 6 species. Outside USSR - *C.bryantae* Gertsch, 1941 (N-America).

*Clubiona (Clubiona) latericia* Kulczynski, 1926.

MATERIAL. Russia, Tuva: 3♂ 11♀, Tandinskij distr., E bank of Chagytay Lake, 1000 m, a swamp, 29.06.1989, DVL; Chita Area, Sokhondo State Reserve: 2♂, 3-5 km SW of Kyra, Kyra River floodplain, meadow, 850 m, 30.05.1991, DVL; 1♂ (BIN 93), id., 19.06.1991, DVL; 1♂ 1♀, Ingoda River floodplain, Ahus dwarf stand, 1350 m, 8-9.06.1991, DVL.

CATALOGUE. Russia: YaKT: Kolyma River lower flow [Mikhailov, 1990a]. MGD: Kolyma River upper flow [Mikhailov, 1990a]. KMCh [Kulczynski, 1926; Schenkel, 1930; Sytshevskaja, 1935\*; Mikhailov, 1990a]. KhAB: Bolshevikheksyrskiy Reserve [Mikhailov, 1990a]. PR: Kedrovaya Pad Reserve [Mikhailov, 1990a]. SAKh: S-Sakhalin [Mikhailov, 1990a].

DISTRIBUTION. USSR - S-, E-Siberia, all Far East. Alaska.

*Clubiona (Clubiona) tsurusakii* Hayashi, 1987.

MATERIAL. Russia, 1♂, S-Kuriles, Kunashir, Ivanovskiy Cape, 6-8.07.1989, V.K.Zinchenko.

DISTRIBUTION. New for the USSR fauna. All Japan.

*Clubiona (Clubiona) kumashirensis* Michailov, 1990.

1990a C.k. Mikhailov [sp.nov.]: 156-157, figs.33-37 (♂).

1991 *Clyubarskii* Mikhailov [sp.nov.]: 227-228, figs. 57-59 (♀), syn.n.

MATERIAL. Russia: 1♀, Iturup, environs of Kurilsk, 20-24.06.1989, AMB.

CATALOGUE. Russia: SAKh: Kunashir [Mikhailov, 1990a, 1991].

DISTRIBUTION. USSR - S-Kuriles.

*Clubiona (Clubiona) wolchongsensis* Paik, 1990.

MATERIAL. Russia, Khabarovsk Prov.: 1♀ (ZIP), environs of Vyatskoe, 26.06.1910, VKS; Primorye: 1♀ (ZIP), Kedrovaya Pad State Reserve, 17.08.1968, FZP.

CATALOGUE. Russia: KhAB: Jewish AO, environs of Khabarovsk [Mikhailov, 1991]. PRI: Bikin River, Khanka distr. [Mikhailov, 1991].

DISTRIBUTION. USSR - S-Far East (continental). S-Korea, N-Korea (original data; IZPAN), China: Liaonin.

*Clubiona (Clubiona) subborealis* Michailov, sp.n.

Figs. 5a-c, 6a-c.

MATERIAL. Holotype 1♂ (ZMUM Ta-4806), Chita Area, Kyra distr., Sokhondo State Reserve, Ernichniy Stream valley, *Betula*, *Larix* forest, 1400-1450m, 2-3.06.1991, DVL. Paratypes: Russia, Chita Area, Sokhondo State Reserve: 1♂ 2♀, as holotype; 1♂, confluence of Larionova Stream and Agutsa River, ca. 1350 m, *Betula*, *Larix* forest, 13.06.1991, DVL; 1♀, id., open grassland in a forest, 2♀ (ZMUM Ta-4807), 7-8 km SW of Buninda, *Larix*, *Betula* forest, ca. 1300 m, 14.06.1991, DVL; Khabarovsk Prov.: 1♂ 4♀, Komsomolsk distr., Komsomolsk State Reserve, mouth of Goryun River, *Larix* stand, 13-16.07.1991, GNG. Mongolia: 2♂ 1♀, S-Khangai Mts., ca. 80 km NE of Ulan-Bator, Tereldsh Kishr, S.Heimer.

DIAGNOSIS. By the palp structure the male is more close to both *C.logunovi* and *C.tsurusakii* differing by the solid, not subdivided ventral branch of the tibial apophysis (in *C.tsurusakii* the entire apophysis is solid). By the structure of the female genitalia *C.subborealis* differ the following character unique for *Clubiona*: double covering of the parts of the copulatory tube (viewed strongly ventrally) a little anteriorly of the copulatory openings, the broad and long copulatory tubes: their length exceeds the epigynal length twice, end parts of the copulatory tubes (near the spermathecae) the parallel to each other.

DESCRIPTION. MALE (6). Total length 4.68+0.58. Carapace length 2.12+0.23, width 1.62+0.14, their ratio 1.31+0.06. Carapace, chelicerae, and legs pale or straw-coloured. Leg measurements:

	I	II	III	IV
Femur	♂ 1.62+0.17	1.66+0.19	1.46+0.20	2.01+0.20
	♀ 1.61+0.21	1.59+0.28	1.48+0.22	2.09+0.28
Patella	♂ 0.83+0.09	0.83+0.09	0.68+0.06	0.82+0.09
	♀ 0.86+0.12	0.86+0.14	0.75+0.12	0.90+0.15
Tibia	♂ 1.50+0.15	1.51+0.16	1.06+0.19	1.63+0.14
	♀ 1.28+0.20	1.30+0.19	0.95+0.12	1.60+0.21
Metatarsus	♂ 1.10+0.15	1.11+0.12	1.26+0.10	2.13+0.18
	♀ 0.93+0.15	0.92+0.17	1.14+0.23	2.02+0.32
Tarsus	♂ 0.73+0.12	0.72+0.07	0.53+0.05	0.70+0.05
	♀ 0.63+0.09	0.64+0.11	0.53+0.07	0.67+0.07

Leg armature: femur I d.1.1.2, II d.1.1.2 (1.1.3), III-IV d.1.1.3, patella IV rl.1, tibia I-II v.2.2, III d.2.2, v.1.1 (1.1.1), IV d.2.2, v.1.1.1, metatarsus I-I v.2, III d.2.1.2, I.1.2 (2.2), v.2.2, IV d.2.1.2, I.2.2, v.2.1.2 (2.2).

Palp as in Fig.5a-c. Lengths of cymbium, tibia, patella, and femur are 0.82+0.06, 0.30+0.08,




Fig.5. *Clubiona (Clubiona) subborealis* sp.n., male palp: a) ventral view; b) extero-ventral view; c) dorsal view. Scale = 0.14.  
Рис.5. *Clubiona (Clubiona) subborealis* sp.n., пальпа самца: а) вентрально; б) наружно-вентрально; в) дорсально. Масштаб 0.14.

$0.33 \pm 0.05$ ,  $0.62 \pm 0.13$ , respectively.

Abdomen: length  $2.73 \pm 0.33$ , width  $1.42 \pm 0.27$ , their ratio  $1.94 \pm 0.37$ ; pale or pale brown.

FEMALE (10). Total length  $6.39 \pm 0.97$ . Carapace length  $2.31 \pm 0.39$ , width  $1.76 \pm 0.27$ , their ratio  $1.32 \pm 0.11$ . Carapace and legs pale, straw or reddish-coloured. Chelicerae reddish, brown or chestnut-coloured. Leg measurements see above. Leg armature as in male, but femur II d.1.1.2, tibia III d.2.2, v.1.1, metatarsus III d.2.1.2 (2.2), l.1.2, v.2.2, IV d.2.1.2, l.2.2, v.2.1.2.

Abdomen: length  $3.95 \pm 0.80$ , width  $2.31 \pm 0.42$ , their ratio  $1.71 \pm 0.17$ ; pale, pale brown or brown. Epigyne as in Fig.6a-c.

DISTRIBUTION AND ECOLOGY. Occurs in *Betula* and *Larix* forests in S-Siberia, more frequently in mountains (1300-1450 m in Sokhondo Reserve).

The *sapporensis* - subgroup.

DISTRIBUTION. E-Asia - 5 species. Out of USSR - *C.manshanensis*, *C.fuzhouensis* (see above).

*Clubiona (Clubiona) sapporensis* Hayashi, 1986.

MATERIAL. Russia, Amur Area, Arkhara distr., Khingan State

Reserve: 1♂ 1♀, Lebedinoe Lake, birch, alder, wild rosa, 2-3.08.1991, DKK; S-Kuriles, Iturup: 2♀, environs of Kurilsk, 20-24.06.1989, AMB.

CATALOGUE. Russia: PRI: Vladivostok, Khasan distr. [Mikhailov, 1990a]. SAKh: Sakhalin [Saito, 1934; Mikhailov, 1990a], Kunashir [Mikhailov, 1990a].

DISTRIBUTION. USSR - S-Far East (both continental and island). Japan: Hokkaido; N-Korea (original data; IZPAN).

*Clubiona (Clubiona) charitonovi* Mikhailov, 1990.

MATERIAL. Russia: 1♂ (MZH) "Siberia, gouv. Jeniseisk, Minussinsk, 1881, Roschier"; Irkutsk Area: 1♂ 2♀ (IZPAN, Kulczynski's collection F.1759), Kultuk; Chita Area: 1♂ (ZMUM), Nizhniy Tsasuchey, Bydykkhen Lake, small shrub, 26.06.1984, SND; Amur Area: 1♀ (ZIP), Gunyaev-Zeiskiy Trakt, Ulung Station, 18.05.1910 and later, Verkhovskaya. 1♂ 1♀ (ZIP), Dep River (left tributary of Zeya River), 07-08.1911, P.Rippas; Khabarovsk Prov.: 1♀ (ZIP) Sredne-Tambovskoe, 18.07.1910, VKS; 1♀ (ZIP) Kunashir, Otradnoe, 31.07.1965, Lukina.

CATALOGUE. Russia: BUR: Selenga distr. [Mikhailov, 1990a]. CHIT: Khilok distr. [Mikhailov, 1990a]. AMR: Khingan Reserve [Mikhailov, 1990a]. KhAB: Bolshekhekhtsyrskiy Reserve [Mikhailov, 1990a]. SAKh: S-Sakhalin, Kunashir [Mikhailov, 1990a].




Fig. 6. *Clubiona (Clubiona) subborealis* sp.n., female: a) epigyne; b) vulva; c) direction of copulatory tube, ventral view. Scale = 0.14.  
Рис. 6. *Clubiona (Clubiona) subborealis* sp.n., самка: а) эпигина; б) вульва; в) направление копуляторного канала, вентрально. Масштаб 0.14.

DISTRIBUTION. USSR - S-Siberia, S-Far East (both continental and island).

*Clubiona (Clubiona) microsapporensis* Michailov, 1990.

MATERIAL. Russia, Khabarovsk Prov.: 1♀ (ZIP), Amur, Schrenk (determined by E.Grube as *Camarantha* Walck.); 1♀ (Kh), Nanayskiy distr., Slavyanka, meadow, *Rosa canina*, 31.05.1990, DKK; Primorye: 1♀ (ZIP), lower flow of Kedrovka River, swampy meadow, 20.06.1968, FZP; 1♀ (ZIP), Kedrovka River midstream, 3.08.1968, FZP; 1♀ (ZIP), Slavyanka, 1911, I.I.Kuznetsov.

CATALOGUE. Russia: AMR: Khingan Reserve [Mikhailov, 1990a].

DISTRIBUTION. USSR - S-Far East (continental).

The *corrugata* - subgroup.

DISTRIBUTION. E-Asia - 3 species. Outside USSR - *C.corrugata* and *C.phragmitoides* (see above).

*Clubiona (Clubiona) kimyongkii* Paik, 1990.

MATERIAL. Russia, Amur Area, Khingan State Reserve: 1♀, Karapcha River, 12 km S of Kundur, *Quercus* forest, litter, 08.1991, GNG; 1♀, Karapcha River, 10.08.1991, DKK; Khabarovsk Prov.: 1♀ (Kh), Nanayskiy distr., Slavyanka, 30.05.1990, DKK; 1♂ 1♀, id, 24.08.1991, DKK; 1♀, Bikin distr., Shivki, coniferous-broad-leaved forest, litter, 08.1990, GNG; Primorye: 3♂ 5♀ (ZIP), Kedrovka River, Kedrovaya Pad State Reserve and its environs, 30.05-15.06.1968, FZP; 1♂ 2♀, Popova Island, Alexeeva Bay, 31.08-8.09.1991, KGM.

CATALOGUE. Russia: KhAB: Bolshekhekhtsyrskiy Reserve [Mikhailov, 1990a]. PRI: Chuguevka, Ussuriyskiy Reserve, Khan-kha distr. [Mikhailov, 1990a].

DISTRIBUTION. USSR - S-Far East (continental).

The *bakurovi* - subgroup.

Includes a single species,

*Clubiona (Clubiona) bakurovi* Michailov, 1990.

MATERIAL. Russia: 1♀ (Kh) Khabarovsk Prov., Khekhtsyrs Mts., 24th km, *Pinus sibirica* - broadleaved forest, litter, 17.05-14.06.1990, DKK; 1♀ (ZIP) Primorye, Kangauz, upstream of Smolyanoy Klyuch, coniferous - broadleaved forest, on grass, 13.07.1968, FZP.

CATALOGUE. Russia: KhAB: Nanayskiy distr., Bolshekhekhtsyrskiy Reserve [Mikhailov, 1990a]. PRI: Chuguevka distr., Chernyshevka [Mikhailov, 1990a]. SAKh: Nituy River [Mikhailov, 1990a].

DISTRIBUTION. USSR - S-Far East (both continental and island). N-Korea (original data; IZPAN).

The *irinae* - subgroup.

Includes a single species,

*Clubiona (Clubiona) irinae* Michailov, 1991.

MATERIAL. Russia, Khabarovsk Prov.: 1♂ (PSU), Kom-somolsk State Reserve, 27.09.1987, Sheverdan; 1♀, id, mouth of Goryun River, *Larix* stand, 13-16.07.1991, GNG; Primorye: 2♀ (IP), Terney Bay, Sakhobe River, 08.1906, N.Palchevskiy; 1♀ (ZIP), Vladivostok, Orlinoe Gnezdo, 04-06.1903, N.Palchevskiy.

CATALOGUE. Russia: AMR: Zeyskiy Reserve, Khingan Reserve [Mikhailov, 1991]. KhAB: Jewish AO, Nanayskiy distr., Bikin distr., Khabarovsk, Khor, Bolshekhekhtsyrskiy Reserve [Mikhailov, 1991]. PRI: Bikin River, Ussuriyskiy Reserve, Kedrovaya Pad Reserve [Mikhailov, 1991]. SAKh: Sakhalin [Mikhailov, 1991].

DISTRIBUTION. USSR - S-Far East (both continental and island).

The *pallidula* - group.

DISTRIBUTION. Holarctic, 8-10 species. *C.tateyamensis* Ono, 1989 and *C.zacharovi* Michailov,

1991 are included here provisionally. Outside USSR - *C.mandschurica* Schenkel, 1953 (sensu Paik, 1990a).

*Clubiona (Clubiona) pallidula* (Clerck, 1757).

*Aranea holosericea* L. - Dwigubsky, 1802; Fischer, 1791.

*Clubiona holosericea* Walck. - Belke, 1853, 1859.

*Camarantha* Koch - Siemaschko, 1861.

*C. holosericea* L. - Belke, 1866; Palmgren, 1943; Spassky, 1927; Vilbaste, 1974; Zhukauskene, 1966.

*C. holosericea* Latr. - Vagner, 1895.

MATERIAL. Russia, Ulyanovsk Area: 1♂ (ZIP), Radishchevo distr., Volga River floodplain, 19.05.1961; Samara Area: 14♂, Zhigulevsk, Yablonevyi Ovrag, 15.06.1992, PMD; Perm Area: different localities, det. SLE; Kurgan Area: 2♂ 1♀, Uval, Tobol River floodplain, 8.06-16.07.1989, NAU; Novosibirsk Area, Novobisibirk distr., Koltsovo: 1♂, *Betula* forest, 8.06.1986, leg. et det. DVL; 1♂ 2♀ (BIN), id, 29.05.1988, leg. et det. DVL; Tuva: 1♀, Kurtushinskiy Mt.Ridge, ca. 10 km NW of Shivilig, 1100-1200 m, *Betula* crowns, 5.07.1990, DVL; 1♂, Shivilig-Khem River valley, poplar-larch forest, 30.05.1990, OVL; 1♂, Erzin distr., Tes-Khem River, 20 km W of Erzin, poplar floodplain forest, 800 m, 31.05.1989, DVL; 1♀, Azas State Reserve, 950 m, *Betula* forest, 1.06.1990, ABR; Buryatia: 3♂ 3♀ (UU), Kabansk distr., Beregovaya, small-leaved forest, *Betula*, 11.06.1983, leg. et det. SND; 6♂ 9♀ (UU), Kabansk distr., Boyarsk, small-leaved forest, 22.06.1983, leg. et det. SND; Chita Area: 1♀ (UU), Mogotuy distr., environs of Tsugol, *Betula* forest, 8.07.1982, leg. et det. SND; Sakhalin: 1♀, Yuzhno-Sakhalinsk, 1.07.1989, AMB. Kazakhstan: 1♀ (OL), Pavlodar Area, Mayskiy distr., Irtysh River floodplain, 15.08.1989, OVL.

CATALOGUE. Russia: KRL [Palmgren, 1943]. LNN [Siemaschko, 1861; Kharitonov, 1928\*; Palmgren, 1943]. KLN: Ostashkov [Fedotov, 1912c]. KLZh [Vagner, 1895: C.p.+ C.b.]. YaRS [L.Becker, 1879; Fedotov, 1915]. YaRS + VOL: Darvinskiy Reserve [Alexeev, 1986]. MSK [Dwigubsky, 1802; Vagner, 1892; Freiberg, 1894\*, 1897\*; Pokrovskiy, 1925\*; Pereleshina, 1928a\*; Mikhailov, 1983ab]. BRN [Kharitonov, 1936\*]. KUR [Pichka, 1965, 1984ab]. LPTs: Galichya Gora State Reserve [Panteleeva, 1982]. BLG: Valuyki [Kulczynski, 1913], "Les na Vorskle" State Reserve [Ovsyannikov, 1937; Marusik, 1984]. VRN [Pichka, 1965, 1967; Pichka, Skufin, 1981]. GRK [Greze, 1915a]. ChVSh [Holzmayer, 1934]. PNZ [Pyatin, 1988]. TTR [Azheganova, Glukhov, 1973; Garkusha, 1977]. RST [Spassky\*, 1912, 1925, 1940; Sorokin, 1977]. KRS: Krasnodar [Sharov et al., 1984; Mikhailov, 1990b], Slavyanka distr. [Mikhailov, 1990b]. STV: environs of Pyatigorsk [Mikhailov, 1990b]. SOA [Mikhailov, 1990b]. ChLB: Troitskiy Reserve [Azheganova, 1951; Utochkin, Golovashkina, 1977], Ilmen State Reserve [Polyanin, Pakhorukov, 1988\*]. TMN: Tura plain [Shlykov, 1975], Tobol floodplain [Shlykov, 1978]. OMS [Spassky, Lavrov, 1928]. TMS [Ermolajev, 1934]. KRN: Boguchany distr. [Izmailova, Verzhutskiy, 1981], Chunoyar [Izmailova, 1989]. KhAB: Jewish AO [Mikhailov, 1991]. Estonia [Vilbaste, 1960, 1972, 1974, 1980, 1987]. "Lifland" [Fischer, 1791]. Latvia [Prieditis, Sternbergs, 1981, Sternbergs, 1981a, 1983. Lithuania [Zhukauskene, 1966]. Byelorussia [Litvinova et al., 1980\*]. Byelovezhskaya Pushcha [Chebotareva, 1987]. MIN [Zhukovets, 1989]. VIT [Pereleshina, 1930\*]. Ukraine: TRN [Gnelitsa, 1989]. ZAK [Polozhentsev, Akimtseva, 1980]. Pripyatskoe Polesye [Polishchuk et al., 1980]. ChRK: Kanev State Reserve [Pichka, 1974]. KhRK [Kirilenko, Legotay,

GOM: Rogachev [Pereleshina, 1928b\*]. Ukraine: LVV [Wajgiel, 1868, 1874; Nowicki, 1874]. LVV + IFR [Bublik, 1981]. ZAK [Legotay, 1958, 1980; Polozhentsev, Akimtseva, 1980]. ZhTM: Radomyshl [Belke, 1866; Thorell, 1875-cit]. KhML: Kamenets-Podolskiy [Belke, 1853, 1859; Thorell, 1875-cit]. ChRK: Kanev State Reserve [Pichka, 1974]. KhRK [Kirilenko, Legotay, 1981]. DNP: "Ekaterinoslav" [Thorell, 1875]. ODS: Odessa [Sharov et al., 1984\*]. KhRS: Askania-Nova State Reserve [Pereleshina, 1927], Chernomorsky State Reserve [Polchaninova, 1988], lower flow of Dnepr River [Talanov, Nazarenko, 1989]. VRSh [Polchaninova, 1992]. Krim [Spassky, 1927; Tolstov, Atanova, 1982]. Moldavia [Talitskiy, Legotay, 1982; Matsyuk, Yanyuk, Verlan, 1984]. Georgia [Mkheidze, 1964; Mikhailov, 1990b]: Batumi [Simon, 1899]. Armenia: Kirovakan [Mikhailov, 1990b]. Azerbaijan [Atakishiev, 1969; Dunin, 1989; Mikhailov, 1990b]. Kazakhstan: EKA [Savelieva\*, 1970, 1979].


DISTRIBUTION. USSR - European part, Caucasus, S of W- and C-Siberia, S-Siberia, S-Far East (except of Primorye). All Europe, N-America: British Columbia, Ontario (Canada), Washington (USA).

*Clubiona (Clubiona) pbragmitis* C.L.Koch, 1843.

*C.terrestris* - Spassky, 1927.

MATERIAL. Russia, Pskov Area: 1♂ (ZIP), Novorzhev distr., near Moshkovo and Filino, summer 1916, Beletskiy, det. V.E.Pereleshina; Arkhangelsk Area: 3♂ 5♀, Kozhozero, 20-25.08.1984, DVL; Tambov Area: 1♀ (MGPU), Muchkap, 25.07.1987, S.V.Makarenkov; Tuva: 7♂ 4♀, Erzin distr., Terkhon Lake, a swamp, 29.05.1989, DVL. Ukraine: some specimens (ZIP) environs of Berdyansk, 1.03-15.05.1938, leg. V.Nikolaev, det.SAS. Kazakhstan: 1♀ (ZIP), Aral Sea, 10.1936, leg.Karpova, det.SAS. Kirghizia: 1♀, Ferganskiy Mt.Ridge, Baubashata Mt.Ridge, environs of Arslanbob, Yarodar, a lake, in *Phragmites* stems, 10.05.1983, SLZ.

CATALOGUE. Russia: KRL [Palmgren, 1943; Uzenbaev, 1986]. LNN [Palmgren, 1943]. PSK [Pereleshina, 1930\*]. KLN: Bologoe [Kharitonov, 1936\*], Belskiy Uezd [Pokrovskiy, 1930], Ostashkov [Fedotov, 1912c]. YaRS + VOL: Darvinskiy State Reserve [Alexeev, 1986]. MSK [Vagner, 1892; Greze, 1915b\*; Pereleshina, 1928a\*; Mikhailov, 1983b]. KUR [Pichka, 1965, 1984b]. BLG: Valuyki [Kulczynski, 1913], "Les na Vorskle" State Reserve [Marusik, 1984]. TTR [Azheganova, Glukhov, 1973; Utochkin, 1977]. KBSh: Zhiguli State Reserve [Krasnobaev, Ovtsharenko, 1986]. AST: Astrakhan State Reserve [Utochkin, 1971]. RST [Spassky\*, 1912, 1925, 1940]. KLM [Minoranskiy, Ponomarev, 1984]. KRS: Slavyanka distr. [Mikhailov, 1990b]. ChIA [Minoranskiy, Ponomarev, Slyusarev, Gramotenko, 1984; Minoranskiy, 1988]. BShK: Bashkirskiy State Reserve [Pakhorukov, Efimik, 1988]. ChLB: Troitskiy Reserve [Azheganova, 1951]. Estonia [Vilbaste, 1957, 1972, 1974, 1980, 1987]. Latvia [Prieditis, Sternbergs, 1981; Sternbergs, 1981a, 1983]. Lithuania [Vaitskute, 1960, 1963; Zhukauskene, 1968]. Byelorussia [Litvinova et al., 1980]. Belovezhskaya Pushcha State Reserve [Chebotareva, 1987]. MIN [Zhukovets, 1989]. VIT [Pereleshina, 1930\*]. Ukraine: TRN [Gnelitsa, 1989]. ZAK [Polozhentsev, Akimtseva, 1980]. Pripyatskoe Polesye [Polishchuk et al., 1980]. ChRK: Kanev State Reserve [Pichka, 1974]. KhRK [Kirilenko, Legotay,

Fig.7. *Clubiona (Clubiona) komissarovi* sp.n., female: a) epigyne; b) vulva. Scale = 0.14.Рис.7. *Clubiona (Clubiona) komissarovi* sp.n., самка: а) эпигина; б) вульва. Масштаб 0.14.

1981; Polchaninova, Astakhova, 1984]. Levoberezhnaya Ukraina: N steppes, SE forest-steppe [Polchaninova, 1984]. KhRS: "Dneprivskiy Uezd" [Spassky, 1927: C.t.\*]. Moldavia [Talitskiy, Legotay, 1982]. Azerbaidjan [Atakishiev, 1969; Dunin, 1989; Mikhailov, 1990b]. "Turanian zoogeographical province" [Spassky, 1952\*]. Kazakhstan: EKA [Savelieva\*, 1970, 1979].

Wrong identifications. Russia: AMR [Azheganova, Stenchenko, 1977]. PRI [Sternbergs, 1988]. SAKh: S-Sakhalin [Saito, 1935].

DISTRIBUTION. USSR - European part, Caucasus, S of W- and C-Siberia, montaneous Middle Asia. S-, C-, N-Europe.

#### *Clubiona (Clubiona) ezoensis* Hayashi, 1987.

MATERIAL. Russia: 1♀ (ZIP 127/1072) "near Kinzi" leg. Shrenk; 1♀ (ZIP), Sakhalin, Due, Ugolnaya Pad, 21.06 (1920th), B.Dombrovskiy.

CATALOGUE. Russia: AMR: Khingan State Reserve [Mikhailov, 1991]. SAKh: Sakhalin, Kunashir [Mikhailov, 1991].

DISTRIBUTION. USSR - S-Far East (both continental and island). Japan: Hokkaido.

#### *Clubiona (Clubiona) langei* Michailov, 1991.

MATERIAL. Russia, Khabarovsk Prov.: 2♂ (ZIP), environs of Troitskoe, 30.06-2.07.1910, VKS; Primorye: 1♀ (ZIP), Vinogradovka, 6.07.1929, A.N.Kirichenko; 1♀ (ZIP), Shkotovo distr., upper flow of Smolyanoy stream, coniferous-broadleaved forest, 13.07.1968, FZP; 1♀ (ZIP), environs of Kangauz (Anisimovka), 5.07.1968, FZP.

CATALOGUE. Russia: KhAB: Bolshevikheksyrskiy State Reserve [Mikhailov, 1991]. PRI: Kedrovaya Pad State Reserve [Mikhailov, 1991].

DISTRIBUTION. USSR - S-Far East (continental).

#### *Clubiona (Clubiona) zucharovi* Michailov, 1991.

MATERIAL. Russia, Primorye: 3♂ (ZIP), Kangauz (Anisimovka), 5.07.1968, FZP.

CATALOGUE. Russia: PRI: Kedrovaya Pad State Reserve [Mikhailov, 1991].

DISTRIBUTION. USSR - USSR - S-Far East (continental).

#### *Clubiona (Clubiona) komissarovi* Michailov, sp.n.

Fig. 7a,b.

MATERIAL. 1♀ (holotype, ZMUM Ta-4808), Russia, Primorye, Popova Isl., Alexeeva Bay, SW slope 0-50 m, broadleaved forest with Siberian pine, litter, 3.09.1991, KGM.

DIAGNOSIS. By the structure of genitalia the female differs from the other members of the *pallidula*-group by the narrow copulatory openings, which are twice less than the breadth of the copulatory tubes. The combination of following characters is of equal importance: the atrium and the second (mesal) part of the spermatheca are of equal breadth and length; the copulatory opening are situated at the level (vertical by viewing ventrally) of the first (lateral) part of the spermathecae (not at the level of the atria, as in *C.sopaiensis*); posterior part of the epigyne exceeds a little behind the epigastric fissure.

DESCRIPTION. FEMALE. Total length 6.1. Carapace length 2.85, width 1.90, their ratio 1.50. Carapace and legs straw-coloured, chelicerae brown. Leg measurements:

	I	II	III	IV
Femur	1.93	1.95	1.63	2.55
Patella	1.03	1.05	0.83	1.00
Tibia	1.63	1.68	1.25	2.00

Metatarsus	1.18	1.23	1.45	2.48
Tarsus	0.73	0.75	0.63	0.75

Leg armature: F I-II d.1.1.2, III d.1.2.3 (1.3.3), IV d.1.1.3, Pt III-IV rl.1, Ti I-II v.2.2, III d.2.2, v.1.1, IV d.2.2, v.1.1.1, Mt I-II v.2, III d.2.1.2, I.1.2, v.2.2, IV d.2.1.2, I.2.2, v.2.1.2.

Abdomen: length 3.40, width 1.95, their ratio 1.74; dark brown. Epigyne as in Fig. 7a,b.

MALE unknown.

DISTRIBUTION AND ECOLOGY. The species is known from the single locality at the small island, ca. 20 km from continent. Apparently it is associated with broadleaved forests.

DERIVATIO NOMINIS. The species is dedicated to my companion during the short field trip in the Far East in 1991, molecular geneticist Andrey Komissarov (Moscow).

#### *Clubiona (Clubiona) sopaikensis* Paik, 1990.

MATERIAL. Russia, Amur Area: 1♀, Arkhara distr., Khingan State Reserve, Lebedinoe Lake, birch, alder, wild rosa crowns, 2.08.1991, DKK; Primorye: 1♀ (ZIP), Khanka Lake, environs of Kamen-Rybolov, 22.07.1968, FZP; 2♀ (ZIP), Shkotovo distr., environs of Kangauz (Anisimovka), 28.06-6.07.1968, FZP.

CATALOGUE. Russia: AMR: Kundur, Sennoy Isls. [Mikhailov, 1991]. KhAB: Jewish AO, Khabarovsk, Khor [Mikhailov, 1991]. PRI: Khorol, Kedrovaya Pad State Reserve, Khasan distr. [Mikhailov, 1991].

DISTRIBUTION. USSR - S-Far East (both continental and island). S-Korea, N-Korea (original data; IZPAN).

#### *Clubiona (Clubiona) odesanensis* Paik, 1990.

MATERIAL. Russia, Khabarovsk Prov.: 1♂, Nanayskiy distr., Slavyanka, broadleaved forest, 2.08.1990, DKK; 1♀, Khabarovsk, Voronezhskie Sopki hills, broadleaved forest, crowns, 4.07.1991, DKK; Primorye: 1♂ (ZIP), Kangauz (Anisimovka), 10.07.1968, FZP; 1♂ (ZIP), midstream of Kedrovka River, 16.06.1968, FZP.

CATALOGUE. Russia: PRI: Ussuriyskiy State Reserve [Mikhailov, 1991].

DISTRIBUTION. USSR - S-Far East (continental). S-Korea, N-Korea (original data; IZPAN).

#### *Clubiona (Clubiona) propinqua* L.Koch, 1879.

1907 *C. subinteriecta* Strand: 562 (♂, non ♀), syn.n.

1909 *C. subinteriecta* - Strand: 35-38 (♂, non ♀).

1991 *Clubiona propinqua* - Mikhailov: 215-217, Figs.30-33 (♂, ♀).

1991 *C. subinteriecta* - Mikhailov: 217-218, Figs.34-38 (♂, ♀).

MATERIAL. Russia, Krasnoyarsk Prov.: 2♀ (BIN 60), 1♂ 2♀, Ermakovo distr., 25-30 km N of Aradan, 1600 m, stony stream, 9.07.1990, DVL; Irkutsk Area: 1♀ (ZIP), Marituy, 25.06.1925, P.V.Tikhomirov; 1♂ 2♀, environs of Slyudyanka, Shamanka Cape,

*Betula* forest, in leaves, 24.07.1990, KGM; Chita Area, Kyra distr., Sokhondo State Reserve: 1♂ 1♀, confluence of Larionova Stream and Agutsa River, ca. 1300 m, moss-grown stony stream, 13.06.1991, DVL; 1♀ (BIN 91), environs of Vershina Bukukuna mine, 1600-1700 m, upper *Larix* forest line, 19-23.06.1991, V.V.Dubatolov; 1♂ (BIN 96), id., 1600 m, stony stream in a forest, 3.06.1991, DVL; 1♀, id., alpine tundra, 1700 m, 28-29.06.1991, S.E.Chernyshev; 1♂ 6♀, id., alpine *Driada*-stony tundra, 1700-1750 m, 1.06.1991, DVL; Amur Area: 1♀ (ZIP), Dep River (left tributary of Zeya River), 07-08.1911, P.Rippas; Primorye: 1♂ 8♀ (ZIP), Shkotovo distr., environs of Kangauz (Anisimovka), coniferous-broadleaved forest, in grass and under stones along a stream, 13.07.1968, FZP; 4♀ (ZIP), id., broadleaved forest, 30.06-5.07.1968, FZP.

CATALOGUE. Russia: KRN: Krasnoyarsk [Koch, 1879; Holm, 1973]. YaKT: Khametakh Lake, 232 km of Khandyga-Magadan highway [Mikhailov, 1991]. MGD: environs of Magadan, upper flow of Kolyma River, Chukotka [Mikhailov, 1991]. KMCh [Sytsevskaja, 1935\*: Mikhailov, 1991]. BUR: Visim River [Mikhailov, 1991]. AMR: Blagoveschensk [Strand\*, 1907, 1909; Mikhailov, 1991], Kundur [Mikhailov, 1991]. KhAB: Bolshekheksyrskiy State Reserve [Mikhailov, 1991]. PRI: Ussuriyskiy State Reserve, Kedrovaya Pad State Reserve [Mikhailov, 1991]. SAKh: S-Sakhalin [Mikhailov, 1991].

DISTRIBUTION. USSR - Siberia E of Yenisey, Far East. N-Korea (original data; IZPAN).

REMARK. As it was noticed by me [Mikhailov, 1991, p.217-218], females of *C.propinqua* and "*C.subinteriecta*" have no differences in epigynal structure. The differences between the males mentioned in my paper ("more rounded embolus" in "*C.subinteriecta*") appear to be wrong, since all the emboli examined by me later were of *propinqua*-type (see Fig.31 in Mikhailov, 1991); changes of their "roundness" are connected with the different viewpoints (not strictly ventral view etc.). So I consider here *C.subinteriecta* as a junior synonym of *C.propinqua*.

#### *Clubiona (Clubiona) pseudogermanica* Schenkel, 1937.

MATERIAL. Russia: 1♂ (ZIP), Primorye, upper flow of Kedrovka River, 7.06.1968, FZP; 1♂, N-Kuriles, Shumshu Isl., Bolshoe Lake, 18-22.08.1989, AMB.

CATALOGUE. Russia: KhAB: Khabarovsk, Bolshekheksyrskiy State Reserve [Mikhailov, 1991]. PRI: Chuguevka distr., Anuchino distr., Ussuriyskiy State Reserve [Mikhailov, 1991]. SAKh: Sakhalin, Kunashir [Mikhailov, 1991].

DISTRIBUTION. USSR - S-Far East (continental), S-, C-Far East (island). Japan (Hokkaido), S-Korea, China (Liaonin, Hunan, Kansu).

#### The *reclusa* - group.

DISTRIBUTION. Holarctic, 13 species. Outside USSR - *C.rosserae* Locket et Millidge, 1953, and 2 American species [Dondale, Redner, 1982].

*Clubiona (Clubiona) reclusa* O.Pickard-Cambridge, 1863.

*Cubicunda* C.L.Koch - Siemaschko, 1861.

*Cgrisea* L.Koch - Bergroth, 1881; Greze, 1909b, 1911.

MATERIAL. Russia, Arkhangelsk Area: 1♀, Kozhozero, 25.08.1984, DVL; Kaluga Area: 2♀, Yukhnov, 4.08.1981, KGM; Krasnoyarsk Prov., Khakassia: 1♀, Askiz distr., ca. 8 km E of Birikchul, floodplain *Salix-Betula* forest, 1000 m, 16.07.1990, DVL; Buryatia: 1♀ (ZIP), Baikal Lake, near Barguzin, Chivyrkuy Gulf, 14.06.1917, L.L.Rossolimo et al. Lithuania: 1♀, Shilutee, Vente Ornithological Station, 1984, RM.

CATALOGUE. Russia: KRL [Palmgren, 1943; Uzenbaev, 1986]. LNN [Siemaschko, 1861; Palmgren, 1943]. KLN: Ostashkov [Fedotov, 1912b], Belskiy Uezd [Pokrovskiy, 1930]. MSK [Greze, 1911\*, 1914; Pereleshina, 1928a\*; Mikhailov, 1983ab]. BRN [Kharitonov, 1936\*]. VRN [Pichka, 1965, 1967, 1977]. TTR [Utochkin, 1977]: Kazan [Kulczynski, 1901]. KOM: Pechoro-Ilychskiy State Reserve [Pakhorukov, 1980]. KST [Mikhailov, 1989]. PRM: Basegi State Reserve [Esyunin, 1991]. SVR [Pakhorukov, 1977]. BShK: Bashkirskiy State Reserve [Pakhorukov, Efimik, 1988]. TMN: S-Yamal [Greze, 1909b; Bakhvalov, Korshunov, 1976-cit; Eskov, 1986\*], between Tobolsk and Obdorsk (now Salekhard) [Bergroth, 1881], Tobolsk [Ermolaev, Samko, 1929\*; Ermolajev, 1937]. KRN: Nizhnyaya Tunguska River [Holm, 1973], Mirnoe [Eskov\*, 1986, 1988], Evenkiya [Eskov, 1988\*]. IRK [Izmailova, 1989]. BUR/ChIT: Muy Kettle [Verzhutskiy, Bessolitsyna, Seryshev, 1985]. Estonia [Vilbaste, 1964, 1972, 1973, 1974, 1980, 1987]. Latvia [Prieditis, Sternbergs, 1981; Sternbergs, 1981a, 1983]. Byelorussia [Litvinova et al., 1980]. MIN [Zhukovets, 1989]. VIT [Pereleshina, 1930\*]. Ukraine: ZAK [Legotay, 1958, 1989]. ChRK: Kanev [Pichka, 1974]. KhRK [Kirilenko, Legotay, 1981]. Kazakhstan: EKA [Savelieva, 1970, 1979].

Wrong identification. Russia, AMR [Azheganova, Stenchenko, 1977].

DISTRIBUTION. USSR - European part, Siberia to Yenisey and Baikal Lake. S-Europe: Balkans, Italy, C-, N-Europe.

#### *Clubiona (Clubiona) subsultans* Thorell, 1875.

Troitskiy Reserve [Azheganova, 1951], Ilmen State Reserve [Pakhorukov, Polyanin, 1987; Polyanin, Pakhorukov, 1988]. TMN: S-Yamal [Kulczynski, 1916], Tobolsk [Ermolaev, Samko, 1929; Ermolajev, 1937: *stagnalis*]. TMS [Ermolaev, 1934: *stagnalis*]. KRN: Stolby State Reserve [Sternbergs, 1977], Mirnoe [Eskov, 1988\*]. BUR [Odenvall, 1901; Izmailova, 1989]. BUR + ChIT: Muy Kettle [Verzhutskiy, Bessolitsyna, Seryshev, 1985]. Estonia [Vilbaste, 1957, 1959, 1964, 1969, 1972, 1973, 1980, 1987]. Latvia [Sternbergs, 1981a]. Lithuania [Zhukauskene, 1968]. Byelorussia: MIN [Zhukovets, 1989]. Ukraine: TRN [Gnelitsa, 1989]. ZAK [Legotay, 1989]. KhRK [Kirilenko, Legotay, 1981]. Georgia: Kolkhida [Mikhailov, 1990b].

DISTRIBUTION. USSR - European part, Caucasus, Siberia to Yenisey. S-Europe (Italy), C-, N-Europe, Afghanistan.

#### *Clubiona (Clubiona) subsultans* Thorell, 1875.

*Cerratica* C.Koch - L.Becker, 1879; Freiberg, 1894; Koch, 1879; Nordmann, 1863; Simon, 1891; Strand, 1906; Wagner, 1892, 1895.

#### *C.decora* - Greze, 1911.

MATERIAL. Russia, Arkhangelsk Area: 1♀, Kozhozero, Letniy Konets, 22.08.1984, DVL; 1♀, Gribanikha, *Picea* forest, 18.07.1972; Krasnodar Prov., Caucasian State Reserve: 1♂ 1♀ (ZIP), S slope of Chugush Mt., 27-29.06.1975, leg. et det. V.I.Ovtsharenko; Novosibirsk Area: different localities; Krasnoyarsk Prov.: 1♀, Evenkiya, Podkamenaya Tunguska River, ca. 70 km upstream of the mouth, Sulomay, 4.10.1989, ABR; Tuva: 2♂ 1♀ (BIN 50), 2♂ 1♀, Tandinskij distr., 1-5 km SW of Khovy-Aksy, Elegest River floodplain, forest of *Picea*, *Betula*, *Salix*, 5-6.05.1990, DVL; Chita Area, Kyra distr., Sokhondo State Reserve: 2♂ 4♀, confluence of Ubur-Ashagley Stream and Ingoda River, 1300-1400 m, *Larix*, *Betula* crowns, 8-9.06.1991, DVL; 1♂, confluense of Larionova stream and Agutsa River, ca. 1300 m, moss-grown stony stream, 13.06.1991, DVL.

CATALOGUE. Russia: MRM [Beer, 1964]. KRL [Palmgren, 1943; Uzenbaev, 1986]. LNN [Kharitonov, 1928\*; Palmgren, 1943]. KLN: Belskiy Uezd [Pokrovskiy, 1930]. KLZh [Vagner, 1895]. YaRS: Yaroslavl [L.Becker, 1879; Fedotov, 1915]. YaRS + VOL: Darvinskiy State Reserve [Alexeev, 1986]. MSK [Vagner, 1892; Freiberg, 1894; Greze, 1911\*: C.d., 1915b; Pereleshina, 1928a\*; Beer, 1968; Mikhailov, 1983b]. BRN [Kharitonov, 1936\*]. VRN [Pichka, 1965, 1967]. GRK [Greze, 1915a]. ChVSh [Holzmayer, 1934]. TTR [Garkusha, 1977; Utochkin, 1977]. KOM: Pechoro-Ilychskiy State Reserve [Pakhorukov, 1980]. KST [Veselova, 1986\*; Veselova, Mikhailov, 1986, 1987; Mikhailov, 1989]. PRM [Kharitonov, 1926a; Kharitonov, Bezzrezeva, 1970; Azheganova, Glukhov, 1981; Gridina, 1992]. Basegi State Reserve [Esyunin, 1991]. SVR [Kharitonov, 1923]. ChLB: Ilmen State Reserve [Pakhorukov, Polyanin, 1987; Polyanin, Pakhorukov, 1988]. TMN: Sosva River valley to Berezov [Simon, 1891; Strand, 1906-cit]. KRN: Krasnoyarsk, Intsarovo [L.Koch, 1879; Holm, 1973], Selivaninsk [Holm, 1973], Boguchany distr. [Izmailova, Verzhutskiy, 1981\*]. IRK [Izmailova, 1989]. BUR [Izmailova, 1989]. Estonia [Silvere, 1960; Vilbaste, 1969, 1972, 1973, 1980, 1987]. Latvia [Sternbergs, 1981a, 1985]. Lithuania [Zhukauskene, 1968]. Byelorussia [Litvinova et

#### *Clubiona (Clubiona) stagnatilis* Kulczynski, 1897.

MATERIAL. Russia, Murmansk Area: 4♂ 2♀, Kandalaksha State Reserve, Luda Malyshka Isl., 21.07, 14.08.1975, Yu.B.Byzova; 1♂, Vereskovaya Luda Isl., 27.07.1976, Yu.B.Byzova; Ryazan Area: 1♀, Okskiy State Reserve, Lubyaniki, 16.06.1981, KYE; Kurgan Area: 1♀, Uval, Tobol River floodplain, 8.06.1989, NAU; Tuva: 2♀, Piy-Khem distr., ca. 5 km NW of Sesarlig, *Larix*, *Salix* forest along a stream, 1100-1200 m, 24.07.1989, DVL; Novosibirsk Area: different localities. Ukraine: 1♀, Rovno, near a stream, on *Phragmites communis*, 4.08.1981, leg. et det. YMM. Kazakhstan: East-Kazakhstan Area: 4♀, Kaindinskiy Bor, Lake Shybyndy-Kul, 12-14.06.1968, LGS; 1♀, Feklistivka, 19.06.1969, LGS. Kirghizia: 2♂, 2 km N of Frunze, Karagachevaya Roshcha, 17.06.1979, SLZ.

CATALOGUE. Russia: KRL [Palmgren, 1943; Uzenbaev, 1986]. LNN [Palmgren, 1943]. MSK [Greze, 1911\*; Pereleshina, 1928a\*; Mikhailov, 1983ab]. ChVSh [Holzmayer, 1934]. STV: Teberda State Reserve [Mikhailov, 1990b]. KST [Veselova, 1986\*; Veselova, Mikhailov, 1986, 1987; Mikhailov, 1989]. PRM [Kharitonov, 1926a; Azheganova, Kopyltsova, Parfenova, 1976; Azheganova, Glukhov, 1981]. SVR [Pakhorukov, 1977]. ChLB:

al., 1980\*]. Belovezhskaya Pushcha [Chebotareva, 1987]. MIN [Zhukovets, 1989]. VIT [Pereleshina, 1930\*]. Ukraine: ZAK [Vysotskaya, 1977]. KVS [Pereleshina, 1930\*]. Levoberezhye [Astakhova, 1978].

Doubtful identifications. Ukraine: KhRS: Dnepr river lower flow [Talanov, Nazarenko, 1989]. Krim [Nordmann, 1863]. Kazakhstan: AAO [Tarabaev, 1979].

DISTRIBUTION. USSR - European part, Caucasus, Siberia to Yenisey and Transbaikalia. S-Europe: Italy, C-, N-Europe.

*Clubiona (Clubiona) norvegica* Strand, 1900.

MATERIAL. Russia, Moscow Area: 1♂, Petrovo-Dalnee, swamp, 28.08.1985, KYE; 1♀ (ZMUM Ta-1710), Kosino, Svyatoe Lake, 17.06.1908, N.S.Greze; Arkhangelsk Area: 1♀, Naryan-Mar, 07.1984, A.V.Tanasevitch; Tyumen Area: 1♀, Yamal, Obskaya Guba, Bereg Sorov, 5.08.1908, B.M.Zhitkov; 3♂, S-Yamal, Shchuchie, 17.07-7.08.1980, E.M.Veselova; 1♂ (PSU), S-Yamal, Khadyta River, *Salix* stand in a depression, 24.07.1981, leg. et det. SLE; Surgut distr., Yuganskiy State Reserve, swamp, 07.1988, V.G.Nonokshonov. Lithuania: 1♂ 2♀, Zhuvintas State Reserve, Eriophoreto-Sphagnetum, 2-28.06.1986, RM.

CATALOGUE. Russia: KRL [Uzenbaev, 1985, 1986, 1987]. LNN [Palmgren, 1943]. SVR [Pakhorukov, 1977].

DISTRIBUTION. Boreo-montaneous, "sphagnoicolous" species. USSR - N-European part, N of W-Siberia. C-, N-Europe, N-America from Alaska to Utah and New-York, USA.

*Clubiona (Clubiona) kulczynskii* Lessert, 1905.

*C.borealis* - Ermolajev, 1934; Fedotov, 1912b; Izmailova, 1989; Kulczynski, 1885; Strand, 1906.

*Curalensis* - Izmailova, 1975, 1989; Pakhorukov, 1979.

MATERIAL. Russia, Komi Republic: 7♀, Kortkeros distr., Dan, *Betula*, *Pinus* forest, and *Filipendula* near a river, 21-22.06.1988, KGM; Perm Area: 1♂ 2♀ (PSU), environs of Perm, cherry, 10-23.05.1987, leg. et det. SLE; Tyumen Area: 1♂ 1♀ (PSU), S-Yamal, Khadyta River, *Salix* floodplain forest, 3.08.1981, leg. et det. SLE; Novosibirsk Area: Toguchin and Kolyvan distr., Altai, Turochan distr.: 1♀, Soda Lake, rock stream, 5.09.1977, BPZ; 1♀, id, *Pinus sibirica* forest, 7.09.1979, BPZ; Krasnoyarsk Prov.: 3♀, Stolby State Reserve, 9.06.1980, KYE; 1♀, id., Mana River, 20.06.1990, ABR; C-Siberian State Reserve (different localities), ABR; Krasnoyarsk Prov., Ermakovo distr.: 1♀, Abakan-Kyzyl highway, Tanzybey, meadow, 20.07.1988, D.L.Grodnitskiy; 5♀, ca. 14 km SW of Tanzybey, 400-500 m, *Picea*, *Betula*, *Pinus sibirica* forest, 13.07.1990, DVL; 1♀, Abakan-Kyzyl highway, 180 km from Abakan, 1000-1200 m, 14.08.1988, D.L.Grodnitskiy; 2♀ (BIN 54), 2-3 km N of Aradan, Us River floodplain, ca. 840 m, 8-9.07.1990, DVL; 2♀ (BIN 57) 1♀, 2-3 km N of Aradan, *Picea*, *Pinus sibirica*, *Betula* forest, 840-850 m, 8-9.07.1990, DVL; 1♀, 25-30 km N of Aradan, 1600 m, stone stream, 9.07.1990, DVL; 2♂ 2♀ (BIN 61) 1♀, Oyskiy Pass, 1500-1600 m, subalpine meadow, 27-28.06, 10-11.07.1990, DVL; 2♀, 4-5 km S of Tanzybey, near Mutnaya River, 380 m, 26.06.1990, DVL; Krasnoyarsk Prov., Khakassia: 4♀, environs of Abakan, Ona River Canyon, 16.06.1990,


ABR; 1♂ 5♀, Askiz distr., 10-15 km SW of Birikchul, meadow, 1400-1500 m, 17-18.07.1990, DVL; 1♀ (BIN 58) 1♂ 6♀, ca. 8 km E of Birikchul, steppe slopes and floodplain *Betula-Salix* forest, 16-18.07.1990, DVL; Tuva: 2♀, Tandinskij distr., Chagytay Lake, 1100 m, sparse stand, 26.06.1989, DVL; 1♂ 3♀, environs of Chagytay Lake, N slope of E Tannu-Ola Mts., 1200-1700 m, forest, 27.06.1989, DVL; 1♂ 5♀, Piy-Khem distr., ca. 5 km NW of Seserlig, ca. 25 km NW of Kyzyl, *Larix*, *Salix* forest along a stream, 1100-1200 m, 24.07.1989, DVL; 1♀ (BIN 49), id., *Larix*, *Salix* forest, litter, 1000-1200 m, 2.05.1990, DVL; 2♀ (BIN 53), ca. 3 km NW of Seserlig, near a stream, *Salix* stand, litter, 1000-1100 m, 29.06.1990, DVL; 2♀, ca. 10 km NW of Seserlig, 1800-2000 m, *Larix*, *Picea*, *Pinus* forest, 25.07.1989, DVL; 1♂ (BIN 52), Kurtushinskiy Mt.Ridge, ca. 10 km NW of Shivilig, meadow, 1100 m, 6.07.1990, DVL; 1♂ 4♀, id., *Betula* crowns, 5.07.1990, DVL; 1♀ 2♀, Azas State Reserve, 1-7.06.1990, ABR; Yakutia: 1♂, Zhigansk, *Larix* taiga, 4.07.1989, KYE; Chita Area, Sokhondo State Reserve: 1♀, W slope of Sagan-Ula, 1700 m, upper forest bound, 06.1982, N.Shevrevya; 1♂ 1♀, confluence of Larionova Stream and Agutta River, meadow, 1300-1400 m, 12.06.1991, DVL; Khabarovsk Prov.: 1♂ (Kh), Verkhnebureinskij distr., Badzhalskiy Mt.Ridge, Siberian dwarf-pine, *Rhododendron*, 1400 m, 11.08.1989, DKK; 1♂, id., Mogdy River, *Larix*, Siberian dwarf-pine sparse stand, 1000 m, 8.08.1989, DKK; Sakhalin: 1♀, Aniva distr., Pik Chehova Mt., 1000 m, 3.07.1986, AMB;

CATALOGUE. Russia: MRM [Fedotov, 1912b; Palmgren, 1943]. KOM: Pechoro-Ilychskiy State Reserve [Pakhorukov, 1980]. KST [Mikhailov, 1989]. PRM [Gridina, 1992]: Basegi State Reserve [Esyunin, 1991]. SVR [Kharitonov, 1926b; Pakhorukov, 1977, 1979]. BShK: Bashkirskiy State Reserve [Pakhorukov, Efimik, 1988\*]. ChLB: Ilmen State Reserve [Pakhorukov, Polyanin, 1987\*; Polyanin, Pakhorukov, 1988\*]. TMN: Polar Ural [Tanasevitch, 1985\*]. TMS [Ermolajev, 1934]. KRN: Mirnoe [Eskov\*, 1981, 1988], Evenkia [Eskov, 1988\*], Boguchany distr. [Izmailova, Verzhutskiy, 1981; Izmailova, 1989: C.Kulczynski]. YaKT: Amga, Chona Rivers [Mikhailov, 1991]. MGD: Kolyma River upper flow, environs of Magadan [Mikhailov, 1991]. KMCh [Kulczynski, 1885; Strand, 1906-cit; Sytshevskaja, 1935\*; Mikhailov, 1991]. IRK [Izmailova, 1989: C.b. + C.u.]: Verkhnee Priangarye [Izmailova, 1975: C.b.]. ChLT: Sokhondo State Reserve [Danilov, Kurtova, 1991]. KhAB: Okhotsk distr. [Mikhailov, 1991]. SAKh: N-Sakhalin [Mikhailov, 1991]. Kazakhstan: EKA [Savelieva, 1972ab, 1976, 1979].

DISTRIBUTION. USSR - N-European part, all Siberia, N-(and montaneous) Far East. C-Europe (sporadic), N-Europe, N-America from Alaska to Colorado and N-Carolina, USA.

*Clubiona (Clubiona) interjecta* L.Koch, 1879.

MATERIAL. Russia, Tyumen Area: 1♀, Novyi Urengoy, *Betula* forest, 19.07.1984, NMP; 1♂, Tarko-Sale, hardwood forest, 24.07.1984, NMP; Krasnoyarsk Prov.: 1♀ (ZMH) Minusinsk, 1881, Roschier; 2♀ (BIN 59), Shushenskoe distr., ca. 3-5 km E of Shushenskoe, Oya River floodplain, mesophytous meadow with sparse willows, 26.06.1990, DVL; Tuva: 3♂ 2♀, environs of Kyzyl, right bank of Yenisey, floodplain forest, litter, 7.06.1989, DVL; 1♂, Piy-Khem distr., ca. 5 km NW of Seserlig, ca. 25 km NW of

Fig.8. *Clubiona (Clubiona) maracandica* Kroneberg, 1875, female lectotype: a) epigyne; b) vulva. Scale = 0.14.Рис.8. *Clubiona (Clubiona) maracandica* Kroneberg, 1875, лектотип, самка: а) эпигина; б) вульва. Масштаб 0.14.

Kyzyl, Larix, Salix forest, along a stream, ca. 1100-1200 m, 24.07.1989, DVL; 1♂, Erzin distr., Eder-Elezin sands, Tere-Khol Lake, Stipa tussocks, 8-9.08.1989, DVL; Yakutia: 1♀, Zhigansk, Larix taiga, 4.07.1989, leg. KYE, det. YMM; 1♀, Sangar, 23-28.07.1989, KYE; 2♂, Aldan River, 3 km upstream of Krestkha-lidzhey, 6.09.1990, IDS; Buryatia: 1♂, Kyakhta distr., Duren, Chikoy River floodplain, 26.06.1986, BPZ; 1♂ (BIN 6), id., 15.08.1986; 1♀ (UU), Pribaikalskiy distr., Turuntaevo, Betula sparse stand, 24.06.1983, SND; 1 (UU), Selenginsk, Pinus, Betula forest, 24.08.1983, SND; 2 (UU), Ulan-Ude, Selenga River floodplain, bird-cherry trees, Salix tussocks, 12.06.1983, SND; Chita Area: 1, Kyra distr., 3-5 km E of Kyra, poplar forest in a floodplain, 800-850 m, 19.06.1991, DVL; 1 (UU), Dahurian State Reserve, Nizhnii Tsasuchey, Pinus forest, in a grass, 26.06.1984, SND; 2 1, Khilok distr., Bada, 12.08.1987, A.P.Rasnitsyn; Amur Area: 1 (ZIP), Dep River (right tributary of Zeya River), 07-08.1911, P.Rippas; Khabarovsk Province: 1 (ZIP), Amur River, near Sredne-Tambovskoe, 18.07.1910, VKS; 1 (Kh), Tsimmermanovskiy Isl. Amur River, broadleaved forest, *Polygonatum*, *Vicia*, 26.06.1990, DKK; 1 (Kh), Ulcha distr., 445th km of Amur River fairway, broadleaved forest, crowns, 26.06.1990, DKK; Primorye: 1 (ZIP), S part of Prihankayskie forest-steppe, environs of Khorol, forest, 18.07.1968, FZP; 1 (ZIP), Prihankayskie forest-steppe, Mo River valley, 21.07.1968, FZP.

Mongolia: 4 ♂, Ulan-Bator, near Tuul River, 48°N, 107°E, S.Heimer.

CATALOGUE. Russia: KRN: Evenkiya [Eskov, 1988\*], Krasnoyarsk [Koch, 1879; Holm, 1973]. YaKT: Yana River mouth [Kulczynski, 1908\*]. MGD: Kolyma River upper flow [Mikhailov, 1991]. KMCh [Sytshevskaja, 1935\*; Mikhailov, 1991]. IRK [Izmailova, 1989]. BUR: Barguzin State Reserve [Sternbergs, 1981c]. BUR + ChIT: Muy Kettle [Verzhutskiy, Bessolitsyna, Seryshev, 1985]. AMR: Blagoveshchensk [Strand\*, 1907, 1909]. KhAB: Okhotsk distr., Khabarovsk, Khor, Jewish AO [Mikhailov, 1991]. PRI: Bikin River [Mikhailov, 1991]. SAKh: Sakhalin [Mikhailov, 1991].

DISTRIBUTION. USSR - all Siberia and Far East. Mongolia.

*Clubiona (Clubiona) maracandica* Kroneberg, 1875.

Fig.8a,b.

1875 Cn. Kroneberg: 16, Tab.2., Fig.8 (♀).

MATERIAL. Uzbekistan, Samarkand: 1♀ (ZMUM Ta-4800, lectotype, designated here), 2♀ (ZMUM Ta-1064, paralectotypes).

CATALOGUE. Uzbekistan: SMR [Kroneberg\*, 1875; Alimdzhanov, Bronstein, 1956-cit; Bronstein, Murtazaev, 1974].

Wrong identifications. Turkmenia: Repetek State Reserve [Kaplin, 1978; Mikhailov, Fet, 1986-cit].

DIAGNOSIS. By the structure of genitalia the species differs from other members of the *reclusa*-group by a combination of the central position of the copulatory openings, the ridged lateral and posterior margins of the copulatory openings, and the strongly horizontal (in a ventral view) posterior edges of the spermathecae.

DESCRIPTION. FEMALE. Total length 6.25. Carapace length 2.50, width 1.50, their ratio 1.67. Carapace and chelicerae reddish, legs straw-coloured. Leg measurements:

	I	II	III	IV
Femur	1.45	1.55	1.35	1.90
Patella	0.83	0.85	0.73	0.88
Tibia	1.23	1.25	0.95	1.48
Metatarsus	0.88	0.95	1.13	1.93
Tarsus	0.58	0.63	0.50	0.63

Leg armature: F I-II d.1.1.2, III-IV d.1.1.3, Pt III-IV rl.1, Ti I-II v.2.2, III d.2.2, v.1.1, IV d.2.2, v.1.1.1, Mt I-II v.2, III d.2.1.2, I.1.2, v.2.2, IV d.2.1.2, I.2.2, v.2.1.2.

Abdomen: length 3.70, width 2.25, their ratio 1.64; greyish-brown. Epigyne as in Fig. 8a,b.

MALE unknown.




Fig.9. *Clubiona (Clubiona) rybini* sp.n., male palp: a) ventral view; b) extero-ventral view; c) tibial apophysis, external view. Scale = 0.14.

Рис.9. *Clubiona (Clubiona) rybini* sp.n., пальпа самца: а) вентрально; б) наружно-вентрально; в) вырост голени, снаружи. Масштаб 0.14.

**DISTRIBUTION.** The species is known only from the type locality. Other data are wrong.

**REMARKS.** Due to the very poor condition of type material, I confine myself to study of only one of paralectotype specimens.

Two Middle Asian representatives of the *reclusa*-group, *Cmaracandica* and *C.rybini* sp.n. differ from the other species by the round copulatory openings closely narrowed and directed to the depth of the epigyne.

#### *Clubiona (Clubiona) rybini* Michailov, sp.n.

Figs. 9a-c, 10a-c.

*Cmaracandica* - Folkina, 1978; Mikhailov, 1992 (part.).

**MATERIAL.** Holotype: 1♂ (ZMUM Ta-4809), Kazakhstan, environs of Alma-Ata, Aksay Canyon, ca. 1300 m, litter, 28.08.1983, YMM, Ch.K.Tarabaev. Paratypes: 6♂ 5♀, as holotype; 1♀, Aksay Canyon, 15.09.1984, SVO; 1♂ 1♀, Alma-Ata State Reserve, Canyon of Sredniy Talgar River, 1840 m, 18.09.1984, A.B.Nenilin, SVO. Kirghizia: 1♂ 1♀ (ZIP), Osh, 1-2.11.1933, V.N.Shnitnikov; 1♂, 2 km N of Frunze, Karagachevaya Roshcha, 17.06.1979, SLZ; 1♀, Frunze, Chon-Aryk, coll. Sarkisova; 1♀ (ZMUM Ta-4810), Kirghizskiy Mt.Ridge, 20 km S of Frunze, Malinovoe Canyon, 1500 m, *Juniperus* sparse stand belt, 27.07.1983, SVO; 1♂, Ferghanskiy

Mt.Ridge, Kara-Alma, 4-6.06.1979, SLZ; 1♂ 3♀, Ferghanskiy Mt.Ridge, environs of Arslanbob, 22-25.10.1986, SLZ; 1♀, W.Tien-Shang Mts., Ugamskiy Mt.Ridge, Kaynar-Say, *Juglans* forest, 23.04.1983, SLZ; 3♂ 2♀ 3♂ sbad, Osh Area, Gulcha, hospital, nest of *Turdus merula*, 5.06.1984, S.N.Rybin; 2♀, Osh Area, Naukat, Yanghi-Bazar, nest of *Turdus merula*, 25.09.1983, S.N.Rybin.

**CATALOGUE.** Kazakhstan: AAO: Zailiyskiy Alatau Mt. Ridge [Folkina, 1978: Cm]. SE-Kazakhstan, Kirghizia [Mikhailov, 1992: Cm].

**DIAGNOSIS.** By the palp structure the male differs from other members of the *reclusa*-group by both the very broad protector (the breadth in the basal and middle parts are equal) and a combination of the single spine-like apophysis of the embolar part and the little-developed back tooth ("hook") of the "harpoon" of tibial apophysis. The female is more close to *Cmaracandica* differing by the copulatory openings situated in the posterior part of the epigynal plate, the anteriorly and laterally ridged margins of the openings, and the posterior edges of the spermathecae mesally projected behind.

**DESCRIPTION. MALE (10).** Total length 4.81+0.92. Carapace: length 2.15+0.36, width 1.53+0.26, their ratio 1.41+0.12. Carapace, chelicerae and legs reddish or straw-coloured. Leg measurements:




Fig.10. *Clubiona (Clubiona) rybini* sp.n., female: a) epigyne; b) vulva; c) epigyne, caudal view. Scale = 0.14.

Рис.10. *Clubiona (Clubiona) rybini* sp.n., самка: а) эпигина; б) вульва; в) эпигина, вид сзади. Масштаб 0.14.

Legs	I	II	III	IV
Femur	♂ $1.44 \pm 0.30$	$1.49 \pm 0.28$	$1.34 \pm 0.22$	$1.78 \pm 0.42$
	♀ $1.52 \pm 0.30$	$1.56 \pm 0.28$	$1.43 \pm 0.33$	$1.99 \pm 0.38$
Patella	♂ $0.79 \pm 0.14$	$0.80 \pm 0.14$	$0.69 \pm 0.11$	$0.78 \pm 0.17$
	♀ $0.89 \pm 0.16$	$0.92 \pm 0.16$	$0.79 \pm 0.10$	$0.93 \pm 0.16$
Tibia	♂ $1.35 \pm 0.22$	$1.49 \pm 0.21$	$0.98 \pm 0.16$	$1.47 \pm 0.32$
	♀ $1.26 \pm 0.24$	$1.32 \pm 0.22$	$0.98 \pm 0.21$	$1.56 \pm 0.25$
Metatarsus	♂ $1.01 \pm 0.22$	$1.06 \pm 0.18$	$1.18 \pm 0.22$	$1.88 \pm 0.47$
	♀ $0.94 \pm 0.18$	$0.97 \pm 0.20$	$1.15 \pm 0.20$	$1.99 \pm 0.35$
Tarsus	♂ $0.66 \pm 0.13$	$0.67 \pm 0.10$	$0.51 \pm 0.06$	$0.62 \pm 0.10$
	♀ $0.64 \pm 0.11$	$0.65 \pm 0.12$	$0.52 \pm 0.10$	$0.67 \pm 0.12$

Leg armature: F I d.1.1.2, II d.1.1.2 (1.1.3), III-IV d.1.1.3, Pt III-IV rl.1, Ti I v.2.2, II v.2.2 (1.2), III d.2.2, v.1.1 (1.1.1), IV d.2.2, v.1.1.1, Mt I-II v.2, III d.2.1.2, I.1.2 (2.2), v.2.2, IV d.2.1.2, I.2.2 (2.1.2), v.2.1.2.

Palp as in Fig.9a-c. Lengths of cymbium, tibia, patella, and femur are  $0.75 \pm 0.10$ ,  $0.25 \pm 0.05$ ,  $0.35 \pm 0.07$ , and  $0.60 \pm 0.16$ , respectively.

Abdomen: length  $2.67 \pm 0.57$ , width  $1.41 \pm 0.37$ , their ratio  $1.91 \pm 0.38$ ; pale or pale brown with both




Fig.11. *Clubiona (Clubiona) saurica* sp.n., female: a) epigyne; b) vulva. Scale = 0.14.

Рис.11. *Clubiona (Clubiona) saurica* sp.n., самка: а) эпигина; б) вульва. Масштаб 0.14.

a lanceolate pattern and a posterior part brown dorsally.

FEMALE (10). Total length  $5.88 \pm 1.82$ . Carapace length  $2.66 \pm 0.48$ , width  $1.85 \pm 0.34$ , their ratio  $1.44 \pm 0.12$ . Carapace and legs reddish or straw coloured. Chelicerae reddish, straw or chestnut-coloured. Leg measurements see above. Leg armature as in male, with following exceptions: F II d.1.1.2, III d.1.1.3 (1.2.3), Ti II v.2.2, III d.2.2, v.1.1, IV d.2.2, v.1.1.1 (1.1), Mt III d.2.1.2 (2.2), I.1.2, v.2.2, IV d.2.1.2, I.2.2, v.2.1.2 (2.1.1).

Abdomen: length  $3.36 \pm 1.15$ , width  $2.04 \pm 0.95$ , their ratio  $1.67 \pm 0.37$ ; pale, sometimes with both a lanceolate pattern and a posterior part brown dorsally. Epigyne as in Fig.10a-c.

DISTRIBUTION. SE-Kazakhstan, Kirghizia.

DERIVATIO NOMINIS. The species is dedicated to the Russian entomologist and parasitologist, Stanislav Nikolaevich Rybin (Osh, formerly N-Osetia).

#### *Clubiona (Clubiona) saurica* Michailov, sp.n.

Fig. 11a,b.

MATERIAL. Kazakhstan, East-Kazakhstan Area: 1♀ (holotype, ZMUM Ta-4811), Zaisan distr., Saur Mts., Akkolka River valley (Kenderlyk River Basin), *Salix* along a stream, 27.06.1990, KYE.

DIAGNOSIS. The female differs from other members of the *reclusa*-group by the very long atrium; the length of atrio-spermathecal tube not




Fig.12. *Clubiona (Clubiona) bashkirica* sp.n., male palp: a) ventral view; b-c) tibial apophysis, extero-ventral (b), external view (c). Scale = 0.14.

Рис.12. *Clubiona (Clubiona) bashkirica* sp.n., пальпа самца: а) вентрально; б-с) вырост голени, наружно-вентрально (б), снаружи (с). Масштаб 0.14.

exceeding 1/3 of atrial length. The posterior edge of the epigynal plate with a tiny protrusion with narrowed copulatory openings of a typical *reclusa* pattern.

**DESCRIPTION. FEMALE.** Total length 5.90. Carapace length 2.40, width 1.75, their ratio 1.37.

Carapace and legs straw-coloured. Chelicerae reddish-brown. Leg measurements:

	I	II	III	IV
Femur	1.50	1.55	1.30	1.85
Patella	0.85	0.88	0.73	0.88
Tibia	2.40	1.28	0.93	1.58
Metatarsus	0.98	0.98	1.13	2.05
Tarsus	0.63	0.68	0.53	0.65

Leg armature: F I-II d.1.1.2, III-IV d.1.1.3, Pt III-IV rl.1, Ti I-II v.22, III d.2.2, v.1.1, IV d.2.2, v.1.1.1, Mt I-II v.2, III d.2.1.2, l.1.2, v.2.2, IV d.2.1.2 (2.2.2), 1.2.2, v.2.1.2.

**Abdomen:** length 3.50, width 2.10, their ratio 1.67; greyish-brown. Epigyne as in Fig. 11a,b.

**MALE unknown.**

**DISTRIBUTION.** The species is known only from the single locality.

*Clubiona (Clubiona) bashkirica* Michailov, sp.n.  
Figs. 12a-c, 13a,b.

**MATERIAL.** Holotype ♂ (ZMUM Ta-4812) Russia, Bashkiria, Burzyanskiy distr., Ural-Tau Mts., Bashkirian State Reserve, Malaya Yamal-Yalgha River, flood plain, moist meadow, 6.08.1986, VEE. Paratypes: 1♂ (ZMUM) as holotype; 2♀ Bashkirian State Reserve, Kazmash Kordon, Uzyan River, moist meadow, 9.08.1986, VEE; 1♀ (ZMUM Ta-4813), Burzyanskiy distr., Shulghan-Tash State Reserve, meadow, 29.06.1985, VEE.

**DIAGNOSIS.** By the palpal structure the male is more close to the *lutescens* group, differing by the slender and short embolus. By the epigynal structure the female is more close to the *reclusa* group, but differs by the oval (not slit-like) copulatory openings.

**DESCRIPTION. MALE (2).** Carapace length




Fig.13. *Clubiona (Clubiona) bashkirica*, female: a) epigyne, b) vulva. Scale = 0.14.

Рис.13. *Clubiona (Clubiona) bashkirica*, самка: а) эпигина, б) вульва. Масштаб 0.14.

2.40-2.45, width 1.70-1.85, their ratio 1.30-1.44. Caparace and chelicerae reddish, legs straw-coloured. Leg measuremets:

Legs:	I	II	III	IV
Femur	♂ 1.83-1.85	1.78-1.88	1.58-1.63	2.13
	♀ 1.55-1.98	1.58-1.90	1.40-1.78	1.93-2.45
Patella	♂ 0.88-0.95	0.93-0.98	0.78-0.80	0.90-0.95
	♀ 0.83-1.13	0.85-1.13	0.70-0.88	0.88-1.10
Tibia	♂ 1.53-1.58	1.53-1.63	1.15-1.18	1.75-1.78
	♀ 1.20-1.50	1.25-1.58	0.95-1.15	1.55-1.88
Metatarsus	♂ 1.18-1.20	1.15-1.23	1.33-1.40	2.13-2.20
	♀ 0.90-1.18	0.93-1.18	1.13-1.40	1.85-2.38
Tarsus	♂ 0.78	0.75-0.78	0.55	0.70-0.75
	♀ 0.68-0.78	0.65-0.80	0.50-0.63	0.65-0.75

Leg armature: F I-II d.1.1.2, II-IV d.1.1.3, Pt III-IV rl.1, Ti I-II v.2.2, III d.2.2, v.1.1, IV d.2.2, v.1.1.1, Mt I-II v.2, III d.2.1.2, 1.1.2, v.2.2, IV d.2.1.2, 1.2.2, v.2.1.2.

Palp as in Fig. 12a-c. Lengths of cymbium, tibia, patella, and femur are 0.75-0.78, 0.25, 0.30-0.38, 0.63, repectively.

Abdomen length 2.35-2.50, width 1.40-1.60, their ratio 1.56-1.68, brown.

FEMALE (3). Caparace length 2.45-2.90, width 1.80-2.00, their ratio 1.30-1.45, reddish brown. Chelicerae brown, legs straw-coloured. eg measurements se above. Leg armature as in male, but Ti II v.2.2 (1.2).

Abdomen length 3.80-4.0, width 2.35-2.50, their ratio 1.58-1.70, grey-brown or brown. Epigyne and vulva as in Fig.13a,b.

REMARKS. The new species occupies an intermediate position between the *lutescens* and *reclusa* groups. The male of *C.bashkirica* can be compared with the first group by the presence of the non-specialized protector at the tegular part of embolus. The shape of the tibial apophysis reminds of a garpoon-like form, especially by the external view (Fig. 12c). In *C.bashkirica* female the interval between the atrium and the spermatheca in the vulva is considerable, but the general composition of vulval structures makes the new species close to the *reclusa* group, specific difference see Diagnosis.

By the leg measurements female specimens from Bashkirian Reserve are more close to each other than Shulghan-Tash Reserve one, the latter being more large.

DISTRIBUTION. The species is known only from moist meadows of S-Urals.

The *caeruleascens* - group.

Includes a single species,  
*Clubiona caeruleascens* L.Koch, 1866.

MATERIAL. Russia, Komi Republic: 1♂ 2♀ (ZMUM),

Kortkeros distr., Dan, *Betula-Pinus* sandy forest, birch leaves, 21.06.1988, KGM; Altay Prov.: 1♀ (BIN 3), Talmenka distr., Larichikhino Lespromkhoz, Shipinskoe Forestry, pine-birch forest, 24.08.1979, BPZ; Krasnoyarsk Prov.: 2♀ (ZMUM), Stolby Reserve, Lalitina Stream valley, litter, 9.06.1980, KYE; 1♀ (ZMUM), Stolby Reserve, *Picea* forest near a stream, 10.06.1980, KYE; 1♀ (ZMUM) environs of Krasnoyarsk, Ermakovo airport, clearing in pine-birch forest, 31.08.1982, KYE; 1♀ (ZMUM) Bolshaya Murta distr., right bank of Yenisey River, opposite of Yukseevo, 10.09.1988, A.V.Gurov, C.M.Loshchev; 1♂ (ZMUM) W-Sayan Mts., Ermakovskoe, pine forest with birches, 19.08.1984, ABR; 1♀ (ZMUM) Ermakovskoe distr., ca. 15 km E of Ermakovskoe, *Pinus-Betula-Populus tremula* forest, 26.06.1990, DVL; Tuva: 1♂ (ZMUM), Tandinskiy distr., near Chagytay Lake, N slope of E-Tannu-Ola, 1200-1700 m, forest, 27.06.1989, DVL; Irkutsk Area: 1♀ (BIN 4), Novaya Lisikha, *Populus tremula-Betula* forest, 28.06.1979, BPZ; 1♂ (ZMUM), Slyudyanka, forest, grass, 6.09.1966, coll.et det. V.P.Tyshchenko; 1♂ 3♀ (ZMUM) Slyudyanka distr., Marituy, *Aibus*, *Betula*, in leaves, 5.07.1988, KGM, I.M.Marova; 3♂ 4♀ (ZMUM) along Marituy Stream, *Aibus*, *Rubus*, in leaves, 23.07.1990, KGM; 1♂ 2♀ (UU) Baykalsk, Solzan, birch forest, 22.08.1981, SND; Buryatia: 2♂ 1♂-inad 3♀ (UU), Selenginsk, birch forest, 25.08.1983, coll. et det. SND; Sakhalin: 2♀ (ZMUM), Yuzhno-Sakhalinsk, 1.07.1989, AMB. Kazakhstan, 1♀ (ZMUM) North-Kazakhstan Area, Sokolovka distr., Bolshaya Malyshka, pine forest, 10.06.1986, coll.et det. DVL.

CATALOGUE. Russia: KRL [Palmgren, 1943; Uzenbaev, 1986]. LNN [Siemaschko, 1861; Palmgren, 1943]. KLN: Belskiy Uezd [Pokrovskiy, 1930]. YaRS [L.Becker, 1879]. MSK [Freiberg\*, 1894, 1897; Greze, 1914; Pokrovskiy, 1925\*; Pereleshina, 1928a\*; Mikhailov, 1983ab]. BRN [Kharitonov, 1936; Lavrov, 1968]. KUR [Pichka, 1965, 1984b]. LPTs: Galichya Gora State Reserve [Panteleeva, 1982]. BLG: Valuyki [Kulczynski, 1913], "Les na Vorskle" State Reserve [Ovsyannikov, 1937; Marusik, 1984]. VRN [Pichka, 1965, 1967; Pichka, Skufyin, 1981]. Middle Povolzhye [Aleynikova, 1964, 1968]. GRK [Greze, 1915a]. ChVSh [Holzmayer, 1934]. MRD [Fatersov, 1984]. TTR [Garkusha, 1977; Utochkin, 1977; Zheltukhina, Utochkin, 1984]. KBSh: Zhiguli State Reserve [Krasnobaev, 1983; Krasnobaev, Ovtsharenko, 1986]. STV: Pyatigorsk [Mikhailov, 1990b]. SOA [Mikhailov, 1990b]. KOM: Pechoro-Ilychskiy State Reserve [Pakhorkov, 1980]. KST [Veselova, 1986\*; Veselova, Mikhailov, 1986, 1987; Mikhailov, 1989]. PRM [Kharitonov, 1926a; Kharitonov, Bezzrezeva, 1970; Azheganova, Kopyltsova, Parfenova, 1976; Azheganova, Glukhov, 1981; Gridina, 1992]. SVR [Pakhorkov, 1977]. BShK: Bashkirskiy State Reserve [Pakhorkov, Efimik, 1988\*]. ChLB: Troitskiy Reserve [Azheganova, 1951; Pakhorkov, 1985], Ilmen State Reserve [Pakhorkov, Polyanin, 1987; Polyanin, Pakhorkov, 1988, 1992]. OMS: Omsk [Spassky, Lavrov, 1928\*]. TMS: Tomsk [Ermolajev, 1934]. ALT: Kulunda Steppe [Litvinchuk, 1980]. KRN: Krasnoyarsk [Koch, 1879; Holm, 1973]. MGD: environs of Magadan [Mikhailov, 1991]. IRK [Izmailova, 1989]. BUR [Izmailova, 1989]. KhAB: Bolshekhekhtsyrskiy State Reserve [Mikhailov, 1991]. SAKh: S-Sakhalin [Mikhailov, 1991]. Estonia [Vilbaste, 1987]. Latvia [Sternbergs, 1979, 1981a, 1983; Prieditis, Sternbergs, 1981]. Byelorussia [Litvinova et al., 1980]. Belovezhskaya Pushcha [Chebotareva, 1987]. MIN [Zhukovets, 1989]. Ukraine: ZAK [Legotay, 1971,

1989; Polozhentsev, Akimtseva, 1980; Legotay, Sekerskaya, 1982]. LVV + IFR [Legotay, Tarasyuk, 1964]. "S-Russia" [Nosek, 1895]. ChRK: Kanev State Reserve [Pichka, 1974]. KhRK [Kirilenko, Legotay, 1981]. KhRS: lower flow of Dnepr River [Talanov, Nazarenko, 1979]. Moldavia [Talitskiy, Legotay, 1982; Matsyuk, Yanyuk, Verlan, 1984]. Georgia [Mikhailov, 1990b]. Armenia [Mikhailov, 1990b]. Azerbaijan [Atakishiev, 1969; Dunin, 1989; Mikhailov, 1990b]. Kazakhstan: EKA [Savelieva\*, 1970, 1979].

DISTRIBUTION. USSR - European part, Caucasus, Siberia, Far East (sporadically). S-Europe: Balkans, Italy, C-, N-Europe, Japan: Hokkaido.

The *chabrovi* - group.

Includes a single species,

*Clubiona (Clubiona) chabrovi* Michailov, 1991.

CATALOGUE. Russia: KhAB: Bolshekhekhtyrskiy State Reserve [Mikhailov, 1991].

DISTRIBUTION. The species is known only from the type locality.

*Clubiona (Clubiona) incertae sedis.*

REMARKS. This artificial grouping includes a lot of E-Asian species: *C.venusta* Paik, 1985 ( $\varphi$  from Korea, resembles the *pallidula* - group), *C.haeinsensis* Paik, 1990 ( $\sigma\varphi$  Korea) and *C.bwangbakensis* Paik, 1990 ( $\sigma$  Korea), both close to the *obesa* and *pallidula* - groups, *C.kasanensis* Paik, 1990 ( $\sigma\varphi$  Korea, may constitute a separate group), and some species described by Bösenberg and Strand, Schenkel, Ono, etc.

## References

- Alexeev V.A. 1986. [To the study of the fauna of terrestrial invertebrates of Darwinian State Reserve during 40 years]/ / Fauna i ekologiya bespozvonochnykh zhivotnykh v zapovednikakh RSFSR. Moscow. P.4-20 [in Russian].
- Aleynikova M.M. 1964. [Soil fauna of different landscapes of Middle Povolzhye]/ / Pochvennaya fauna Srednego Povolzhya. Moscow, Nauka Publ. P.5-51 [in Russian].
- Aleynikova M.M. 1968. [Soil-dwelling invertebrates of different forest types in Srednee Povolzhye]/ / Materialy po faune i ekologii pochvoobrazuyushchikh bespozvonochnykh. Kazan, Kazan State Univ. Press. P.3-97 [in Russian].
- Aleynikova M.M., Tyschenko V.P. 1969. [Fauna and landscape distribution of spiders (mainly connected with a soil) in Middle Povolzhye]/ / Problemy pochvennoy zoologii. Materialy Tretiego Vsesoyuznogo soveshchaniya. Moscow, Nauka Publ. P.15-16 [in Russian].
- Alimdzhanov R.A., Bronstein Ts.G. 1956. [Invertebrate animals of Zeravshan River Valley]. Tashkent - Samarkand, AN UzbSSR Press. 348 pp. [in Russian].
- Anchipanova Ya.Ya., Sternbergs M.T. 1984. [Peculiarities of the development of the spider *Pholidromus aureolus* (Cl, 1757) (Aranei, Thomisidae)]// Trudy Latv. selskokhoz. akad. No.213. P.5-9 [in Russian].
- Andreeva E.M. 1975. Distribution and ecology of spiders (Aranei) in Tadzhikistan// Fragm. faun. Mus. zool. Warszawa. T.20. No.19. P.323-352.
- Andreeva E.M. 1976. [Spiders of Tadzhikistan. Fauna and zonal-ecological distribution]. Dushanbe, Donish Publ. 193 pp. [in Russian].
- Astakhova E.V. 1974. [To the study of the spider fauna of Kharkov and Poltava Areas]// Vestn. Kharkovsk. Univ. No.105 (Biol. No.5). P.94-97 [in Russian].
- Astakhova E.V. 1978. [To the arachnological subdivision of the steppes of Levoberezhnaya Ukraine]// Vestn. Kharkovsk. Univ. No.164. P.71-72 [in Russian].
- Atakishiev T.A. 1969. [Spiders as symbionts of the honey bee]// Uchenye zapiski Kazansk. gos. veterin. inst. Vol.105. P.317-323 [in Russian].
- Azheganova N.S. 1951. [To the spider fauna of Troitskiy forest-steppe Reserve]// Izv. Est.-Nauch. Inst., Perm Gos. Univ. T.13. No.2-3. P.137-156 [in Russian].
- Azheganova N.S. 1968. [Short identification book of spiders (Aranei) of the forest and forest-steppe zones of USSR]// Opredelitel po faune SSSR, izdavaemye Zool. Inst. AN SSSR. Leningrad, Nauka Publ. No.98. 149 pp. [in Russian].
- Azheganova N.S., Glukhov M.F. 1981. [Fauna and ecology of spiders of the environs of the biological station (V. Kvazhva)]// Fauna i ekologiya nasekomykh. Perm, Perm State Univ. Press. P.21-32 [in Russian].
- Azheganova N.S., Gorshkov P.K. 1973. [Spiders of the burrows of carnivorous mammals of Volzhsko-Kamskiy State Reserve]// Uch. zap. Perm. ped. inst. Vol.109. P.61-68 [in Russian].
- Azheganova N.S., Kopylitsova L.S., Parfenova T.V. 1976. [Spiders of forest lower strata of the environs of the biological station V. Kvazhva]// Biogeografiya i kraevedenie. Perm. No.4. P.42-48 [in Russian].
- Azheganova N.S., Stenchenko T.I. 1977. [To the spider fauna of Amur Area]// Voprosy arakhnoentomologii. Perm, Perm State Univ. Press. P.106-111 [in Russian].
- Babidorich M.M. 1980. [To the study of spiders (Aranei) in the orchards of Moldavia]// Entomofagi vrediteley rasteniy. Kishinev. P.42-46 [in Russian].
- Babidorich M.M. 1983. [Annotated list of entomophages of the codling moth]// Inform. Bull. Vostochno-Palearkticheskoy Sektii MOBB. No.7. P.35-53 [in Russian].
- Bakhvalov V.F., Korshunov Yu.P. 1976. [To the spider (Aranei) fauna of southern Yamal]// Trudy Biol. Inst. Sibirs. Otdel. AN SSSR. No.18. P.38-46 [in Russian].
- Balogh J. 1940. Zur Kenntnis der Spinnenfauna der Nordostkarpaten// Fragm. faun. Hung. Vol.3. No.3. P.71-74.
- Becker A. 1888. Die Spinnen und fortgesetzte Mitteilungen über bei Sarepta vorkommende Insekten// Bull. Soc. Nat. Moscou. Nouv. ser. Vol.2. No.2. P.373-379.
- Becker L. 1879. Aranéides de Russie récueillies à Jaroslaw (Russie centrale) par M. Kokouyew// Ann. Soc. ent. belg. Vol.22. C.R. P.XLVII-XLVIII.
- Beer S.A. 1964. [On the fauna and ecology of spiders of Murman Area]// Zool. Zhurn. Vol.43. No.4. P.525-533 [in Russian].
- Beer S.A. 1968. [On the terrestrial fauna of spiders of Moscow Area]// Ibidem. Vol.47. No.1. P.131-134 [in Russian].
- Belimov G.T. 1975. [Some data on the arachnofauna of Middle Lena valley]// Biol. resursy Yakutii. Yakutsk. P.42-43 [in Russian].
- Belke G. 1853. Quelques mots sur le climat et la faune de Kamieniec-Podolski// Bull. Soc. Imp. Nat. Moscou. Vol.26. No.2. P.410-437.
- Belke G. 1859. Esquisse de l'histoire naturelle de Kamienietz-Podolski précédée d'un coup d'oeil sur les travaux des naturalistes des provinces occidentales de la Russie et du royaume de Pologne au XIXe siècle [1]// Ibidem. Vol.31. No.3. P.97-158.
- Belke G. 1866. Notice sur l'histoire naturelle du district de Radomysl (Gouvernement de Kief) [2]// Ibidem. Vol.39. No.2. P.491-526.

- Bergroth E. 1881. [Captures d'Aranéides en Sibérie]// Ann. Soc. ent. Belg. Vol25. C.R. P.X.
- Bösenberg W., Strand E. 1906. Japanische Spinnen// Abh. Senckenberg. naturforsch. Ges. Bd.30. H.1. S.93-422, Taf.III-XVI.
- Bronstein Ts.G., Murtazaev A.B. 1974. [Materials to the spider fauna of Samarkand Area]// Voprosy zashchity rasteniy. Trudy Uzbechk. Univ. Nov.ser. No.247. P.124-143 [in Russian].
- Bublik I.M. 1981. [To the ecology of spiders (Aranei) of ecosystems of Ciscarpathia]// Visn. Lvivsk. Univ. Ser. biol. No.12. P.86-90 [in Ukrainian].
- Bukovskiy V.I. 1940. [Population of invertebrates, mainly pests, of oak foliage in Crimean State Reserve]// Trudy Krymsk. Gos. Zapovednika. Vol2. P.39-169 [in Russian].
- Bulyginskaya M.A., Kalinkin V.M. 1985. [Effect of chemical treatment against the codling moth *Laspeyresia pomonella* L. (Lepidoptera, Tortricidae) on the accompanying species of Lepidoptera and their natural enemies]// Entomol. Obozr. 1985. Vol64. No.3. P.441-449 [in Russian].
- Chebotareva N.A. 1987. [Species composition and biotopical distribution of spiders (Aranei) of Belovezhskaya Pushcha]. Deposited in VINITI 13.07.87. No.5009-B87. 23 pp. [in Russian].
- Danilov S.N., Kurtova O.G. 1991. [Materials to the spider fauna (Aranei) of Sokhondo State Reserve]// Entomologicheskie problemy Baykalskogo regiona. Tez. dokl. regionaln. shkoly-seminara (6-8 avg. 1991, Ulan-Ude). P.34-35 [in Russian].
- Dmitrienko V.K. 1987. [Numerosity of predatory herpetobionts as an indicator of habitat optimality]// Ekologicheskaya otsekha mestoobitanii lesnykh zhivotnykh. Novosibirsk. P.16-29 [in Russian].
- Dondale C.D., Redner J.H. 1982. The sac spiders of Canada and Alaska (Araneae: Clubionidae and Anyphaenidae)// The insects and arachnids of Canada. Ottawa, Agriculture Canada. Pt.9. 194 pp.
- Dunin P.M. 1984. [Fauna and ecology of spiders (Aranei) of Apsheron Peninsula (Azerbaijanian SSR)]// Fauna i ekologiya paukoobraznykh. Perm, Perm State Univ. Press. P.45-60 [in Russian].
- Dunin P.M. 1989. [Fauna and altitudinal distribution of spiders (Arachnida, Aranei) of the Azerbaijan part of the southern macroslope of the Caucasus Major]// Fauna i ekologiya paukov i skorpionov. Arakhnologicheskiy sbornik. Moscow, Nauka Publ. P.31-39 [in Russian].
- Dwigubsky I. 1802. Primitiae faunae Mosquensis. Mosqua. P.I-VIII, 9-222.
- Ermolaev V.N. [Ermolajev W.] 1934. Materialen zur Spinnenfauna Westsibiriens. III. Die Spinnen der Stadt Tomsk// Folia zool. hydrobiol. Vol7. Pt.1. P.130-147.
- Ermolaev V.N. [Ermolajev W.] 1937. Beiträge zur Spinnentier-Fauna der Stadt Tobolsk (West-Sibirien)// Festschr. Strand. 1937. Bd.2. S. 519-524.
- Ermolaev V.N., Samko K.P. 1929. [Addition to the spider list of Tobolsk Town]// Byull. Obschch. izucheniya kraya pri Muzei Tobolskogo Severa. No.1-2. P.37-39 [in Russian].
- Eskov K.Y. 1981. [Analysis of the spatial distribution of spiders in Cis-Yeniseyan taiga]// Zool. Zhurn. Vol60. No.3. P.353-362 [in Russian].
- Eskov K.Y. 1986. [Spider fauna (Aranei) of the hypoarctic zone of Siberia]// Yuzhnye tundry Taymyra. Leningrad, Nauka Publ. P.174-191 [in Russian].
- Eskov K.Y. 1988. [Spiders (Aranei) of Middle Siberia]// Materiały po faune Sredney Sibiri i prilezhashchikh rayonov Mongoli. Moscow, Institute of Animal Evolutionary Morphology and Ecology. P.101-155,185 [in Russian].
- Esyunin S.L. 1991. [Arachnids of "Basegi" State Reserve (Pseudoscorpiones, Opiliones, Aranei, Parasitiformes: Ixodidae - annotated list of species)]// Flora i fauna zapovednikov SSSR. Moscow. No.38. 38 pp.[in Russian].
- Fatersov S.Ya. 1984. [Effect of forest fires on the species composition and numerosity of spiders]// Problemy pochvennoy zoologii. Tez. dokl. Vostmogo Vses. soveshch. Kniga 2. Ashkhabad. P.134-135 [in Russian].
- Fedotov D.M. 1912a. [To the spider fauna of Volyn Gouvernement]// Russk. Entomol. Obozr. Vol12. No.3. P.390-406 [in Russian].
- Fedotov D.M. 1912b. [To the spider fauna of Murman and Novaya Zemlya]// Ezheg. Zool. Muz. Imper. Akad. Nauk. Vol16. No.4. P.449-474, pl. VIII [in Russian].
- Fedotov D.M. 1912c. [Materials to the spider fauna of Tver Gouvernement]// Trudy Presnovodnoy Biol. Stants. St.-Peterb. obshch. estestvoispyt. Vol.3. P.53-134, tab.I-III [in Russian].
- Fedotov D.M. 1915. [To the spider fauna of Yaroslavl Gouvernement]// Ezheg. Zool. Muz. Imper. Akad. Nauk. Vol20. No.3. P.445-456 [in Russian].
- Fet V.Ya. 1983. [Spider fauna (Aranei) of SW-Kopetdagh Mts.]// Entomol. Obozr. Vol62. No.4. P.835-845 [in Russian].
- Fet V.Ya. 1985. [Zoogeographical analysis of the spider fauna of SW-Kopetdagh Mts.]// Fauna i ekologiya paukov SSSR. Trudy Zool. Inst. AN SSSR. Vol139. P.72-77 [in Russian].
- Fischer J.B. 1791. Versuch einer Naturgeschichte von Livland. Königsberg. 826 S., Tab.I-IV.
- Folkina M.Ya. 1978. [On the species composition of spiders predators of fruit aphids in orchards of Zailiyskiy Alatau Mt. Ridge]// Izv. AN KazSSR. Ser. biol. No.2. P.23-26 [in Russian].
- Freiberg P.R. 1894. [Araneae of Moscow Gouvernement on the materials collected by the Committee for the study of the fauna of Moscow Gouvernement in summer 1893]// Izv. Obshch. lyubit. estestvozn., antropol. i etnogr. Vol86/ Dnevn. Zool. otd. Obshch. Vol2. No.1-2. P.4-11 [in Russian].
- Freiberg P.R. 1897. [Araneida collected on the right bank of Oka River]// Ibidem. Vol2. No.5. P.23-39 [in Russian].
- Garkusha T.A. 1977. [Spider population of a spruce-broadleaved forest of the southern taiga subzone (Raifa, Tartarian ASSR)]// Voprosy arakhnoentomologii. Perm, Perm State Univ. Press. P.112-117 [in Russian].
- Garkusha T.A. 1978. [Soil fauna as a component of biogeocenoses of dried peatbogs. Spiders]// Pochvennaya fauna i biologicheskaya aktivnost osushennykh i rekultiviruemых torfyanikov. Moscow, Nauka Publ. P.28-36 [in Russian].
- Gnelitsa V.A. 1989. [Species composition of spiders of riparian biotopes of forest-steppe of Ternopol Area]// Problemy obshchey i molekularnoy biologii (Kiev). No.8. P.80-83 [in Russian].
- Greze N.S. 1909a. [Spiders of Don Area]// Trudy studencheskogo kruzhka po izucheniyu rodnoy prirody pri Moskovskom Univ. Vol4. P.99-111 [in Russian].
- Greze [Grese] N. 1909b. Die Spinnen der Halbinsel Jamal// Annu. Mus. Zool. Acad. Sci. S.Petersb. Vol14. P.325-331.
- Greze N.S. 1911. [Additions to the list of spiders of Moscow Gouvernement]// Dnevnik Zoologicheskogo otdeleniya Obshchestva lyubiteley estestvoznaniya, antropologii i etnografii. Vol3. No.10. P.55-61 [in Russian].
- Greze N.S. 1914. [Spiders (Araneae). List of spiders gathered in Podolskiy Uezd]// Estestvenno-istoricheskaya kollektiya gr. E.P.Sheremetevoy. Moscow. No.8. 15 pp. [in Russian].
- Greze N.S. 1915a. [Spiders of Nizhniy Novgorod Gouvernement]// Izvestiya Moskovsk. entomol. obshch. Vol1. P.104-115 [in Russian].
- Greze N.S. 1915b. [Spiders collected on the snow]// Ibidem. P.116-123 [in Russian].
- Gridina T.I. 1992. [Spatial interaction between *Formica* ants and

- arachnids under different landscape conditions]// Fauna i ekol paukov, skorpionov i lozhnoskorponov SSSR. Trudy Zool inst. AN SSSR. Vol.226. P.29-37 [in Russian].
- Holm A. 1973. On the spiders collected during the Swedish expedition to Novaya Zemlya and Yenisey in 1875 and 1876// Zool Scripta. Vol.2. No.3. P.71-110.
- Holzmayer O.P. 1934. [Materials to the spider fauna of Chuvashian Republic]// Uch. zap. Kazansk. univ. Vol.94. No.4. P.104-142 [in Russian].
- Izmailova M.V. 1975. [Fauna and ecology of spiders of Verkhnee Priangarye]. Autoreferate of the Thesis on Candidate of Biological Sci. Degree. Irkutsk. 15 pp. [in Russian].
- Izmailova M.V. 1989. [Spider fauna of the south of Eastern Siberia]. Irkutsk, Irkutsk State Univ. Press. 182 pp. [in Russian].
- Izmailova M.V., Verzhutskiy B.N. 1981. [Araneocomponent (Arthropoda, Aranei) of taiga geosystems]// Fauna i ekologiya nazemnykh chlenistonogikh Sibiri. Irkutsk. P.115-130 [in Russian].
- Kalinkin V.M. 1985. [Influence of useful arthropods on harmful lepidopteran numbers in apple orchards of Transcarpathia]// Byull. VIZR. No.59. P.11-18 [in Russian].
- Kaplin V.G. 1978. [Complexes of soil invertebrates of sandy deserts of the southern subzone (Eastern Karakum taken as an example). Ashkhabad, Ylym Publ. 160 pp. [in Russian].
- Karsch F. 1879. Baustoffe zu einer Spinnenfauna von Japan// Verh. naturh. Ver. preuss. Rheinl. Bd.36. S.57-105, pl.I.
- Kauri H. 1938. Ämblikulisi kevadises faunaaспектis Tarumaa-Jöepere ümbruses Virumaal// Miscellaneous I. Acta Inst. Mus. Zool. Univ. Tartuensis. No.2. P.47-48.
- Kharitonov D.E. 1923. [To the spider fauna of the eastern slope of the Urals]// Izv. Biol. NII pri Permsk. Univ. Vol.2. No.1. P.25-36 [in Russian].
- Kharitonov D.E. 1926a. [Materials to the spider fauna of Perm Gouvernement]// Ezhegodn. Zool. Muz. AN SSSR za 1925. P.103-136, tab.V [in Russian].
- Kharitonov D.E. 1926b. [Materials to the spider fauna of Verkhoturskiy Ural]// Izv. Biol. NII pri Permsk. Univ. Vol.5. Вып.2. P.49-60, 1 tab. [in Russian].
- Kharitonov D.E. 1928. [Materials to Araneina fauna of Leningrad Gouvernement]// Ibidem. Vol.6. No.1. P.1-8 [in Russian].
- Kharitonov D.E. 1932. Katalog der russischen Spinnen. Leningrad, AN SSSR Publ. 206 pp. [in both German and Russian].
- Kharitonov D.E. 1936. [Addition to the catalogue of Russian spiders]// Uch. zap. Permsk. Univ. Vol.2. No.1. P.167-225 [in Russian].
- Kharitonov[Charitonov] D.E. 1937. Contribution to the fauna of Crimean spiders// Festschr. Strand. Bd.3. S.127-140.
- Kharitonov D.E., Berezneva N.A. 1970. [Materials to the spider fauna of "Preduralye" Uchleskhoz]// Ibidem. No.206. P.195-200 [in Russian].
- Kirilenko V.A., Legotay M.V. 1981. [To the study of Aranei fauna in eastern forest-steppe of Ukraine]// Fauna i ekologiya nasekomykh. Perm, Perm State Univ. Press. P.45-54 [in Russian].
- Koch L. 1866. Die Arachniden-Familie der Drassiden. H.6. Nürnberg. S.257-304, Tab.XI-XII.
- Koch L. 1878. III. Kaukasische Arachnoideen// O.Scheider. Naturwissenschaftliche Beiträge zur Kenntnis der Kaukasusländer. Dresden, Sitzber. Isis (Beiträge). S.36-71.
- Koch L. 1879. Arachniden aus Sibirien und Novaja Semlja eingesammelt von der schwedischen Expedition im Jahre 1875// Kongl Svenska Vet.-Akad. Handl. Bd.16. H.5. S.1-136, Taf.I-VII.
- Kokhmanyuk F.S., Yaroshchuk L.S. 1981. [To the question of the spider and myriapod fauna of Belovezhskaya Pushcha]// Zhivotniy mir Belorusskogo Polesya, okhrana i ratsionalnoe ispolzovanie. Vtoraya itogovaya nauchnaya konferentsia. Gomel. P.11-12 [in Russian].
- Kolosvary G. von. 1939. Neue Beiträge zu Ungarn-Spinnenfauna// Zool. Anz. Bd.126. H.7/8. S.205-207.
- Krasnobaev Yu.P. 1983. [To the spider fauna of Zhiguli State Reserve]// Problemy ratsionalnogo ispolzovaniya i okhrany prirodного kompleksa Samarskoy Luk. Kuybyshev. P.83-86 [in Russian].
- Krasnobaev Yu.P. 1992. [Spiders of stony steppes of Zhiguli State Reserve]// Fauna i ekol. paukov, skorpionov i lozhnoskorponov SSSR. Trudy Zool inst. AN SSSR. Vol.226. P.83-90 [in Russian].
- Krasnobaev Yu.P., Ovcharenko V.I. 1986. [To the question of the spider fauna (Aranei) of Kuybyshev Area]// Ekologicheskie issledovaniya v Sredнем Povolzhье. Kuybyshev. P.89-98 [in Russian].
- Kroneberg A.I. 1875. [A.P.Fedchenko. Voyage in Turkestan Spiders Araneae]// Izvestiya Obshchestva lyubiteley estestvoznanii, antropologii i etnografii. Vol.19. No.3. P.I-IV, 1-55, 5 tab. [in Russian].
- Kulczynski W. 1885. Araneae in Camtschadalia a Dre B.Dybowski collectae// Pam.Wydz. mat-prz. Akad. Umiej. T.11. P.1-60, 1-6, Tab.IX-XI.
- Kulczynski W. 1901. Arachnoidea// Dritte Asiatische Forschungreise des Grafen Eugen Zichy. Bd.II. Zoologische Ergebnisse. S.313-369, Tab.XII-XIII.
- Kulczynski V. 1908. Araneae et Oribatidae expeditionum Rossicarum in insulas Novo-Sibiricas annis 1885-1886 et 1900-1903 susceptarum// Mem. Acad. Imp. Sci. St.-Petersbourg. Ser.VIII. Vol.18. No.7. P.I-IV, 1-97, PLI-III.
- Kulczynski W. 1913. Arachnoidea// Velitchkovsky. Faune de district de Walouky du gouvernement de Woronege (Russie). Pt.10. P.1-30.
- Kulczynski W. 1916. Araneae Sibiriae Occidentalis Arcticae// Mem. Acad. Imp. Sci. Petrograd. Ser.VIII. Vol.28. No.11. P.1-44, Tab.I-II.
- Kulczynski W. 1926. Arachnoidea camtschadalica// Annu. Zool. Mus. AN URSS. Vol.27. No.1. P.29-72, Tab.II-III.
- Lavrov M.T. 1968. [Fauna of forest soils and the ways of its management]. Moscow, Lesnaya Promyshlennost. 131 pp. [in Russian].
- Legotay M.V. 1958. [Some data on the arachnofauna of Transcarpathia]// Doklady i soobshcheniya Uzhgorodsk. Univ. Ser.biol. No.2. P.27-30 [in Russian].
- Legotay M.V. 1959. [Additional data on the arachnofauna of Transcarpathia]// Ibidem. No.3. P.53-56 [in Russian].
- Legotay M.V. 1973. [Spiders of the Ukrainian Carpathians]. Autoreferate of the Thesis of Candidate of Biological Sci. Degree. Kharkov. 21 pp. [in Russian].
- Legotay M.V. 1980. [Spiders (Aranei) on wheat fields of Transcarpathia]// Entomofagi vrediteley rasteniy. Kishinev. P.28-33 [in Russian].
- Legotay M.V. 1989. [Materials to the spider fauna (Arachnida, Aranei) of Transcarpathia]// Fauna i ekologiya paukov i skorpionov. Arakhnologicheskiy sbornik. Moscow, Nauka Publ. P.16-30 [in Russian].
- Legotay M.V., Sekerskaya N.P. 1982. [Spiders in orchards]// Zashchita rasteniy. No.7. P.48-51 [in Russian].
- Legotay M.V., Sekerskaya N.P. 1989. [Spiders]// Poleznaya fauna plodovogo sada. Spravochnik. Moscow, Agropromizdat Publ. P.218-227, 294-297 [in Russian].
- Legotay M.V., Tarasyuk G.D. 1964. [Ecological distribution of the arachnofauna of Ciscarpathia]// Ekologiya nasekomykh i drugikh nazemnykh bespozvonochnykh Sovetskikh Karpat. Uzhgorod. P.54-59 [in Russian].
- Litvinova A.N. 1978. [Fauna and numbers of crown-dwelling spiders]// Biologicheskie osnovy osvoeniya, rekonstruktsii i okhrany zhivotnogo mira Belorussii. Tezisy Chetvertoy

- zoologicheskoy konferentsii Belorusskoy SSR Minsk. P.180-181 [in Russian].
- Litvinova A.N., Shlyakhtyonok A.S., Ovtsharenko V.I. 1980. [On the spider fauna (Aranei) of Byelorussia]. Deposited in VINITI. No.3114-80. 11 pp. [in Russian].
- Litvinova A.N., Shlyakhtyonok A.S., Ovtsharenko V.I. 1981. [Stratification of spiders in pine forests]// Vestsi AN BSSR Ser. biyaL. N. No.6. P.98-102, 143 [in Byelorussian].
- Litvinchuk L.N. 1980. [Alder leaf-rolling sawfly]. Novosibirsk, Nauka Publ., Siberian Division. 48 pp. [in Russian].
- Lukianov N. 1897. [List of spiders (Araneina, Pseudoscorpionina and Phalangina) living in the South-Western Province and adjacent gouvernements of Russia]// Zapiski Kievskogo obshchestva estestvoispytateley. Vol.14. No.2. P.1-19 [in Russian].
- Marusik Yu.M. 1984. [To the spider fauna of "Les na Vorskle" State Reserve]// Fauna i ekologiya paukoobraznykh. Perm, Perm State Univ. Press. P.132-138 [in Russian].
- Marusik Yu.M. 1992. [Zoogeographical peculiarities of the Upper Kolyma araneofauna]// Fauna i ekol. paukov, skorpionov i lozhnoskorionov SSSR. Trudy Zool. inst. AN SSSR. Vol.226. P.125-127 [in Russian].
- Matsuda M. 1991. Two newly recorded species of the genus *Clubiona* (Araneae: Clubionidae) from Japan// Atypus. No.97. P.22-26.
- Matsyuk V.A., Yanyuk V.A., Verlan E.T. 1984. [Materials to the study of the spider fauna (Aranei) in Moldavian orchards]// Entomofagi vrediteley rasteniy. [Part 2]. Kishinev. P.7-13 [in Russian].
- Mikhailov K.G. 1983a. [Spiders (Arachnida, Aranei) of forest litter of Zvenigorod Biostation of Moscow State University]// Fauna i ekologiya pochvennykh bespozvonochnykh Moskovskoy oblasti. Moscow, Nauka Publ. P.52-67 [in Russian].
- Mikhailov K.G. 1983b. [Catalogue of spiders (Arachnida, Aranei) of Moscow Area]// Ibidem. P.67-85 [in Russian].
- Mikhailov K.G. 1989. [Materials to the spider fauna (Arachnida, Aranei) of the territory between Volga and Kama Rivers]// Fauna i ekologiya paukov i skorpionov: Arakhnologicheskiy sbornik. Moscow, Nauka Publ. P.52-57 [in Russian].
- Mikhailov K.G. 1990a. The spider genus *Clubiona* Latreille 1804 in the Soviet Far East, 1 (Arachnida, Aranei, Clubionidae) // Korean Arachnol. Vol.5. No.2. P.139-175.
- Mikhailov K.G. 1990b. The spider genus *Clubiona* Latreille 1804 in the Caucasus, USSR (Arachnida: Araneae: Clubionidae) // Senckenberg. biol. Bd.70. H.4/6.S.299-322.
- Mikhailov K.G. 1991. The spider genus *Clubiona* Latreille 1804 in the Soviet Far East, 2 (Arachnida, Aranei, Clubionidae) // Korean Arachnol. Vol.6. No.2. P.207-235.
- Mikhailov K.G. 1992. [Spiders of the genus *Clubiona* Latreille 1804 (Aranei Clubionidae) of the Soviet Union (faunistic review)]// Fauna i ekol. paukov, skorpionov i lozhnoskorionov SSSR. Trudy Zool. inst. AN SSSR. Vol.226. P.60-69 [in Russian].
- Mikhailov K.G., Fet V.Ya. 1986. [Materials to the spider fauna (Aranei) of Turkmenia. I. Families Anyphaenidae, Sparassidae, Zoridae, Clubionidae, Micariidae, Oxyopidae]// Fauna, sistematika i filogeniya bespozvonochnykh zhivotnykh. Sbornik trudov Zool. Muzeya MGU. Vol.24. P.168-186 [in Russian].
- Mikityuk V.F. 1982. [Spiders in agroecosystems of Northern Cis-Chernomoria]// Formirovanie zhivotnogo i mikrobnogo naseleniya agrotsenozov. Tezisy dokladov Vsesoyuznogo soveshchaniya, Pushchino, 14-16 sent. 1982. P.66-67 [in Russian].
- Miller F. 1971. Pavouci - Araneida// Klič Zviřeny ČSSR. Praha. Vol.4. P.51-306.
- Minoranskiy V.A. 1988. [Materials to the spider fauna (Aranei) of Checheno-Ingushetia]// Fauna i ekologiya paukoobraznykh. Perm, Perm State Univ. Press. P.34-42 [in Russian].
- Minoranskiy V.A., Gramotenko V.P., Ponomarev A.V. 1977. [Some data on the distribution of spiders in Rostov Area]// Voprosy arakhnoentomologii. Perm: Perm State Univ. Press. P.92-105 [in Russian].
- Minoranskiy V.A., Ponomarev A.V. 1984. [Materials to the spider fauna of Kalmykia]// Fauna i ekologiya paukoobraznykh. Perm, Perm State Univ. Press. P.82-92 [in Russian].
- Minoranskiy V.A., Ponomarev A.V., Gramotenko V.P. 1981. [On spiders of populated areas]// Fauna i ekologiya nasekomykh. Perm, Perm State Univ. Press. P.33-44 [in Russian].
- Minoranskiy V.A., Ponomarev A.V., Slyusarev V.V., Gramotenko V.P. 1984. [To the spider fauna (Aranei) of Checheno-Ingushetia]// Izv. Severo-Kavkazsk. nauchn. tsentra vyssh. shkoly. Estestv. nauki. No. 4. P.76-81 [in Russian].
- Mkheidze T.S. 1964. [Spiders]// Zhivotniy mir Gruzii. Vol.2. Chlenistonogie. Tbilisi, AN GruzSSR Publ. P.48-116 [in Georgian].
- Nordmann A. von. 1863. Erstes Verzeichniss der in Finnland und Lapland bisher gefundenen Spinnen, Araneae// Bidr. Finl. Naturkannedom. Vol.8. P.1-39.
- Nosek A. 1895. Seznam českých a moravských Pavouků// Vest. Kral. české spol. náuk. No.3. P.1-56.
- Nowicki M. 1870. Zapiski fauniczne// Spraw. kom. fizyogr. Krakow. Vol.4. P.1-30.
- Nowicki M. 1874. Dodatek do fauny pajęczaków Galicyi// Ibidem. Vol.8. P.1-11.
- Odenval E. 1901. Araneae nonnulae Sibiriae Transbaicalensis// Ofvers. Finska Vet.-Soc. Forhandl. Vol.43. P.255-273, 1 tab.
- Olshvag V.N., Malozymov Yu.A. 1987. [Hortobionous arthropod population in montaneous tundra of the southern Urals]// Fauna i ekologiya nasekomykh Urala. Sverdlovsk, Ural State Univ. Press. P.121-130 [in Russian].
- Ovsyannikov A.G. 1937. [To the spider fauna of Kursk Area]// Uch. zap. Permsk. Univ. Vol.3. No.1. P.89-93 [in Russian].
- Paik K.Y. 1990a. Korean spiders of the genus *Clubiona* (Araneae: Clubionidae). I. Description of eight new species and five undescribed records from Korea// Korean Arachnol. Vol.5. No.2. P.85-129.
- Paik K.Y. 1990b. Korean spiders of the genus *Clubiona* (Araneae: Clubionidae) II. On the clubionid spiders reported from Korea before the report I// Ibidem. Vol.6. No.1. P.63-89.
- Pakhorukov N.M. 1977. [Materials of the spider fauna of the eastern slope of northern Urals]. Deposited in VINITI No. 4208-77. 25 pp. [in Russian].
- Pakhorukov N.M. 1979. [Spiders of northern Urals (eco-faunistic review). Autoreferate of Thesis of Candidate of Biological Sci. Degree. Leningrad. 22 pp. [in Russian].
- Pakhorukov N.M. 1980. [Spiders of Yaksha profile]// Vzaimosvyazi komponentov lesnykh i bolotnykh ekosistem. Leningrad, Nauka Publ. P.160-171 [in Russian].
- Pakhorukov N.M. 1985. [Characteristics of spider complexes of forest-steppe biocoenoses of southern Transurals]// Fauna i ekologiya paukov SSSR. Trudy Zool. Inst. AN SSSR. Vol.139. P.92-98 [in Russian].
- Pakhorukov N.M., Efimik V.E. 1988. [To the spider fauna of Bashkirian State Reserve]// Fauna i ekologiya paukoobraznykh. Perm, Perm State Univ. Press. P.15-34 [in Russian].
- Pakhorukov N.M., Polyanin A.B. 1987. [Materials to the spider fauna of Ilmen State Reserve]// Fauna, ekologiya bespozvonochnykh zhivotnykh Chelyabinskoy oblasti. Sverdlovsk, Ural Div. of USSR Academy of Sci. P.13-21 [in Russian].
- Panteleeva N.Yu. 1982. [To the study of spiders of "Galichya

- Gora" State Reserve]// Issled. rastit. i zhivot. mira zapoved. Galichya Gora. Voronezh. P.89-92 [in Russian].
- Peelle M.L., Saito S. 1932. Spiders from the Southern Kurile Islands. I. Araneidae from Iturup// J. Fac. Sci. Hokkaido Imp. Univ. Ser.6. Vol2. No.2. P.83-96.
- Pereleshina V.I. [Pereleshina W.] 1927. Beitrag zur Kenntnis der Spinnenfauna von Askania Nova// Trav. Mus. Zool Acad. Sc. Ukraine. No.3. P.53-56.
- Pereleshina V.I. 1928a. [Spider fauna of the environs of Bolshevo Biostation]// Zapiski biostantsii v Bolsheve. No.2. P.1-74, 1 tab. [in Russian].
- Pereleshina V.I. 1928b. [Materials to the study of the spider fauna of Byelorussia]// Matali da vyvuchennya flery i fauny Belarusi. Minsk. Vol2. P.34-48 [in Byelorussian].
- Pereleshina V.I. 1930. [Materials to the spider fauna of western and southwestern parts of Eastern Europe]// Ezheg. ZoolMuz. AN SSSR. Vol31. No.3-4. P.359-391 [in Russian].
- Pichka V.E. 1965. [On the ecology of spiders of Central forest-steppe]// Zool. Zhurn. Vol44. No.4. P.527-536 [in Russian].
- Pichka V.E. 1967. [Observations on spiders of Voronezh State Reserve]// Trudy Voronezh State Reserve. No.15. P.86-93 [in Russian].
- Pichka V.E. 1974. [On the fauna and ecology of spiders of environs of Kanev (forest-steppe of USSR)]// Vestn. zool. No.6. P.23-30 [in Russian].
- Pichka V.E. 1977. [About insect parasites of spiders, their cocoons and nests]// Vestn. zool. No.3. P.92-93 [in Russian].
- Pichka V.E. 1984a. [To the spider fauna of Central-Chernozym State Reserve]// Fauna i ekologiya paukoobraznykh. Perm, Perm State Univ. Press. P.68-77 [in Russian].
- Pichka V.E. 1984b. [To the fauna and ecology of spiders of Central-Chernozym State Reserve]// Ekologo-faunisticheskie issledovaniya Tsentralnoy lesostepi Evropeyskoy chasti SSSR. Moscow. P.65-75 [in Russian].
- Pichka V.E., Skufin K.V. 1981. [Additions to the spider fauna of Central forest-steppe]// Vestn. zool. No.6. P.7-15 [in Russian].
- Pokrovskiy S.V. 1925. [Spider fauna of Klyazma River Basin]// Zapiski biostantsii v Bolsheve. No.1. P.19-25 [in Russian].
- Pokrovskiy S.V. 1930. [Spiders of Belskiy Uezd of Smolensk Gouvernement]// Trudy obshchestva izucheniya prirody Smolenskogo kraja. Vol5. P.115-128 [in Russian].
- Polchaninova N.Yu. 1988. [Spiders of Chernomorian State Reserve]// Fauna i ekologiya paukoobraznykh. Perm, Perm State Univ. Press. P.42-51 [in Russian].
- Polchaninova N.Yu. 1992. [Spiders of Provalskaya Steppel]// Fauna i ekol. paukov, skorpionov i lozhnoskorpionov SSSR. Trudy Zool. inst. AN SSSR. Vol226. P.98-104 [in Russian].
- Polchaninova N.Yu., Astakhova E.V. 1984. [To the study of the araneofauna of riparian biocenoses of the south of the forest-steppe zone of Levoberezhnaya Ukraine]// Vestn. Kharkovsk. Univ. No.262. P.85-86 [in Russian].
- Polishchuk V.V., Travyanko V.S., Tavolzhanova T.I., Kobenek G.V. 1980. [Invertebrate fauna of water bodies of Pripyatian Polesye and its changes under the influence of amelioration]// Issled. po entomol. i akarol. na Ukraine. Tez. dokl. Vtorogo syezda Ukr. Entomol. Obshch. Uzhgorod, 1980. Kiev. P.55-56 [in Russian].
- Polozhentsev P.A., Akimtseva N.A. 1980. [Spiders (Aranei) of forest biocenoses of Transcarpathia]// Entomol. Obozr. Vol59. No.2. P.448-450 [in Russian].
- Polyanin A.B., Pakhorukov N.M. 1988. [Spider fauna of forest zone of southern Transuralia]// Fauna i ekologiya paukoobraznykh. Perm, Perm State Univ. Press. P.61-80 [in Russian].
- Polyanin A.B., Pakhorukov N.M. 1992. [Spiders the of herb-undershrub stratum of some biocenoses of Ilmen State Reserve]// Fauna i ekol. paukov, skorpionov i lozhnoskorpionov SSSR. Trudy Zool. inst. AN SSSR. Vol226. P.12-18 [in Russian].
- Ponomarev A.V. 1981. [Eco-faunistical characteristics of spiders of the semidesert zone of USSR European part]. Autoreferate of the Thesis of Candidate of Biological Sci. Degree. Leningrad, State Univ. 26 pp. [in Russian].
- Prieditis A.P., Sternbergs M.T. 1981. [Spider fauna (Aranei) in apple-trees agrobiocenoses]// Zashchita rasteniy ot vrediteley i bolezney. Trudy Latviyskoy selskokhoz. akad. Elgava. No.188. P.9-12 [in Russian].
- Prieditis A.P., Sternbergs M.T. 1982. [Lower level of activity of some spider species]// Borba s vredit. i boleznyami kartofelya, plodoovoshchnykh i polevykh kultur. Trudy Latviyskoy selskokhoz. akad. Elgava. No.200. P.25-28 [in Russian].
- Pristavko V.P., Zhukovets E.M. 1987. [Spiders (Aranei) as the subject of ecological monitoring in Berezina State Reserve]// Entomol. Obozr. Vol66. No.1. P.184-189 [in Russian].
- Pyatin M.A. 1988. [Species composition, numbers and biotopical distribution of spiders in Penza Area]// Fauna i ekologiya paukoobraznykh. Perm, Perm State Univ. Press. P.80-87 [in Russian].
- Rosca A. 1930. Contributiuni la conoasterea Arachnidelor din Bukovina// Bul. Fac. sti. Cernauti. Vol4. No.2. P.201-219.
- Rosca A. 1936. Fauna Araneelor din Bukovina (sistematica, ecologia si raspandirea geographică) // Ibidem. Vol10. P.123-216.
- Saito S. 1932. Descriptions of two new species of Araneida from Northern Kurile Islands// Trans. Sapporo Nat. Hist. Soc. Vol12. No.2-3. P.100-104.
- Saito S. 1933. Araneida from the northern Kurile islands, with descriptions of two new species// Bull. Biogeogr. Soc. Jap. Vol4. No.2. P.122-132.
- Saito S. 1934. A supplementary note on spiders from southern Saghaliens, with descriptions of two new species// Trans. Sapporo Nat. Hist. Soc. Vol13. No.3. P.326-340.
- Savelieva L.G. 1970. [Fauna and zoogeographical connections of spiders of East-Kazakhstan Area]// Biologiya i geografiya. Alma-Ata. No.6. P.78-88 [in Russian].
- Savelieva L.G. 1972a. [New species and subspecies of Clubionidae (Aranei) from East-Kazakhstan Area]// Zool. Zhurn. Vol51. No.9. P.1404-1407 [in Russian].
- Savelieva L.G. 1972b. [Ecologico-faunistical characteristics of spiders of East-Kazakhstan Area]. Autoreferate of the Thesis of Candidate of Biological Sci. Degree. Leningrad, State Univ. 23 pp. [in Russian].
- Savelieva L.G. 1976. [Zonal-stational distribution of spiders in the territory of East-Kazakhstan Area]// Biol. nauki. Alma-Ata, Kazakhstan. Pedagog. Inst. 1974. No.1. P.50-54 [in Russian].
- Savelieva L.G. 1979. [Zoogeographical complexes of spiders (Aranei) from Eastern Kazakhstan]// Priroda i khozyaystvo Vostochnogo Kazakhstana. Alma-Ata. P.139-148 [in Russian].
- Schenkel-Haas E. 1924. 11. Araneina und Opiliones// Dampf A. Zur Kenntnis der estlandischen Hochmoorfauna (II. Beitrag) // Sitz.-Ber. naturf. Ges. Dorpat. Bd.31. H.1-2. S.57-65.
- Schenkel E. 1930. Die Araniden der schwedischen Kamtschatka-Expedition 1920-1922// Ark. f. Zool. Bd.21A. No.15. P.1-33.
- Sharov A.A., Izhevskiy S.S., Prokofyeva E.A., Mikhailov K.G. 1984. [Spiders as predators of *Hybantria cunea* in the south of USSR European part]// Zool. Zhurn. Vol63. No.3. P.392-398 [in Russian].
- Sharova I.Kh. 1970. [Soil mesofauna of leaved forests in Podmoskovye]// Fauna i ekol. zhivotnykh. Uch. zap. Mosk. gos. ped. inst. No.394. P.3-20 [in Russian].

- Shlykov N.B. 1975. [Biogeographical characteristics of spiders of Tura Plain]// Nauch. tr. Tyumensk. Univ. No.16. P.46-52 [in Russian].
- Shlykov N.B. 1978. [Araneological characteristics of three biotopes of Tobol River floodplain]// Ekol. zhivotnykh i faunistika. Tyumen. P.41-46 [in Russian].
- Shlyakhtyonok A.S. 1986. [Spider fauna (Aranei) of meadow biogeocenoses of Berezina Biosphere Reserve]. Deposited in VINITI 08.04.86. No.2499-B. 11 pp. [in Russian].
- Siemaschko J.M. 1861. Verzeichniss der in der Umgegend v. St. Petersburg vorkommenden Arachniden// Horae Soc. Ent. Ross. Vol.1. P.117-137.
- Silvere A.-P. 1960. Must-kärbsenäpi (*Muscicapa hypoleuca* Pall.) nidikolide faunast. I// Loodusuuri jate Seltsi Aastaraamat. K.53. L.170-187.
- Simon E. 1891. Liste des Arachnides récueillis par Rabot dans la Sibérie occidentale en 1890// Bull. Soc. Zool. Fr. Vol.16. P.107-109.
- Simon E. 1899. Araneeae Caucasicae// Ed. G.I.Radde. Kollektii Kavkazskogo Muzeya (Museum Caucasicum). Vol.1. Zoologiya. Tiflis. P.477- 478 [in Russian].
- Song D.X. 1987. Spiders from agricultural regions of China. Beijing, Agricultural Publ. House. 376 pp.
- Sorokin N.S. 1977. [Entomophages of the potato beetle (*Leptinotarsa decemlineata* Say) and their influence on the numbers of pests in Rostov Area]. Autoreferate of the Thesis of Candidate of Biological Sci. Degree. Leningrad. 25 pp. [in Russian].
- Spassky S. 1912. Die Spinnen des Don-Gebietes// Zool. Anz. Bd.40. H.6-7. S.179-188.
- Spassky S.A. 1914. [Spiders of Don Area. II]// Izv. Donsk. politekhn. inst. Novocherkassk. Div.2. Vol.3. No.2. P.85-97 [in Russian].
- Spassky S.A. 1919. Die Spinnen des Dongebietes. II// Zool. Anz. Bd.50. H.6-7. S.147-159.
- Spassky S.A. 1925. [An identification book of spiders of Don Area]. Novocherkassk, "Znanie" Publ. 62 pp., 2 tab. [in Russian].
- Spassky S.A. 1927. [Materials to the spider fauna of Tauric Gouvernement]// Izv. Donsk. inst. selsk. khoz. i melior. Vol.7. P.66-80 [in Russian].
- Spassky S.A. 1937. [Materials to the spider fauna of Black Sea coast]// Sborn. nauch.-issled. rabot Azovo-Chernomorsk. selskokhoz. inst. No.5. P.131-138 [in Russian].
- Spassky S.A. 1940. [Spiders]// Priroda Rostovskoy oblasti. Rostov-na-Donu, Rostizdat Publ. P.193-202 [in Russian].
- Spassky S.A. 1952. [Spiders of Turanian zoogeographical province]// Entomol. Obozr. Vol.32. P.192-205 [in Russian].
- Spassky S.A., Lavrov S.D. 1928. [Materials to the spider fauna of Western Siberia]// Trudy Sibirsk. inst. selsk. khoz. i lesovedstva. Vol.10. No.2. P.203-213 [in Russian].
- Spassky S.A., Shnitnikov V.N. 1937. [Materials to the spider fauna of Kazakhstan]// Materialy po vredityam zhivotnovodstva i faune preimushchestvenno Yuzhnogo Kazakhstana. Trudy Kazakhsk. filiala AN SSSR. No.2. P.264-300 [in Russian].
- Sternbergs M.T. 1977. [Materials to the spider fauna (Aranei) of Stolby State Reserve]// Trudy zapov. "Stolby". No.11. P.87-90 [in Russian].
- Sternbergs M.T. 1979. [Structure and dynamics of the spider fauna of Aegopodio-Tiliatum litter of "Moritssala" Reserve]// Tr. Zool. inst. AN SSSR. Vol.85. P.54-59 [in Russian].
- Sternbergs M.T. 1981a. Materialia par Latvijas zirneklu faunu. V. Dzimta Clubionidae// Latv. Ent. No.24. P.56-59 [in Latvian].
- Sternbergs M.T. 1981b. Moricsalas rezervata melnaksnu garsas zemsedzes zirneklu// Mezsaimnieciba un Mezrupmeciba No.3. P.40-42 [in Latvian].
- Sternbergs M.T. 1981c. [Materials to the spider fauna (Aranei) of Barguzin State Reserve]// Fauna i ekol. nazemnykh chlenistonogikh Sibiri. Irkutsk. P.130-133 [in Russian].
- Sternbergs M.T. 1982. [Spiders (Aranei) of oak forest litter of "Moritssala" Reserve]// Izuchenie okhranyaemykh prirodn. terr. Latv. SSR. Riga, Zinatne Publ. P.86-88 [in Russian].
- Sternbergs M.T. 1983. [Spiders (Chelicera, Aranei)]// Prirod. rezervat Moritssala. Flora i fauna. Riga, Avots Publ. P.41-47 [in Russian].
- Sternbergs M.T. 1985. [Spiders (Aranei) of soil cover and litter of Latvian pine forests]// Fauna i ekologiya paukov SSSR. Trudy Zool. Inst. AN SSSR. Vol.139. P.141-146 [in Russian].
- Sternbergs M.T. 1988. [Materials to the spider fauna of Primorye Province]// Fauna i ekologiya paukoobraznykh. Perm, Perm State Univ. Press. P.92-97 [in Russian].
- Strand E. 1906. Die arktischen Araneeae, Opiliones und Chernes// Fauna Arctica. Bd.4. H.3. S.433-478.
- Strand E. 1907. Vorläufige Diagnosen süd- und ostasiatischer Clubioniden, Ageleiden, Pisauriden, Lycosiden, Oxyopiden und Salticiden// Zool. Anz. Bd.31. H.17-18. S.558-570.
- Strand E. 1909. Süd- und ostasiatische Spinnen. II// Abh. naturforsch. Ges. Gorlitz. Bd.26. S.1-128.
- Sytshhevskaja V.I. 1935. Etude sur les Araignées de la Kamtchatka// Folia zool. hydrobiol. Vol.8. No.1. P.80-103.
- Talanov V.P., Nazarenko S.V. 1989. [To the spider fauna of Lower Dnepr]// Vses. soveshch. po probleme kadastra i uchyota zhivotn. mira. Tez. dokl. Chast 4. Opyt kadastrovoy kharakteristiki, materialy k kadastru po bespozvonochnym zhivotnym. Ufa, Bashkirian Press. P.262-264 [in Russian].
- Talitskiy V.I., Legotay M.V. 1982. [Spiders (Aranei) in fruit plantations of Moldavia]// Fauna i ekol. entomofagov. Kishinev. P.25-31 [in Russian].
- Tanasevitch A.V. 1985. [To the study of spiders (Aranei) of the Polar Urals]// Fauna i ekologiya paukov SSSR. Trudy Zool. Inst. AN SSSR. Vol.139. P.52-62 [in Russian].
- Tarabaev Ch.K. 1979. [Spiders as apple-tree crown-dwellers in piedmonts of Zailiyskiy Alatau Mt.Ridge]// Novosti entomol. Kazakhstana. Trudy Kazakhsk. otd. Vses. entomol. obshch. Deposited in VINITI No.3415-79. P.119-125 [in Russian].
- [Terrestrial invertebrates of recreational spruce forests of Podmoskovye]. 1989./ Ed. by B.R.Striganova. Moscow, Nauka Publ. 224 pp. [in Russian].
- Thorell T. 1875. Verzeichnisse Südrussischer Spinnen// Horae Soc. Ent. Ross. T.11. P.39-122.
- Tolstova Yu.S., Atanov N.I. 1982. [Effect of chemical pesticides on the arthropod fauna of a fruit orchard. I. Long-term effect of pesticides on an agroecosystem]// Entomol. Obozr. Vol.61. No.3. P.441-453 [in Russian].
- Tsellarius A.Yu., Shorokhov V.V. 1985. [Species composition and biogeographical distribution of herpetobiontic spiders of Kivach State Reserve (Karelian ASSR)]// Fauna i ekologiya paukov SSSR. Trudy Zool. Inst. AN SSSR. Vol.139. P.84-91 [in Russian].
- Tyshchenko V.P. 1971. [An identification book of spiders of the European part of USSR]// Opredeliteli po faune SSSR, izdavaemye Zool. inst. AN SSSR. Leningrad, Nauka Publ., Leningrad Division. No.105. 281 pp. [in Russian].
- Utochkin A.S. 1971. [To the spider fauna of Astrakhan State Reserve]// Uch. zap. Permsk. Univ. No.249. P.154-157 [in Russian].
- Utochkin A.S. 1977. [Spiders of Saratovo Forestry of Volga-Kama State Reserve]// Voprosy arakhnoentomologii. Perm, Perm State Univ. Press. P.69-80 [in Russian].
- Utochkin A.S., Golovashkina L.N. 1977. [To the fauna and

- distribution of spiders among phytocenoses of Troitskiy Reserve]// Ibidem. P.81-91 [in Russian].
- Utochkin A.S., Pakhorukov N.M. 1976. [Materials to the spider fauna of the Reserve]// Trudy Pechoro-Ilychskogo zapov. VoL13. P.78-88 [in Russian].
- Uzenbaev S.D. 1985. [Fauna and ecology of spiders (Aranei) of a dried mesotrophic bog biocenosis]// Nasekomye i fitopatogennye grify v lesnykh ekosistemakh. Petrozavodsk. P.81-110 [in Russian].
- Uzenbaev S.D. 1986. [To the spider fauna (Aranei) of Karelia]// Fauna i ekologiya chlenistonogikh Karelii. Petrozavodsk. P.7-28 [in Russian].
- Uzenbaev S.D. 1987. [Ecology of predatory arthropods of a mesotrophic bog]. Petrozavodsk, Karelian Filial of USSR Acad. Sci. 128 pp. [in Russian].
- Vagner V.A. 1892. Araneina// Dwigubsky. Primitiae Faunae Mosquensis [Second Ed.]. Opyt kataloga predstaviteley Moskovskoy fauny. M. P.117-120 [in Russian].
- Vagner V.A. 1895. [List and description of the collection on the biology of spiders]// Kollektii Zool. Kab. Warshawsk. Univ. Vol4. P.1-20, tab.I-II [in Russian].
- Vaitskute Ya.I. 1960. [Spiders (Araneida) of young pine forests of the S part of Lithuanian SSR]// Trudy AN LitSSR Ser.C. No.3 (23). P.133-144 [in Lithuanian].
- Vaitskute Ya.I. 1963. [Analysis of the spider distribution using Student's criterium]// Ibidem. No.1. P.65-78 [in Russian].
- Verzhbitskiy E. 1902. [On the spiders of Caucasian Province]// Zapiski Kievskogo obshchestva estestvoispytateley. Vol17. No.2. P.461-504, tab.VI [in Russian].
- Verzhutskiy B.N., Bessolityna E.P., Seryshev A.A. 1985. [Cadrat of araneopopulation of kettles of Stanovoy Mt.Ridge]// Nazemnye chlenistonogie Sibiri i Dalnego Vostoka. Irkutsk, Irkutsk Univ. Press. P.117-134 [in Russian].
- Veselova E.M. 1986. [Structure of invertebrate population of the herb-undershrub layer]// Kologrivskiy les (ekologicheskie issledovaniya). Moscow, Nauka Publ. P.88-110 [in Russian].
- Veselova E.M., Mikhailov K.G. 1986. [Structure of spider population (Arachnida, Aranei) of the herb-undershrub layer in phytocenoses of the territory between Volga and Kama Rivers]// Ekologiya (Sverdlovsk). No.6. P.46-51 [in Russian].
- Veselova E.M., Mikhailov K.G. 1987. [About the vertical structure of a spider population]// Ispolzovanie i okhrana resursov flory i fauny SSSR. Doklady MOIP 1985, Zoologiya i botanika. Moscow, Nauka Publ. P.83-85 [in Russian].
- Vilbaste A. 1957. Endla rabade Ämblikefauna (Endla rabamaastiku uurimistöödest) // Loodusuurijate Selsti aastaraamat. Tartu. Vol.50. L.115-118 [in Estonian].
- Vilbaste A. 1959. Avaste soo rohurinde ämblikufaunast// Entomol. kogumik. Tartu. Vol.1. L.29-36 [in Estonian].
- Vilbaste A. 1960. Ämblikke pmeie elamius// Eesti Loodus. No.6. L.339-340, tahr.XXI [in Estonian].
- Vilbaste A. 1964. Eesti ämblikefauna struktuurist ja sesoonsetest muutustest// ENSV Ted. Akad. toimetised. BiolSer. Vol13. No.4. L.284-301 [in Estonian].
- Vilbaste A. 1965. Suveaspekti ämblikefaunast kultuurniitude// Ibid. Vol14. No.3. L.329-337 [in Estonian].
- Vilbaste A. 1969. Eesti madalsoode ämblikefauna struktuurist ja sesoonsetest muutustest// Ibid. Vol18. No.4. L.390-407 [in Russian].
- Vilbaste A. 1972. Eesti rabade ämblikefauna struktuurist ja sesoonsetest muutustest// Ibid. Vol21. No.4. L.307-326 [in Estonian].
- Vilbaste A. 1973. Eesti siirdesoometsade ämblikefauna struktuurist ja sesoonsetest muutustest// Ibid. Vol22. No.3. L.210-225 [in Estonian].
- Vilbaste A. 1974. Väinamere laidude Ämblikefaunast// Loodusvaatlusi 1973. Tallin. Vol1. L.132-145 [in Estonian].
- Vilbaste A. 1980. The spider fauna of Estonian mires// ENSV Ted. Akad. toimetised. BiolSer. Vol29. No.4. L.313-327.
- Vilbaste A. 1987. Eesti ämblikud (Aranei) [Annoteeritud nimestik]. Tallinn, Valgus Publ. 178 lap. [in Estonian].
- Vysotskaya S.O. 1977. [Microbiocenosis of nests of (*Sciurus vulgaris carpathicus*)// Parazitol sbornik AN SSSR. Vol27. P.181-191 [in Russian].
- Vysotskaya S.O. 1978. [Analysis of biocenotic relations between the ectoparasites of *Microtus arvalis* Pall. and its nest-dwellers of E-Carpathians (Zakarpatskaya Area USSR)]// Ibidem. Vol28. P.73-95 [in Russian].
- Wajgiel L.P. 1868. Spis pajakow// Spraw. kom. fizyogr. Krakow. Vol2. P.153-155.
- Wajgiel L.P. 1874. Pajeczaki galicyjskie (Arachnoidea Haliciae). Kolomyia (Kolomea). 36 pp.
- Yaginuma T. 1957. Spiders from Hokkaido and Rishiri Island// Acta Arachnol. Vol14. No.2. P.51-61, plIII-IV.
- Zheltukhina V.I., Utochkin A.S. 1984. [Spiders of Raifa Forestry of Volga-Kama State Reserve]// Fauna i ekologiya paukobraznykh. Perm, Perm State Univ. Press. P.61-68 [in Russian].
- Zhukauskene Ya.I. 1966. [Soil invertebrates of Kurshu Nyaria Spit (7. Spiders)]// Trudy AN LitSSR. Ser.C. No.2 (40). P.193-199 [in Russian].
- Zhukauskene Ya.I. 1968. [Spiders (Araneida) of Lake Zhuvintas bank]// Zapovednik Zhuvintas. Vilnius, Mintis Publ. P.223-229 [in Russian].
- Zhukovets E.M. 1989. [To the spider fauna (Arachnida, Aranei) of Verkhneberezinskaya Lowland]// Fauna i ekologiya paukov i skorpionov. Arakhnologicheskiy sbornik. Moscow, Nauka Publ. P.58-62 [in Russian].
- Zonstein S.L. 1984. [To the fauna and ecology of spiders (Aranei) of lower strata of walnut-fruit forests of southern Kirghizia]// Entomol issled. v Kirghizii. Frunze, Ilim Publ. No.17. P.144-151 [in Russian].
- Zubko T.L., Roshchinenco V.I. 1981. [To the spider fauna of some regions of Udmurtian ASSR]// Fauna i ekologiya zhivotnykh Udmurtskoy ASSR i prilezhashchikh rayonov. Izhevsk. P.48-57 [in Russian].