

## Redescription of *Oecobius tadzhikus* Andreeva et Tyshchenko, 1969 (Aranei: Oecobiidae)

## Переописание *Oecobius tadzhikus* Andreeva et Tyshchenko, 1969 (Aranei: Oecobiidae)

Yuri M. Marusik<sup>1</sup>, Mikhail M. Omelko<sup>2,3</sup>, Seppo Koponen<sup>4</sup>  
Ю.М. Марусик<sup>1</sup>, М.М. Омелько<sup>2,3</sup>, С. Копонен<sup>4</sup>

<sup>1</sup> Institute for Biological Problems of the North of the Russian Academy of Sciences, Portovaya Str. 18, 685000 Magadan, Russia. E-mail: yurmar@mail.ru

<sup>2</sup> Far Eastern Federal University, Sukhanova Str., 8, Vladivostok 690950 Russia. Email: omelkom@gmail.com

<sup>3</sup> Gornotaezhnaya Station FEB RAS, Gornotaezhnoe Vil., Ussuriyski Dist., Primorski Krai 692533 Russia.

<sup>4</sup> Zoological Museum, University of Turku, FI-20014, Turku, Finland. E-mail: sepko@utu.fi

<sup>1</sup> Институт биологических проблем Севера ДВО РАН, Портовая 18, Магадан 685000, Россия.

<sup>2</sup> Дальневосточный Федеральный университет, ул. Суханова 8, Владивосток, 690950, Россия.

<sup>3</sup> Горнотаёжная станция им. В.Л. Комарова ДВО РАН, с. Горнотаёжное, Уссурийский район, Приморский край 692533, Россия.

KEY WORDS: Araneae, spider, Oecobiinae, Central Asia, Tajikistan, endogyne.

КЛЮЧЕВЫЕ СЛОВА: Araneae, паук, Oecobiinae, Средняя Азия, Таджикистан, эндогина.

**ABSTRACT.** In this paper we provide detailed re-description of *Oecobius tadzhikus* Andreeva et Tyshchenko, 1969 based on the types. Endogyne of this species is described for the first time.

**РЕЗЮМЕ.** Приводится детальное переописание *Oecobius tadzhikus* Andreeva et Tyshchenko, 1969 на основе типовой серии. Эндогина этого вида описана впервые.

### Introduction

*Oecobius* Lukas, 1846, with 84 species [Platnick, 2014; World Spider Catalog, 2015], is the largest genus of the family (110 species). The genus is restricted to the Holarctic Region. Only four species are known outside this region (from India, Burkina Faso, Chad and Ghana), and all of them most likely do not belong to *Oecobius*. *Oecobius* is a rather poorly studied genus. There are only three revisional publications on this group, by Shear [1970], Shear & Benoit [1974] and Wunderlich [1995] covering the Nearctic, Africa, and the Mediterranean region, respectively.

Twenty-eight species of *Oecobius* are known from a single sex: 16 from females and 12 from males [Platnick, 2014]. Half of the species are known from a single taxonomic record, and 20 species were addressed in two taxonomic publications connected to each other (same specimens were studied, or same figures were used) [Platnick, 2014].


More than half of *Oecobius* species occur in the Mediterranean, but the highest species diversity (34 species) was observed in the Canary Islands. Only three

species of *Oecobius* are known to occur in Central Asia: *O. nadiæ* (Spassky, 1936), *O. przewalskyi* Hu et Li, 1987, and *O. tadzhikus* Andreeva et Tyshchenko, 1969. Only the former species is well known; it is widespread in the region and was mentioned in several publications. *Oecobius przewalskyi* is known only from a type series of 5 females from the eastern part of the Xizang Province of China [Hu, 2001]. *Oecobius tadzhikus* is also known from its type series only, which was collected in the western Tajikistan. Although the latter species was addressed in two taxonomical publications, both were based on the same specimens and featured same figures. These published figures were very schematic and illustrated female habitus, epigyne and prolateral view of the male palp. Recently, we got an opportunity to examine the types of *Oecobius tadzhikus* and decided to redescribe it, since it is a poorly known species: the existing illustrations are schematic, and the description was published only in Russian.

### Methods

The photographs were taken using an Olympus SZX16 stereomicroscope with an Olympus E-520 camera and prepared using CombineZP software at the Zoological Museum of the University of Turku. The species description format follows Shear [1970].

Abbreviations: ALE — anterior lateral eyes, AME — anterior median eyes, PLE — posterior lateral eyes, PME — posterior median eyes. Leg articles: F — femur, Mt — metatarsus, Pt — patella, T — tarsus, Ti — tibia. All measurements are given in millimeters.



Figs 1–6. Habitus and male palp of *Oecobius tadjhikus*: 1 — habitus of male; 2 — habitus of female holotype; 3 — male palp, anteroventral; 4 — male palp, retrolateral; 5 — male palp, ventral; 6 — male palp, prolateral. Abbreviations: *Al* — loop of the spermophor on anterior part of the tegulum; *At* — anterior part of the tegulum; *Ce* — claw like extension of *Ta*; *Pl* — loop of the spermophor on prolateral side of the tegulum; *Ta* — terminal apophysis.

Рис. 1–6. Габитус и пальпа *Oecobius tadjhikus*: 1 — габитус самца; 2 — габитус самки (голотип); 3 — пальпа самца, спереди-снизу; 4 — пальпа самца, ретролатерально; 5 — пальпа самца, снизу; 6 — пальпа самца, пролатерально. Сокращения: *Al* — петля семенного канала в передней части тегулюма; *At* — передняя часть тегулюма; *Ce* — когтевидный вырост *Ta*; *Pl* — петля семенного канала на пролатеральной части тегулюма; *Ta* — терминальный отросток.

## Taxonomy


*Oecobius tadjhikus* Andreeva et Tyshchenko, 1969  
Figs 1–13.

*O. t.* Andreeva, Tyshchenko, 1969: 376, f. 3 (♂♀).

*O. t.*: Andreeva, 1976: 22, f. 19–22 (♂♀, copy of figures from Andreeva & Tyshchenko [1969]).

TYPES: Holotype ♀ and paratypes 1♂ 1♀ 3 juv., Tajikistan, Chiluchor-chashma, under stones on slope, 8.05.1965 (E. Martynova) from Zoological Institute, St. Petersburg (Russia), examined.

NOTE. Although holotype is mentioned in Andreeva & Tyshchenko [1969], both females were found in one vial without a specification which of them is holotype.



Figs 7–12. Epigyne of paratype female of *Oecobius tadjikus*: 7–8 — ventral; 9–10 — posterior, 11 — dorsal; 12 — posterior-dorsal. Scale = 0.2 mm if not otherwise indicated. Abbreviations: *Fc* — fertilization chamber; *Fd* — fertilization duct; *Id* — insemination duct; *Ok* — oval chamber of the receptacle; *Sc* — scape; *Sr* — sack-like chamber of the receptacle; *Tr* — tube-like part of the receptacle.

Рис. 7–12. Эпигина паратипа *Oecobius tadjikus*: 7–8 — снизу; 9–10 — сзади, 11 — сверху; 12 — сзади-сверху. Масштаб 0,2 мм, если не указано иначе. Сокращения: *Fc* — оплодотворительная камера; *Fd* — оплодотворительный канал; *Id* — копулятивный канал; *Ok* — овальная камера рецептакулы; *Sc* — скапус; *Sr* — мешковидная камера рецептакулы; *Tr* — трубковидная часть рецептакулы.

**DIAGNOSIS.** Male of *Oecobius tadjikus* differs well from other Palearctic congeners by a strongly developed anterior part of tegulum (*At*) being as large as the “embolic division” (much smaller in other species) and having a loop of spermophor on prolateral side, lacking in other congeners. Females differ from all other species occurring in Central Asia and Palearctic by very large receptacles (Figs 8, 11–12) and transverse fertilization chambers (*Fc*, Figs 10–12).

**DESCRIPTION.** *Male.* Total length 2.73. Carapace wider than long: 0.98 long, 1.13 wide; dirty brown without distinct pattern due to poor preservation (Fig. 1). According to original description, carapace had a brown spot. Sternum, maxillae, labium and clypeus yellow. Sternum outlined by a thin gray line. Clypeal projection evenly rounded. Carapace highest behind eye area. AME round, the largest in size, separated by a distance less than their single diameter. ALE small, adjoined with AME. PLE elongate, located close to AME. PME irregular, opalescent, separated a distance equal to their doubled diameter. Abdomen yellow with a longitudinal stripe crossed by two transverse stripes. Legs yellow without spots or rings.

Leg measurements in male/female: FI 0.93/1.13, FII 0.93/1.10, FIII 1.05/1.13, FIV 1.15/1.38, PtI 0.30/0.38, PtII 0.38/0.40, PtIII 0.28/0.43, PtIV 0.30/0.43, TiI 0.83/0.78, TiII 0.85/0.88, TiIII 0.90/0.88, TiIV 1.00/1.00, MtI 0.75/0.78, MtII 0.88/0.90, MtIII 0.95/1.00, MtIV 1.13/1.13, TI 0.65/0.63, TII 0.73/0.70, TIII 0.65/0.65, TIV 0.63/0.63; total length: leg I 3.46/3.70, leg II 3.77/3.98, leg III 3.83/4.09, leg IV 4.21/4.57.

Palp as in Figs 3–6, bulb as large as cymbium; tegulum with a very large anterior part (*At*); spermophor very long, encircling whole tegulum, and forming two circular loops (coils) on the anterior part of tegulum (*Al*) and an elongated loop (*Pl*) in the prolateral-basal part of tegulum. Terminal apophysis (*Ta*) large, with square-like base and claw-like extension (*Ce*).

*Female* (holotype). Total length 3.08. Carapace 1.0 long, 1.13 wide. Carapace dirty-brown without distinct pattern. Sternum, maxillae and labium yellow. Clypeal projection evenly rounded. Abdomen yellow with longitudinal stripe crossed by pair transverse stripes. ALE biggest, located close to AME. AME second in size, separated by a distance slightly less than their single diameter. PME much bigger than in males, irregular, opalescent, separated from each other a distance equal to their diameter. PLE small, located close both to PLE and AME. Carapace with a large round spot at the anterior part. Clypeus with a pair of black dots. Sternum, maxillae and labium yellow. Sternum outlined by a thin gray line like in males. Dorsal side of abdomen light brown, covered by white spots, with a dark longitudinal stripe in anterior part. Ventral side without white spots. Legs yellow. Tibia with poorly visible spots at ventral sides.

Epigyne as in Figs 7–9, with a kind of scape (*Sc*). Ventral and posterior part of epigynal plate with transverse wrinkles. Insemination ducts (*Id*) long, connected to a complex receptacle composed of three parts: oval chambers (*Or*), wide tube-like part (*Tr*) and weakly sclerotized large sack part (*Sr*). Sack part connected






Fig. 13. Distribution records of *Oecobius tadjikus*: Open circles refer to doubtful records from Turkmenistan.  
 Рис. 13. Распространение *Oecobius tadjikus*: Незалитые кружки — сомнительные указания в Туркмении.

with fertilization chamber (*Fc*) by a long duct (*Fd*). Fertilization chamber transverse.

**COMMENTS ON POSITION OF THE SPECIES.** Since *Oecobius tadjikus* has a rather unusual epigyne (large receptacles, transverse fertilization chambers) and male palp (a very long spermophor forming several loops unknown in other congeners), it can very likely belong to a different genus.

**DISTRIBUTION.** Besides Tajikistan, *Oecobius tadjikus* was reported also from the western Turkmenistan [Mikhailov, Fet, 1994], but these records could refer to another species (V. Fet, pers. comm.).

**ACKNOWLEDGEMENTS.** The English of an earlier draft was kindly checked by Victor Fet (Huntington, USA).

## References

- Andreeva E.M., Tyshchenko V.P. 1969. [On the fauna of spiders (Araneae) from Tadjikistan. Haplogynae, Cribellatae, Ecribellatae Trionychae (Pholcidae, Palpimanidae, Hersiliidae, Oxyopidae)] // Entomologicheskoe Obozrenie. Vol.48. No.2. P.373–384 [in Russian].
- Andreeva E.M. 1976. [Spiders of Tajikistan]. Dushanbe. 196 p. [in Russian]
- Hu J.L. 2001. [Spiders in Qinghai-Tibet Plateau of China]. Henan Science and Technology Publishing House. 658 pp. [in Chinese].
- Mikhailov K.G. 2013. The spiders (Arachnida: Aranei) of Russia and adjacent countries: a non-annotated checklist // Arthropoda Selecta. Supplement 3. 262 pp.
- Mikhailov K.G., Fet V. 1994. Fauna and zoogeography of spiders (Aranei) of Turkmenistan // Fet V., Atamuradov K.I. (eds.). Biogeography and Ecology of Turkmenistan. Dordrecht: Kluwer Academic Publisher. P.499–524.
- Platnick N.I. 2014. The World Spider Catalog. Version 15. American Museum of Natural History, New York. Available from: [http://research.amnh.org/iz/spiders/catalog\\_15.0/index.html](http://research.amnh.org/iz/spiders/catalog_15.0/index.html) (accessed 28 March 2015).
- Shear W.A. 1970. The spider family Oecobiidae in North America, Mexico, and the West Indies // Bulletin of the Museum of Comparative Zoology. No.140. P.129–164.
- Shear W.A., Benoit P.L.G. 1974. New species and new records in the genus *Oecobius* Lucas from Africa and nearby islands (Araneae: Oecobiidae: Oecobiinae) // Revue Zoologique Africaine. T.88. P.706–720.
- World Spider Catalog. 2015. Available from: <http://wsc.nmbe.ch/> (accessed 28 March 2015).
- Wunderlich J. 1995. Zu Taxonomie und Biogeographie der Arten der Gattung *Oecobius* Lucas 1846, mit Neubeschreibungen aus der Mediterraneis und von der Arabischen Halbinsel (Arachnida: Araneae: Oecobiidae) // Beiträge zur Araneologie. Bd.4. S.585–608.

Responsible editor K.G. Mikhailov