

Deep-sea fauna of European seas: An annotated species check-list of benthic invertebrates living deeper than 2000 m in the seas bordering Europe. Aplacophora

D.L. Ivanov¹, A.H. Scheltema²

¹ *Zoological Museum, Moscow State University, Bol'shaja Nikitskaja str. 6, Moscow 125009, Russia. E-mail: ivanovdl@zmmu.msu.ru*

² *Woods Hole Oceanographic Institution, Woods Hole, MA 02543 USA.*

ABSTRACT: An annotated check-list is given of Aplacophora species occurring deeper than 2000 m in the seas bordering Europe. The check-list is based on published data. The check-list includes 18 species. For each species synonymy, data on localities in European seas and general species distribution are provided. Station data are presented separately in the present thematic issue.

How to cite this article: Ivanov D.L., Scheltema A.H. 2014. Deep-sea fauna of European seas: An annotated species check-list of benthic invertebrates living deeper than 2000 m in the seas bordering Europe. *Aplacophora // Invertebrate Zoology*. Vol.11. No.1. P.8–12.

KEY WORDS: deep-sea fauna, European seas, Aplacophora.

Глубоководная фауна европейских морей: аннотированный список видов донных беспозвоночных, обитающих глубже 2000 м в морях, окружающих Европу. Арпласорфора

Д.Л. Иванов¹, А.Г. Шелтема²

¹ *Зоологический музей МГУ им. М.В. Ломоносова, ул. Большая Никитская, 6, Москва 125009, Россия. E-mail: ivanovdl@zmmu.msu.ru*

² *Woods Hole Oceanographic Institution, Woods Hole, MA 02543 USA.*

РЕЗЮМЕ: Приводится аннотированный список видов Арпласорфора, обитающих глубже 2000 м в морях, окружающих Европу. Список основан на опубликованных данных. Список насчитывает 18 видов. Для каждого вида приведены синонимия, данные о нахождении в европейских морях и сведения о распространении. Данные о станциях приводятся в отдельном разделе настоящего тематического выпуска.

Как цитировать эту статью: Ivanov D.L., Scheltema A.H. 2014. Deep-sea fauna of European seas: An annotated species check-list of benthic invertebrates living deeper than 2000 m in the seas bordering Europe. *Aplacophora // Invertebrate Zoology*. Vol.11. No.1. P.8–12.

КЛЮЧЕВЫЕ СЛОВА: глубоководная фауна, европейские моря, Арпласорфора.

Phylum Mollusca
Class Aplacophora
Subclass Solenogastres
Family Acanthomeniidae

Genus *Acanthomenia* Thiele, 1913

TYPE SPECIES: *Acanthomenia gaussiana* Thiele, 1913 (by monotypy).
 COMPOSITION: 2 species.
 DISTRIBUTION: 2000–4300 m, Atlantic — 1, Antarctic (Davis Strait) — 1.
 CENTRE OF DIVERSITY: unknown.

***Acanthomenia arcuata* Scheltema, 1999**

Acanthomenia arcuata Scheltema, 1999: 7–10, fig. 1A, B, 2, 3A, B, 4.
 LOCALITIES: INCAL (CENTOB), St. DS 01, DS 02, DS 05, DS 06, DS 07, DS 08, DS 09, DS 15, DS 16, CP 01, CP 02, CP 03, CP 04, CP 05, CP 06, CP 07, OS 01, OS 06, OS 07, OS 08, WS 01, WS 07, WS 09 (Scheltema, 1999).
 DISTRIBUTION: West European Basin.
 DEPTH RANGE: 2068–4327 m.

Family Dondersiidae

Genus *Lyratoherpia* Salvini-Plawen, 1978

TYPE SPECIES: *Lyratoherpia carinata* Salvini-Plawen, 1978.
 COMPOSITION: 2 species.
 DISTRIBUTION: 385–2091 m, Atlantic — 1, Antarctic (Ross Sea) — 1.
 CENTRE OF DIVERSITY: unknown.

***Lyratoherpia incali* Scheltema, 1999**

Lyratoherpia incali Scheltema, 1999: 11–15, fig. 1C, E, 3C, D, 5, 6.
 LOCALITIES: INCAL (CENTOB), Sta. DS-01 (Scheltema, 1999).
 DISTRIBUTION: Type locality.
 DEPTH RANGE: 2091 m.

Family Macellomeniidae

Genus *Macellomenia* Simroth, 1893

TYPE SPECIES: *Paramenia palifera* Pruvot, 1893 (by monotypy).
 COMPOSITION: 2 species.
 DISTRIBUTION: 80–2897 m, Atlantic — 2, Mediterranean Sea — 1.
 CENTRE OF DIVERSITY: unknown.

***Macellomenia aciculata* Scheltema, 1999**

Macellomenia aciculata Scheltema, 1999: 25–27, fig. 11, 12.
 LOCALITIES: INCAL (CENTOB), St. DS-09, WS-02 (Scheltema, 1999).
 DISTRIBUTION: West European Basin.
 DEPTH RANGE: 2498–2897 m.

Family Simrothiellidae

Genus *Kruppomonia* Nierstrasz, 1905

TYPE SPECIES: *Kruppomonia minima* Nierstrasz, 1905 (by monotypy).
 COMPOSITION: 4 species.
 DISTRIBUTION: 250–4327 m, Atlantic — 4, Mediterranean Sea — 1.
 CENTRE OF DIVERSITY: unknown.

Kruppomonia levis

Scheltema et Schander, 2000

Kruppomonia levis Scheltema, Schander, 2000: 134–137, fig. 6F, H, 12A–C, F, G, 13A, 14A–D.
 LOCALITIES: INCAL (CENTOB), Sta. DS 14, DS 16, OS 07, OS 08; BIOGAS VI, St. DS 76, DS 77 (Scheltema, Schander, 2000).
 DISTRIBUTION: West European Basin.
 DEPTH RANGE: 4228–4327 m.

Kruppomonia delta

Scheltema et Schander, 2000

Kruppomonia delta Scheltema, Schander, 2000: 137–138, fig. 6G, 12D, E, H–K, 13B, 14E–J.
 LOCALITIES: INCAL (CENTOB), Sta. DS 15, DS 16, OS 06, OS 08 (Scheltema, Schander, 2000).
 DISTRIBUTION: West European Basin.
 DEPTH RANGE: 4211–4327 m.

Genus *Plawenia*

Scheltema et Schander, 2000

TYPE SPECIES: *Simrothiella schizoradulata* Salvini-Plawen, 1978 (OD).
 COMPOSITION: 3 species.
 DISTRIBUTION: 2000–6000 m, Atlantic — 2, Antarctic — 1.
 CENTRE OF DIVERSITY: unknown.

Plawenia sphaera

Scheltema et Schander, 2000

Plawenia sphaera Scheltema, Schander, 2000: 140–141, fig. 15D, E, 16 sclerites 2, 8–11, 17C, D, J, 18B, G, H, K, 19D–G.
 LOCALITIES: INCAL (CENTOB), Sta. DS 01, DS 02, DS 05, CP 01, CP 02; Sarsia St. 50 (Scheltema, Schander, 2000).
 DISTRIBUTION: West European Basin.
 DEPTH RANGE: 1739–2503 m.

Genus *Simrothiella* Pilsbry, 1898

TYPE SPECIES: *Solenopus margaritaceus* Koren et Danielsen, 1978 (SD).
 COMPOSITION: 1 species.
 DISTRIBUTION: 70–2173 m, NE Atlantic — 1.
 CENTRE OF DIVERSITY: unknown.

Simrothiella margaritacea**(Koren et Danielssen, 1877)***Simrothiella margaritacea* Scheltema, Schander, 2000: 143–144, fig. 11A–C, 20–22.

LOCALITIES: “Chain-106”, St. 316 (Scheltema, Schander, 2000).

DISTRIBUTION: West European Basin.

DEPTH RANGE: 70–2173 m.

Genus *Spioenia* Arnofsky, 2000TYPE SPECIES: *Spioenia spiculata* Arnofsky, 2000 (by monotypy).

COMPOSITION: 1 species.

DISTRIBUTION: 2070–4800 m, Atlantic — 1.

CENTRE OF DIVERSITY: unknown.

Spioenia spiculata* Arnofsky, 2000Spioenia spiculata* Arnofsky, 2000: 113–117, fig. 1–3.

LOCALITIES: INCAL (CENTOB), Sta. DS 02, DS 05, DS 06, DS 07, DS 08, DS 09, CP 01, CP 02, CP 05, CP 06, CP 07, CP 08, CP 09, OS 02, OS 06, WS 01, WS 02 (Arnofsky, 2000).

DISTRIBUTION: West European Basin.

DEPTH RANGE: 2068–4829 m.

Subclass Caudofoveata**Family Scutopodidae****Genus *Scutopus* Salvini-Plawen, 1968**TYPE SPECIES: *Scutopus ventrolineatus* Salvini-Plawen, 1968.

COMPOSITION: 4 species.

DISTRIBUTION: Cosmopolitan (?), 40–3542 m, Atlantic — 3, Pacific — 1.

CENTRE OF DIVERSITY: unknown.

Scutopus robustus* Salvini-Plawen, 1970Scutopus robustus* Salvini-Plawen, 1970: 3–5, 2, 4; Ivanov, Scheltema, 2001: 2, 2A.

LOCALITIES: POLYMEDE I, St. KR 04/13, KR 10/31; POLYMEDE II, St. DS 21/45 (Salvini-Plawen, 1977).

DISTRIBUTION: A temperate, Iceland-East Atlantic, near-continental species, living on shelf and upper slope.

DEPTH RANGE: 50–3542 m [very common in the Atlantic and the western Mediterranean at 50 to 685 m. Three records in the Mediterranean at depth of 2415, 2917 and 3542 m (Salvini-Plawen, 1977) seem doubtful (Ivanov, Scheltema, 2001)].

Genus *Psilodens* Savini-Plawen, 1977TYPE SPECIES: *Limifossor elongatus* Salvini-Plawen, 1972.

COMPOSITION: 2 species.

DISTRIBUTION: 1609–3760 m, Pacific — 1, Atlantic — 2.

CENTRE OF DIVERSITY: unknown.

Psilodens tenuis* Salvini-Plawen, 1977Psilodens tenuis* Salvini-Plawen, 1977: 415–417, 1–3.

LOCALITIES: POLYMEDE I, St. DS 14 (Salvini-Plawen, 1977).

DISTRIBUTION: Known from type locality only — Atlantic, off the Strait of Gibraltar.

DEPTH RANGE: 2500 m.

Family Prochaetodermatidae**Genus *Prochaetoderma* Thiele, 1902**TYPE SPECIES: *Chaetoderma radulifera* Kowalevsky, 1901 (by monotypy).

COMPOSITION: 6 species.

DISTRIBUTION: 50–2290 m. Atlantic — 4, Mediterranean Sea — 1, Arabian Sea — 1, (Scheltema, 1985; Scheltema, Ivanov, 2000; Ivanov, Scheltema, 2002; Ivanov, Scheltema, 2008).

CENTER OF DIVERSITY: unknown.

Prochaetoderma yongei* Scheltema, 1985Prochaetoderma yongei* Scheltema, 1985: 501–509, fig. 3a–c, 4a–c, 6–8, 14y.

LOCALITIES: INCAL, St. DS 01, DS 02; CP 07; BIOGAS IV, St. DS 52 (Scheltema, Ivanov, 2000); BIOICE, St. 2856, 2859 (Ivanov, Scheltema, 2001).

DISTRIBUTION: North American Basin, West European Basin, Cape Verde Basin, and Cape Basin (Scheltema, 1985; Scheltema, Ivanov, 2000; Scheltema, Ivanov, 2009).

DEPTH RANGE: 450–2270 m.

Genus *Chevroderma* Scheltema, 1985TYPE SPECIES: *Chevroderma turnerae* Scheltema, 1985 (OD).

COMPOSITION: 10 species.

DISTRIBUTION: 1180–8390 m. Atlantic — 4, Indian Ocean — 5, Pacific — 2. (Scheltema, 1985; Scheltema, Ivanov, 2000; Ivanov, Scheltema, 2002; Ivanov, Scheltema, 2008).

CENTRE OF DIVERSITY: unknown.

Chevroderma turnerae* Scheltema, 1985Chevroderma turnerae* Scheltema, 1985: 512–516, fig. 3 l–p, 5b, d, f, 10, 15t.

LOCALITIES: “Chain-106”, St. 323, 326, 328; 330; INCAL (CENTOB), St. DS 09, DS 10, DS 11, DS 13, DS 16, CP 09, CP 10, CP 11, CP 12, OS 01, OS 02, OS 03, OS 04, OS 06, OS 07, OS 08, WS 05, WS 07, WS 08, WS 09, WS 10; NORATLANTE, St. B16, 113-EO13; ABYPLAINE (CENTOB), St. DS 07, DS 09, DS 10; BIOGAS III, St. DS 45, DS 47, DS 48, DS 50; BIOGAS IV, St. DS 55, DS 56, DS 60, KR 33, KR 35; BIOGAS V, St. DS 66, DS 67; BIOGAS VI, St. DS 77, DS 78, DS 79, DS 80, DS 82,

DS 85, CP 14, CP 15, CP 16, CP 17, CP 19, CP 20, CP 22, CP 22A; BIOGAS IX, St. KG 178; BIOGAS XI, St. KG 203, KG 206, KG 207, KG 208, KG 209, KG 210, KG 211, KG 212, KG 213, KG 215, KG 216, KG 218 (Scheltema, 1985; Scheltema, Ivanov, 2000).

DISTRIBUTION: Lower continental slope to mostly abyssal depths throughout the Atlantic Ocean.

DEPTH RANGE: 2180–5320 m (Scheltema, 1985; Scheltema, Ivanov, 2000; Ivanov, Scheltema, 2008).

***Chevroderma gauson* Scheltema, 1985**

Chevroderma gauson Scheltema, 1985: 516–519, fig. 3q–t, 5a, 11, 15g.

LOCALITIES: “Chain-106”, St. 328, 330; INCAL, St. WS 03, OS 02 (Scheltema, 1985; Scheltema, Ivanov, 2000).

DISTRIBUTION: West European Basin.

DEPTH RANGE: 4426–4829 m.

Claviderma Ivanov et Scheltema, 2000

Rhabdoderma — Scheltema, 1989: 56 [non Reiss, 1888, in Pisces, a fossil coelocanth].

TYPE SPECIES: *Rhabdoderma australe* Scheltema, 1989 (OD).

COMPOSITION: 10 species.

DISTRIBUTION: 100–3800 m. Atlantic — 7, Indian Ocean — 2, southwest Pacific — 1 (Scheltema, Ivanov, 2000; Ivanov, Scheltema, 2002; Ivanov, Scheltema, 2008).

CENTRE OF DIVERSITY: unknown.

***Claviderma gladiatum* (Salvini-Plawen, 1992)**

Prochaetoderma gladiatum Salvini-Plawen, 1992: 324, fig. 10, 24.

LOCALITIES: Sarsia, St. 50.

DISTRIBUTION: Bay of Biscay and Cape Verde, Guinea, Angola, and Cape Basins (Salvini-Plawen, 1992; Scheltema, Ivanov, 2000).

DEPTH RANGE: 1267–3814 m.

Genus *Spathoderma* Scheltema, 1985

TYPE SPECIES: *Spathoderma clenchi* Scheltema, 1985 (OD)

COMPOSITION: 6 species.

DISTRIBUTION: 80–5300 m. Atlantic — 3, Indian — 2, Northeast Pacific — 1 (Scheltema, Ivanov, 2000; Ivanov, Scheltema, 2002).

CENTRE OF DIVERSITY: unknown.

***Spathoderma clenchi* Scheltema, 1985**

Spathoderma clenchi Scheltema, 1985: 509–511, figs. 3D–G, 4D, F, 9, 14c.

LOCALITIES: “Chain-106”, St. 321, 323; INCAL, St. DS 01, DS 02, DS 05, DS 06, DS 07, DS 08,

DS 09, DS 10, CP 01, CP 02, CP 03, CP 04, CP 05, CP 06, CP 07, OS 01, WS 02; BIOGAS IV, St. DS 51, DS 61, DS 62, DS 63, DS 64; BIOGAS V, St. DS 70; BIOGAS VIII, St. KG 142, KG 144, KG 145, KG 157; BIOGAS IX, St. KG 173, KG 174, KG 178, KG 179, KG 181, KG 183, KG 185, BIOGAS XI, St. KG 183, KG 185; “Mac 81”, Box no. 14 (Scheltema, 1985; Scheltema, Ivanov, 2000); BIOICE, St. 2856, 2859, 2860, 2863 (Ivanov, Scheltema, 2001).

DISTRIBUTION: North American Basin, Iceland, West European Basin to Cape Basin.

DEPTH RANGE: 1150–3350 m.

***Spathoderma grossum* Scheltema et Ivanov, 2000**

Spathoderma grossum Scheltema, Ivanov, 2000: 359, figs. 20C, 29E, F, 30, 31.

LOCALITIES: INCAL (CENTOB), St. DS 16, OS 08, WS 09; ABYPLAINE (CENTOB), St. CP 11, CP 19, DS 01, DS 02, DS 04, DS 07, DS 09, DS 11, DS 13 (Scheltema, Ivanov, 2000).

DISTRIBUTION: From West European Basin to Cape Basin (Scheltema, Ivanov, 2000); probably anti-Atlantic (unpublished data).

DEPTH RANGE: 4184–5320 m.

Family Chaetodermidae

Genus *Falcidens* Salvini-Plawen, 1968

TYPE SPECIES: *Falcidens crossotus* Salvini-Plawen, 1968 (OD).

COMPOSITION: 27 species.

DISTRIBUTION: Cosmopolitan, 3–5600 m, Atlantic — 13, Indian Ocean — 2, Pacific Ocean — 10, Arctic — 3.

CENTRE OF DIVERSITY: unknown.

***Falcidens aequabilis* Salvini-Plawen, 1972**

Falcidens aequabilis Salvini-Plawen, 1972: 32–33, 3–7.

LOCALITIES: POLYMEDE I, Sta. KR-04/13, KR-05/15, KR-10/31; POLYMEDE II, Sta. DS-01/2, DS-24/77 (Salvini-Plawen, 1972, 1977).

DISTRIBUTION: Mediterranean Sea (Salvini-Plawen, 1972, 1977, 1997).

DEPTH RANGE: 2430–3542 m.

***Falcidens thorensis* Salvini-Plawen, 1971**

Falcidens thorensis Salvini-Plawen, 1971: 30; Ivanov, Scheltema, 2001: 5, 2C.

LOCALITIES: Thor, St. 164 (Salvini-Plawen, 1971); BIOICE, St. 2855, 2856 (Ivanov, Scheltema, 2001).

DISTRIBUTION: North Atlantic, lower bathyal species.

DEPTH RANGE: 1096–2150 m.

References

- Arnofsky P. 2000. *Spiomenia spiculata*, gen. et sp. nov. (Aplacophora: Neomeniomorpha) collected from the deep waters of the West European Basin // *The Veliger*. Vol.43. No.2. P.110–117.
- Ivanov D.L., Scheltema A.H. 2001. Distribution of known caudofoveate species (Mollusca, Aplacophora) around Iceland // *Ruthenica*. Vol.11. No.1. P. 1–6.
- Ivanov D.L., Scheltema A.H. 2002. Prochaetodermatidae of the Indian Ocean collected during Soviet VITYAZ cruises 1959–1964 (Mollusca: Aplacophora) // *Molluscan Resarch*. Vol.22. P.183–202.
- Ivanov D.L., Scheltema A.H. 2008. Western Atlantic Prochaetodermatidae from 35°N south to the Argentine Basin including the Gulf of Mexico (Mollusca: Aplacophora) // *Zootaxa*. Vol.1885. P.60.
- Salvini-Plawen L., von. 1968. Über Lebendbeobachtungen an Caudofoveata (Mollusca, Aculifera) nebst Bemerkungen zum System der Klasse // *Sarsia*. Vol.31. P.105–126.
- Salvini-Plawen L., von. 1970. Die norwegischen Caudofoveata (Mollusca, Aculifera) // *Sarsia*. Vol.45. P.1–15.
- Salvini-Plawen L., von. 1971. Die nordatlantischen Caudofoveata des Zoologischen Museums zu Kopenhagen (Mollusca, Aculifera) // *Steenstrupia*. Vol.2. No.2. P.27–34.
- Salvini-Plawen L., von. 1972. Die Caudofoveata des Mittelmeeres und das Genus *Scutopus* (Mollusca, Aculifera) // B. Battaglia (ed.). *Fifth European Marine Biology Symposium*. Picin Editore, Padova. P.27–51.
- Salvini-Plawen L., von. 1977. Caudofoveata (Mollusca) des Forschungsprojektes Polymede // *Bull. Mus. Nat. Hist. Nat.* Vol.3. No.447 (Zoologie 310). P.413–421.
- Salvini-Plawen L., von. 1992. On certain Caudofoveata from the Vema-Expedition // E. Gittenberger, J. Goud (eds.). *Proc. Ninth Intern. Malacol. Congr. Edinburgh: Nat. Mus. Scotland*. P.317–333.
- Salvini-Plawen L., von. 1997. Fragmented knowlege on West-European and Iberian Caudofoveata and Sole-nogastres // *Iberus*. Vol.15. P.33–50.
- Scheltema A.H. 1985. The Aplacophoran family Prochaetodermatidae in the North American Basin, including *Chevroderma* n. g. and *Spathoderma* n. g. (Mollusca; Chaetodermomorpha) // *Biol. Bull.* Vol.169. P.484–529.
- Scheltema A.H. 1999. New Eastern Atlantic neomenioid aplacophoran molluscs (Neomeniomorpha, Aplacophora) // *Ophelia*. Vol.51. No.1. P.1–28.
- Scheltema A.H., Ivanov D.L. 2000. Prochaetodermatidae of the Eastern Atlantic Ocean and Mediterranean Sea (Mollusca, Aplacophora) // *J. Moll. Stud.* Vol.66. P.313–362.
- Scheltema A.H., Ivanov D.L. 2009. A natural history of the deep-sea aplacophoran *Prochaetoderma yongei* and its relationship to confamilials (Mollusca, Prochaetodermatidae) // *Deep Sear Research Part II: Tropical Studies in Oceanography*. Vol.55. No.19–20. P.1856–1864.
- Scheltema A.H., Schander C. 2000. Discrimination and phylogeny of Solenogaster species through the morphology of hard parts (Mollusca, Aplacophora, Neomeniomorpha) // *Biol. Bull.* Vol.198. P.121–151.

Responsible editors

A.V. Gebruk, H. Thiel, M. Thurston