

Deep-sea fauna of European seas: An annotated species check-list of benthic invertebrates living deeper than 2000 m in the seas bordering Europe. Crinoidea

Alexander N. Mironov*, Nadia Améziane, Marc P. Eléaume****

* *P.P. Shirshov Institute of Oceanology, Russian Academy of Sciences, Nakhimovsky Pr., 36, 117997, Russia. E-mail: miron@ocean.ru*

** *Muséum national d'Histoire naturelle, Département Milieux et Peuplements Aquatiques, UMR 7208-BOREA MNHN-CNRS-UPMC-IRD, 43 rue Cuvier, 75005 Paris, France. E-mails: ameziane@mnhn.fr, eleaume@mnhn.fr*

ABSTRACT: An annotated check-list is given of Crinoidea species occurring deeper than 2000 m in the seas bordering Europe. The check-list is based on published data. The check-list includes 21 species. For each species synonymy, data on localities in European seas and general species distribution are provided. Station data are presented separately in the present thematic issue.

How to cite this article: Mironov A.N., Améziane N., Eléaume M.P. 2014. Deep-sea fauna of European seas: An annotated species check-list of benthic invertebrates living deeper than 2000 m in the seas bordering Europe. Crinoidea // *Invert. Zool.* Vol.11. No.1. P.112–119.

KEY WORDS: deep-sea fauna, European seas, Crinoidea.

Глубоководная фауна европейских морей: аннотированный список видов донных беспозвоночных, обитающих глубже 2000 м в морях, окружающих Европу. Crinoidea

А.Н. Миронов*, Н. Амезиане, М.П. Элеом****

* *Институт океанологии им. П.П. Ширшова РАН, Нахимовский просп. 36, Москва, 117997, Россия. E-mail: miron@ocean.ru*

** *Muséum national d'Histoire naturelle, Département Milieux et Peuplements Aquatiques, UMR 7208-BOREA MNHN-CNRS-UPMC-IRD, 43 rue Cuvier, 75005 Paris, France. E-mails: ameziane@mnhn.fr, eleaume@mnhn.fr*

РЕЗЮМЕ: Приводится аннотированный список видов Crinoidea, обитающих глубже 2000 м в морях, окружающих Европу. Список основан на опубликованных данных. Список насчитывает 21 вид. Для каждого вида приведены синонимия, данные о нахождении в европейских морях и сведения о распространении. Данные о станциях приводятся в отдельном разделе настоящего тематического выпуска.

Как цитировать эту статью: Mironov A.N., Améziane N., Eléaume M.P. 2014. Deep-sea fauna of European seas: An annotated species check-list of benthic invertebrates living deeper than 2000 m in the seas bordering Europe. Crinoidea // *Invert. Zool.* Vol.11. No.1. P.112–119.

КЛЮЧЕВЫЕ СЛОВА: глубоководная фауна, европейские моря, Crinoidea.

Phylum Echinodermata**Class Crinoidea****Order Comatulida**

REMARKS: Classification of crinoids used in the present work is in accordance with that in Hess, Messing (2011).

Family Antedonidae**Genus *Leptometra* A.H. Clark, 1908**

COMPOSITION: 2 species (Clark A.H., 1923a). Type species: *L. phalangium* (J. Müller, 1841). Unpublished molecular data suggest that the two *Leptometra* species are synonyms.

DISTRIBUTION: Atlantic from the northern coast of Scotland and Hebrides to Cap Blanc, the Mediterranean; perhaps also off Brazil.

DEPTH RANGE: from 42 to 1292 m (?2789–2834 m).

***Leptometra phalangium* (J. Müller, 1841)**

Leptometra phalangium — Clark A.H., Clark A.M., 1967: 553–564, fig. 2:a, b.

REMARKS: The average depth of occurrence is 300 m (Clark A.H., Clark A.M., 1967), the only occurrence below 2000 m (Clark A.M., 1977) is erroneous. The specimen dredged on the “Atlantis II” cruise 59, St. 211 (33°57'N, 15°08.2'E) was deposited at the Smithsonian Institution under reference number E16294. Both the label and the list of the “Atlantis II” stations were checked by Cynthia Ahearn. It was concluded that the specimen cited by A.M. Clark (1977) was dredged from 500–509 m depth, not 2834–2789 m.

LOCALITIES: “Atlantis II”, cruise 59, St. 211 (Clark A.M., 1977).

DISTRIBUTION: Mediterranean, 70–1292 m.

DEPTH RANGE: 500–509 m (?2789–2834 m).

Genus *Thaumatometra* A.H. Clark, 1908

COMPOSITION: 9 species (Clark A.H., Clark A.M., 1967). Type species: *T. tenuis* (Clark A.H., 1907).

DISTRIBUTION: North Pacific (1 species), northwest Pacific (4 species), Indonesia (1 species), southwest Pacific (1 species); North Indian Ocean (1 species), southwest Indian Ocean (1 species), Atlantic from Iceland and Baffin Bay to Brazil (2 species).

CENTRE OF DIVERSITY: northwest Pacific (4 species).

DEPTH RANGE: 146–3229 m.

Thaumatometra septentrionalis**A.H. Clark, 1918**

Thaumatometra septentrionalis — Clark A.H., Clark A.M., 1967: 752–753.

LOCALITIES: INGOLF, St. 11, 18, 36 (Clark A.H., 1923a; Clark A.H., Clark A.M., 1967).

DISTRIBUTION: Baffin Bay to Iceland.

DEPTH RANGE: 2075–2623 m.

Genus *Trichometra* A.H. Clark, 1908

COMPOSITION: 2 species. Type species: *T. cubensis* (Pourtalès, 1869)

DISTRIBUTION: northwest Pacific (1 species), off Hawaiian Islands (1 species); Atlantic (2 species) from Iceland and Davis Straits to the Bay of Biscay and Brazil.

DEPTH RANGE: 223–2834 m.

***Trichometra cubensis* (Pourtalès, 1869)**

Trichometra cubensis — Clark A.M., 1970: 46–48, fig. 17; Clark A.M., 1980: 195–197; Messing, Dearborn, 1990: 25–26, fig. 17, 18; Hemery et al., 2013: 513..

REMARKS: A.H. Clark (1923a) recorded *Trichometra delicata* at the “Ingolf” station 18.61°44'N, 30°29'W, 2075 m. A.M. Clark (1970) considered this small specimen conspecific with *T. cubensis*. It is possible that smaller specimens from “Helga” stations in the Bay of Biscay and from west of Ireland reported as *T. delicata* will also prove to be conspecific with *T. cubensis* (Harvey et al., 1988).

LOCALITIES: INGOLF, St. 24, and perhaps St. 18, 36 (Clark A.H., 1923a); “Discovery”, St. 7711#66 (Clark A.M., 1980); BIOGAS III, St. 1 CV 24; BIOGAS IX, St. 1 CP 33; BIOGAS XI CP 37 (Roux, pers. com.).

DISTRIBUTION: Iceland, the Faeroe Islands southwards to Morocco, and Greenland, Davis Strait to the West Indies, northwest Pacific, off Hawaiian Islands.

DEPTH RANGE: 210–2834 m (Meyer et al., 1978; Clark A.M., 1980).

***Trichometra delicata* A.H. Clark, 1911**

Trichometra delicata — Clark A.M., 1970: 48–49, fig. 18.

LOCALITIES: “Travailleur”, St 13, (Folin, 1887; Clark A.H., 1911; Marshall, Rowe, 1981); BI- AÇORES, St. 252 (unpublished).

DISTRIBUTION: Ireland to Portugal.

DEPTH RANGE: 700–2700 m.

Family Pentametrocrinidae**Genus *Pentametrocrinus* A.H. Clark, 1908**

COMPOSITION: 6 species (Messing, Hansson, 2012). However Messing (2008) listed 12 species, including 3 undescribed ones.

DISTRIBUTION: northwest Pacific (1 species), northeast Pacific (1 species, Messing, 2008), west Pacific (1 species), southwest Pacific (2 species, McKnight, 1977), Indian Ocean (2 species), Atlantic (1 species, Messing, 1978).

CENTRE OF DIVERSITY: Not recognizable.

DEPTH RANGE: 374–3320 m (or 188–4851 m in accordance with Messing, 2008).

***Pentametrocrinus atlanticus* (Perrier, 1883)**

Pentametrocrinus atlanticus — Messing, 1978: 700–708, figs. 1–5, 8–11, 13–16; Clark A.M., 1980: 203–204; Messing, Dearborn, 1990: 26.

LOCALITIES: “Talisman”, St. “45” (30°01′N, 09°25′W, depth 2215 m) (Folin, 1887; Clark A.H., 1911).

REMARKS: *Pentametrocrinus atlanticus* is one of four amphi-Atlantic comatulids that occur in the tropical western Atlantic. The other three are *Trichometra cubensis*, *Zenometra columnaris* and *Atelecrinus balanoides* (Messing, 1978; Messing, White, 2001). A.H. Clark obviously took data of “Talisman” stations from Smith (1889). However, the official list of Commandant Parfait published in the *Annales Hydrographiques* (1883) differs considerably from that of Smith. The reasons for this discrepancy are given in Bauchot et al. (1971).

DISTRIBUTION: northeast Atlantic from southwest of Ireland to north-west Africa, western Atlantic from the Blake Plateau to Martinique (Lesser Antilles) (Clark A.M., 1980).

DEPTH RANGE: 374–2115 m.

Genus *Thaumatoctrinus***P.H. Carpenter, 1883**

COMPOSITION: 6 species (Clark A.H., Clark A.M., 1967). Type species: *T. renovatus* Carpenter, 1883.

DISTRIBUTION: northwest Pacific (1 species), west Pacific (2 species), South Australia (1 species), Antarctic (1 species), Indian Ocean (1 species), Atlantic (1 species).

CENTRE OF DIVERSITY: West Pacific (2 species).

DEPTH RANGE: 649–3290 (Clark A.H., 1923a).

Thaumatoctrinus jungerseni**A.H. Clark, 1923**

Thaumatoctrinus jungerseni — Clark A.H., 1923a: 13–17, fig. 2–4; Clark A.M., 1980: 204–205.

REMARKS: Distinction between *T. jungerseni* and *T. renovatus* remains unclear (Clark A.H., Clark A.M., 1967; Clark A.M., 1980), only their geographic distribution (northern and southern hemisphere respectively) separates the two species.

LOCALITIES: “INGOLF”, St. 18 (Clark A.H., 1923a); “Discovery”, St. 7711#57, 7711#62, 7711#85 (Clark A.M., 1980); “Discovery”, St. 9640#1 (Gage et al., 1983); “Challenger”, St. SWT 12, SWT 13 (Gage et al., 1983).

DISTRIBUTION: Denmark Strait and the Rockall Trough to southern Ireland.

DEPTH RANGE: 823–3757 m.

Family Thalassometridae**Genus *Thalassometra* A.H. Clark, 1907**

COMPOSITION: 19 species (Clark A.H., 1950; McKnight, 1977; Clark A.M., 1980; Eléaume et al.,

2004). Type species: *Thalassometra villosa* A.H. Clark, 1907.

REMARKS: A.M. Clark (1980) suggested that a single damaged specimen identified as *T. omissa* and referred by Koehler (1909) to the *basicurva* group (i.e. *Thalassometra* of A.H. Clark, 1907), could refer to the Antedonidae. One of us (MPE) examined the specimen and confirmed that it belongs to the family Thalassometridae.

DISTRIBUTION: North Pacific (1 species), northwest Pacific (2 species), west Pacific (4 species), southwest Pacific (2 species), east Pacific (3 species), Indian Ocean (2 species), Antarctic and Subantarctic (2 species), Atlantic (3 species).

CENTRE OF DIVERSITY: west Pacific (4 species).

DEPTH RANGE: 148–4265 m.

Thalassometra lusitanica**(P.H. Carpenter, 1884)**

Stiremetra lusitanica — Clark A.H., 1950: 121–126; *Thalassometra lusitanica* — Clark A.M., 1980: 191–193.

LOCALITIES: “Princesse Alice”, St. 1116 (Koehler, 1909), ABYPLAINE, St. CP 68 (unpublished); CYMOR II, DR 24 (Roux, pers. com.).

DISTRIBUTION: Bay of Biscay (47–49°N, 09–11°W; unpublished data) to Morocco.

DEPTH RANGE: 914–2165 m.

Family Zenometridae**Genus *Zenometra* A.H. Clark, 1907**

COMPOSITION: 1 species (Messing, White, 2001). Type species: *Z. columnaris* (Carpenter, 1881).

DISTRIBUTION: Blake Plateau, northern Straits of Florida and Caribbean Sea (Messing, Dearborn, 1990), west of Gibraltar (Messing, White, 2001), Cape Basin (unpublished).

DEPTH RANGE: 504–4015 m.

Zenometra columnaris**(P.H. Carpenter, 1881)**

Zenometra columnaris — Clark A.H., Clark A.M., 1967: 496–499; Messing, Dearborn, 1990: 26, fig. 12; Messing, White, 2001: 163–164, figs. 1c, 2a, 3.

REMARKS: This amphi-Atlantic comatulid occurs much deeper in the eastern Atlantic (3465–4014 m) than in the western Atlantic (504–1034 m).

LOCALITIES: “Shackleton”, cruise 3, St. 169 (Messing, White, 2001).

DISTRIBUTION: as for the genus.

DEPTH RANGE: 504–4015 m.

Family Bathyrinidae**Genus *Bathyrinus* Thomson, 1872**

COMPOSITION: 9 species (Mironov, 2008). Type species: *B. gracilis* Thomson, 1872.

DISTRIBUTION: northwest Pacific (3 species), southwest Pacific (2 species), east Pacific (2 species), Indian Ocean (1 species), Antarctic (1 species), Atlantic (3 species), Arctic (1 species). *B. volubilis* is the deepest crinoid species found in the northwest Pacific at depths 8175–9375 m (Mironov, 2000).

CENTRE OF DIVERSITY: Not recognizable.

DEPTH RANGE: 520–9345 m.

Bathycrinus aldrichianus

Thomson, 1876

Bathycrinus campbellianus — Carpenter, 1884: 238–240, fig.15; pl. 7a, 22, 23; pl.8; *Bathycrinus serratus* — Clark A.H., 1908: 205–207, fig.1; *Bathycrinus aldrichianus* — Macurda, Meyer, 1976: 647–667, figs. 1–5; pl. 1–5; Eléaume et al., 2012: 3–4, fig. 1.

REMARKS: The Atlantic species *B. aldrichianus* and *B. gracilis* are very closely related and may perhaps ultimately prove to be identical. In the northeast Atlantic, *B. aldrichianus* is known only as fragments of two stems (Clark A.H., 1949; Macurda, Meyer, 1976), therefore identification is questionable.

LOCALITIES: “Atlantis”, cruise 152, St. 30 (Clark A.H., 1949).

DISTRIBUTION: west Atlantic between 38° 20'N and 37°13'S, central Atlantic, and Gulf of Guinea in the East Atlantic.

DEPTH RANGE: 3317–5860 m.

Bathycrinus carpenteri (Danielssen et Koren, 1877)

Ilycrinus carpenteri — Danielssen, 1893: 1–23, pls. 1–5; Clark A.M., 1970: 14–18, fig.3.

LOCALITIES: Norwegian North-Atlantic Expedition 1876–1878, St. 40, 51, 53, 205, 295, 303 (Danielssen, 1893); INGOLF, St. 113 (Clark, 1923a); Swedish Arctic Expedition, Spitzbergen, 1898, St. 26, 27 (Gislén, 1938); “Sadko”, St. 1935–59, 1938–97, 1938–101 (Djakonov, 1946); “Ob”, St. 6, 22, 45 (Koltun, 1964); “F.Litke”, St. 32 (Baranova, 1964); “Polarstern” ARK VIII/2 (Arctic EPOS), St. 078, 108 (Piepenburg et al., 1996); “Polarstern” ARK XV/1, St. 1/7.99, Dive 4 VICTOR 6000, St. 2/7.99, Dive 4 VICTOR 6000 (unpublished); “Jean Charcot” (NORBI Swedish-French Expedition.), St. 1 (DS 02, CP 01, CP 02), St. 2 (CP 04), St. 5 (CP 09, CP 10, DS 11), St. 10 (CP 16), St. 11 (CP 17) (Dahl et al., 1976; Roux, pers. com.).

DISTRIBUTION: Arctic, from the Norwegian and Greenland Seas to the Laptev Sea.

DEPTH RANGE: 520–3800 m.

Bathycrinus gracilis

Thomson, 1872

Bathycrinus gracilis — Carpenter, 1884: 243–245, fig. 16, pl. 8a, figs 1–3; Clark A.M., 1980: 206–207, fig. 5; Roux et al., 2002: 822, Fig. 7c–d.

LOCALITIES: “Porcupine” 1869, St. 37 (Wyville Thomson, 1875; Carpenter, 1884); “Princesse Alice” 1896, St. 749 (Koehler, 1909); “Chain”, cruise 106, St. 328 (Clark A.M., 1977); “Meteor”, cruise 3, St. 24, 30 AT, 38 AT (Clark A.M., 1980); “Challenger”, St. ES 27 (Harvey et al., 1988).

DISTRIBUTION: West European and Iberian Basins, Porcupine Abyssal Plain, Porcupine Seabight and Rockall Trough.

DEPTH RANGE: 2880–5275 m.

Monachocrinus A.H. Clark, 1917

COMPOSITION: 6 species. According to A.M. Clark (1982), the genus includes 7 recent species. The species *M. perrieri* (Koehler, Vaney, 1910) is regarded herein as a junior synonym of *M. recuperatus*. Type species: *M. sexradiatus* A.H. Clark, 1917. *Ilycrinus recuperatus* Perrier, 1885 was claimed to be the type species by Roux et al. (2002).

DISTRIBUTION: Atlantic (3 species), southwest Pacific (2 species), Indian Ocean (1 species). North of the Azores to South Africa, also the West Indies; in the Pacific the New Zealand and New Caledonia regions; in the Indian Ocean the Bay of Bengal.

CENTRES OF DIVERSITY: Not recognizable.

DEPTH RANGE: 1000–4850 m.

Monachocrinus recuperatus (Perrier, 1885)

Ilycrinus recuperatus — Perrier, 1886: 273, figs. 193, 242, 2; *Bathycrinus perrieri* — Koehler, Vaney, 1910: 27–28, figs. 1–2; *Monachocrinus coelus* — Clark A.H., 1923b: 229, pl. 8:1; *Monachocrinus recuperatus* — Gislén, 1951: 52–53, fig.1; Roux et al., 2002: 823, Fig. 7e.

LOCALITIES: “Talisman” 1883, St. 136, St. 40 (Folin, 1887; Koehler, Vaney, 1910); “Michael Sars”, 1910, St. 25 A (Murray, Hjort, 1912; Grieg, 1932); Swedish Deep-Sea Expedition, St. 387 (Gislén, 1951); BIOGAS 3, station without number (47° 32.3'N, 09°36'W, depth 4256 m); BIOGAS 5, St. 4 (CP 05); St. 6 (CP 07); BIOGAS 6, St. 1 (CP 09); St. 3 (CP 13); St. 3 (CP 14); St. 3 (DS 76) POLYGAS A, St. 3 (DS 20); INCAL, St. 4 (Roux, 1985; Roux, pers. comm.).

DISTRIBUTION: north of the Azores to South Africa.

DEPTH RANGE: 1620–4850 m.

REMARK: For stations of “Talisman” see remark in *Pentametrocrinus atlanticus*. Longitudes in Koehler, Vaney (1910) are incorrect since they were calculated relative to the Paris meridian which is 02°20' east of the Greenwich meridian.

Monachocrinus sexradiatus

A.H. Clark, 1917

Monachocrinus sexradiatus — Clark A.H., 1923a: 19, 45, figs. 5–8.

LOCALITIES: INGOLF: St. 18 (Clark A.H., 1923a).

DISTRIBUTION: southwest of Iceland.

DEPTH RANGE: 2075 m.

Family Septocrinidae

Genus *Zeuctocrinus* A.M. Clark, 1973

COMPOSITION: 2 species (Mironov, 2000).

Type species: *Z. gisleni* A.M. Clark, 1973.

REMARKS: Mironov (2002) refers the genus *Zeuctocrinus* to the family Septocrinidae Mironov, 2002.

DISTRIBUTION: northeastern Atlantic and northwestern Pacific (Volcano Trench).

DEPTH RANGE: 2000–6785 m.

Zeuctocrinus gisleni A.M. Clark, 1973

Zeuctocrinus gisleni — Clark A.M., 1973: 277–281, fig. 5; pl. 2; Roux, 1977: 34, pl. I: 6–8; pl. III: 1–7; Roux et al., 2002: 823, Fig. 7a–b.

LOCALITIES: “Discovery”: St. 7711#66, 8511#2 (Clark A.M., 1973, 1980); “Thalassa” 1970, St. W 425 (Roux, 1977).

DISTRIBUTION: northeastern Atlantic from Ireland to north-west Spain (41°49′–53°11′N).

DEPTH RANGE: 2000–2574 m.

Family Bourgueticrinidae

Genus *Democrinus* Perrier, 1883

COMPOSITION: 7 species. According to Roux et al. (2002), the genus includes 9 recent species. Following Messing, Dearborn (1990), the two species, *D. rawsonii* (Pourtalès, 1874) and *D. brevis* Clark, 1909, are considered here as synonyms of *D. confifer* Clark, 1909. Type species: *D. parfaiti* Perrier, 1883.

DISTRIBUTION: northwest Pacific (1 species), west Pacific (3 species), southwest Pacific (1 species), Indian Ocean (1 species), west Atlantic (1 species), northeast Atlantic (1 species).

DEPTH RANGE: 66–4260 m.

CENTRE OF DIVERSITY: west Pacific (3 species).

Democrinus parfaiti Perrier, 1883

Rhizocrinus rawsoni — Döderlein, 1912: 12–13, fig. 4, pl. 4, 7; pl. 9, 2. Non *R. rawsoni* Pournalès, 1874; *Rhizocrinus magnus* — Gislén, 1947: 4–9, figs. 1, 2; *Democrinus parfaiti* — Roux, 1977: 39–40, figs. 4, 9, 10, 11B, 16A, pl. II: 6–8, pl. V, VI; Roux et al., 2002: 823, Fig. 8f–g.

REMARKS: It is difficult to maintain *D. parfaiti* as a valid species distinct from *D. rawsonii* (Pournalès, 1874) from the western Atlantic.

LOCALITIES: “Porcupine”, St. 43 (Thomson, 1875; Carpenter, 1884, Clark A.M., 1977); “Talisman”, St. 135 (unpublished); “Princesse Alice”, 1885, St. 515 (Koehler, 1909); SKAGERAK Exp.,

St. 28V (Gislén, 1947); “Thalassa” 1970, St. W 425, Z 453 (Roux, 1977); “Shackleton”, cruise 3, St. 152, 154, 175 (Clark A.M., 1977); “Challenger”, St. SBC 216 (Gage et al., 1983; Harvey et al., 1988); BIOGAS, St. 1 (DS 61) (Roux, 1985; pers. comm.), ABYPLAINE, St. CP 69 and CP 68 (unpublished).

DISTRIBUTION: southwest Ireland and the Rockall Trough to the Azores, Morocco and Canary Islands, including Ormonde Bank (Gorringe Ridge). Probably occurs in the western Atlantic.

DEPTH RANGE: 1430–4260 m.

Genus *Conocrinus* d’Orbigny, 1850

COMPOSITION: 7 recent species (Roux et al., 2002) and numerous fossils. Type species: *D. thorenti* (d’Archiac, 1846); Eocene.

DISTRIBUTION: west Pacific (3 species), northeast Atlantic (2 species), North Atlantic (1 species).

DEPTH RANGE: 290–3135 m.

CENTRE OF DIVERSITY: west Pacific (3 species).

Conocrinus lofotensis (Sars, 1868)

Rhizocrinus lofotensis — Clark A.H., 1923a: 23–37, figs. 12–58; Clark A.M., 1970 (pt.): 19–22, fig. 4.a–d; *Conocrinus lofotensis* — Roux, 1985: 480.

LOCALITIES: INGOLF, St. 11, 18, 37 (Clark A.H., 1923a).

DISTRIBUTION: Norwegian coast (70°N) and Iceland to Ireland, Davis Strait to Massachusetts (?Florida).

DEPTH RANGE: 140–3135 m.

Conocrinus cabiochi Roux, 1976

Conocrinus cabiochi — Roux, 1976: 758–759, fig. 2.

LOCALITIES: “Thalassa” 1973, St. Z 453 (Roux, 1976; Marshall et al., 1981); CYAGOR (submersible “Cyana”), Pl. 22, BIOGAS 4, St. 1 DS 61 (Roux, 1985, pers. com.).

DISTRIBUTION: Bay of Biscay and Iberian Basin.

DEPTH RANGE: 1975–2700 m.

Family Phrynocrinidae

Genus *Porphyrocrinus* Gislén, 1925

COMPOSITION: 4 species (Clark, 1982; Roux et al., 2002; Messing, 2007). Type species: *P. verrucosus* Gislén, 1925.

DISTRIBUTION: west Pacific (1 species), southwest Indian Ocean (1 species), northeast Atlantic (2 species), central Atlantic (1 species).

DEPTH RANGE: 218–2780 m.

CENTRE OF DIVERSITY: northeast Atlantic (2 species).

Porphyrocrinus thalassae Roux, 1977

Porphyrocrinus thalassae — Roux, 1977: 34–38, figs. 1A, 2, 3, 5, 11, pl. I; Clark A.M., 1982: 124, 125, fig. 1; Roux et al., 2002: 824, Fig. 8a–b; Hemery et al., 2013: 515.

LOCALITIES: “Thalassa” 1970, St. W 425 (Roux, 1977); “Discovery”, St. 8511#2 (Clark A.M., 1982).

DISTRIBUTION: Porcupine Bank to northwest Spain.

DEPTH RANGE: 800–2584 m.

Order Isocrinida

Family Isselicerinidae

Genus *Endoxocrinus* (*Diplocrinus*)

A.H. Clark 1923

COMPOSITION: 3 species (David et al., 2006).

DISTRIBUTION: northwest, west and southwest Pacific (1 species), west Atlantic (2 species), northeast Atlantic (1 species).

DEPTH RANGE: 187–2070 m.

Endoxocrinus (*Diplocrinus*) *wyvillethomsoni* (Jeffreys, 1870)

Diplocrinus (*Annacrinus*) *wyvillethomsoni* — Macurda, Roux, 1981: 170, 171, 173–178, 181, 182, text-figs. 2d, 5B; pls. 7–14, 16, 7–12; *Endoxocrinus* (*Annacrinus*) *wyvillethomsoni* — Roux et al., 2002: 820, Fig. 5f,g; *Endoxocrinus* (*Diplocrinus*) *wyvillethomsoni* — David et al., 2006: 39.

LOCALITIES: “Thalassa” 1973, St. Z 453 (Conan et al., 1977).

DISTRIBUTION: Bay of Biscay to Morocco, Madeira, the Canary Islands and Azores.

DEPTH RANGE: 1246–2070 m.

REMARKS: Conan et al. (1981) noted that the depth of 2133 m for a specimen in the Museum of Berlin sampled by “Porcupine” off Portugal (Clark A.H., 1912) is likely to be erroneous.

Order Hyocrinida

Family Hyocrinidae

Genus *Anachalypsicrinus* Clark, 1973

COMPOSITION: 2 species (Roux, 1990; Roux et al., 2002). Type species: *A. nefertiti* A.M. Clark, 1973.

DISTRIBUTION: Mid-Atlantic region from 53°N to 36°N, off Newfoundland and the Bay of Biscay.

DEPTH RANGE: 1850–2810 m.

Anachalypsicrinus nefertiti

A.M. Clark, 1973

Anachalypsicrinus nefertiti — Clark A.M., 1973: 269–274, figs. 1 a–e; Roux, 1980a: 35, 53, pl. III: 1–4; Mironov, Sorokina, 1998: 24–27, fig. 8, pl. 10.4; Hemery et al., 2013: 515.

LOCALITIES: “Discovery”, St. 7711#66 (Clark, 1973); “Akademik Mstislav Keldysh”, St. 3988 (unpublished). Deep sea photographs: CYMOR II: Shamrock Canyon; Austell; Chateau Roi Arthur; Pendragon Escarpment (Roux, 1985, 1987; pers. comm.); FAMOUS expedition, 36°56.5′–

36°57.5′N, 33°05′–33°06′W, 2220, 2500, 2650 and 2700 m; 36°49.6′–36°51.5′N, 33°15′–33°16.5′W, 2519, 2600, 2630 and 2810 m (Roux, 1980b).

DISTRIBUTION: Mid-Atlantic from 53°N to 36°N, Bay of Biscay.

DEPTH RANGE: 2000–2810 m.

Genus *Gephyrocrinus*

Koehler et Bather, 1902

COMPOSITION: 2 species (Roux, Lambert, 2011).

DISTRIBUTION: northeast Atlantic (1 species) and northeast Pacific (1 species). Perhaps it occurs also in the equatorial part of the Mid-Atlantic Ridge (1°47′N, 24°26′W) at the depth 3330 m.

DEPTH RANGE: 1420–2445 (?3330) m.

Gephyrocrinus grimaldii

Koehler et Bather, 1902

Gephyrocrinus grimaldii — Koehler, 1909: 256–264, pl. I: 12; pl. XXXII: 1–9; Roux et al., 2002: 821, Fig. 10c,d; Roux, Bohn, 2010: 429–434, Figs. 2–8; *Hyocrinus* (*Gephyrocrinus*) *grimaldii* — Roux, 1980a: 33–34, 39, 43, 49, pl. I; text-fig. 1.

LOCALITIES: Deep sea photographs, CYMOR II, Shamrock Canyon, Pendragon Escarpment (Roux, 1985; pers. comm.).

DISTRIBUTION: Bay of Biscay to Madeira and the Canary Islands. Perhaps it occurs also in the equatorial part of the Mid-Atlantic Ridge (1°47′N 24°26′W).

DEPTH RANGE: 1420–2445 (?3330) m.

Acknowledgments. Data on stations were kindly made available by the following colleagues: Cynthia Ahearn (Smithsonian Institution) — stations of ATLANTIS II and “Chain”; Michel Roux (Université de Reims) — BIOGAS, FAMOUS and CYMOR 2; Gunther Krause (Alfred-Wegener-Institut, Bremerhaven) — “Polarstern”; Michel Segonzac (IFREMER) — CYAGOR; David Billett (Southampton Oceanography Centre) — “Discovery”. To all these persons we extend our thanks. The present study was supported by grant of the German Volkswagen Foundation 1/73 638 and by grant of the Russian Foundation of Basic Research No. 05-05-65283.

References

- Baranova Z.I. 1964. [Echinoderms collected by the icebreaker “F. Litke” in 1955] // Trudy Arkticheskogo i Antarkticheskogo Nauchno-Issledovatel'skogo Instituta [Trans. Arct. Antarct. Sci. Res. Inst.]. Vol.259. P.55–372 [in Russian].
- Bauchot M.-L., Iwamoto T., Geisdoerfer P., Rannou M. 1971. Etude critique des résultats des expéditions scientifiques du „Travailleur” et du „Talisman”. Nouvel examen des Macrouridae (Téléostéens, Gadiformes)

- // Bull. Mus. Nat. Hist. Nat. Paris. 3e sér. Vol.14. No.14. P.653–669.
- Carpenter P.H. 1884. Report on the Crinoidea. The stalked crinoids // Rep.Sci. Res. Voyage of H.M.S. “Challenger” (Zoology). Vol.11. No.32. P.1–440.
- Clark A.H. 1907. New genera of recent free crinoids // Smiths. Miscellaneous Collect. Vol.50. No.3. P.343–364.
- Clark A.H. 1908. New stalked crinoids from the eastern coast of North America // Proc. U.S. Nat. Mus. Vol.34. No.1607. P.205–208.
- Clark A.H. 1913. The crinoids of the Museum für Naturkunde Berlin // Proc. U.S. Nat. Mus. Vol.43. No.1937. P.381–410.
- Clark A.H. 1923a. Crinoidea // Danish Ingolf Expedition. Vol.4. No.5. P.1–58.
- Clark A.H. 1923b. The Echinoderm fauna of South Africa // Ann. South Afric. Mus. Vol.13. No.7. P.221–435.
- Clark A.H. 1949. Echinoderms from the mid-Atlantic dredged by the “Atlantis” in the summer of 1948 // J. Washing. Acad. Sci. Vol.39. No.11. P.371–379.
- Clark A.H. 1950. A monograph of the existing crinoids. 1. The comatulids, Part 4c. Superfamily Tropiometrida (the families Thalassometridae and Charitometridae) // Bull. U.S. Nat. Mus. Vol.82. No.1. 4c. P.1–383.
- Clark A.H., Clark A.M. 1967. A monograph of the existing crinoids. 1. The comatulids, Part 5. Suborders Oligophreata (concluded) and Macrophreata // Bull. U.S. Nat. Mus. Vol.82. No.1. 5. P.xiv + 1–860.
- Clark A.M. 1970. Echinodermata, Crinoidea // Mar. Invertebr. Scandinavia. Vol.3. P.1–55.
- Clark A.M. 1973. Some new taxa of recent stalked Crinoidea // Bull. Brit. Mus. (Nat. Hist.). Zoology. Vol.25. No.7. P.267–288.
- Clark A.M. 1977. Notes on deep-water Atlantic Crinoidea // Bull. Brit. Mus. (Nat. Hist.). Zoology. Vol.31. No.4. P.159–186.
- Clark A.M. 1980. Crinoidea collected by the Meteor and Discovery in the NE Atlantic // Bull. Brit. Mus. (Nat. Hist.). Zoology. Vol.38. No.4. P.187–210.
- Clark A.M. 1982. Inter-relationships of the recent stalked non-isocrinid Crinoidea // Mem. Austral. Mus. Vol.16. P.121–128.
- Conan G., Roux M., Sibuet M. 1981. A photographic survey of a population of the stalked crinoid *Diplocrinus (Annaerinus) wyvillethomsoni* (Echinodermata) from the bathyal slope of the Bay of Biscay // Deep-Sea Res. Vol.28. No.A(5). P.441–453.
- Danielssen D.C. 1893. Crinoidea // Den Norske Nordhavs-Expedition 1876–1878. Zoology. Vol.21. P.1–28.
- Dahl E., Laubier L., Sibuet M., Stromberg J.-O. 1976. Some quantitative results on benthic communities of the deep Norwegian Sea // Astarte. Vol.9. P.61–79.
- Djakonov A.M. 1946. [New and interesting species of Echinodermata from the northern Arctic Ocean] // Trudy dreifuyustchei ekspeditsii Glavsevmorputi na ledokolnom parahode “G. Sedov” 1937–1940. [Transact. Drift. Exped. Glavsevmorputi on the ice-breaker “G. Sedov” 1937–1940]. Vol.3. P.324–332 [in Russian].
- Döderlein L. 1912. Die gestielten Crinoiden der deutschen Tiefsee-Expedition // Wissenschaftliche Ergebnisse der Deutschen Tiefsee-Expedition auf dem Dampfer “Valdivia” 1898–1899. Bd.17. H.1. S.1–34.
- Eléaume M., Améziane N., Park Y.-H. 2004. Re-evaluation of the systematics of two deep-sea species of Thalassometra (Echinodermata: Crinoidea) and its biogeographical implications // Journ. Nat. Hist. Vol.38. No.15. P.1949–1968.
- Eléaume M., Bohn J.-M., Roux M., Améziane N. 2012. Stalked crinoids (Echinodermata) collected by the R/V Polarstern and Meteor in the south Atlantic and in Antarctica // Zootaxa. Vol.3425. P.1–22.
- de Folin L. 1887. Sous les mers. Campagnes d’explorations du “Travailleur” et du “Talisman”. Paris: Baillière et Fils. 340 p.
- Gislén T. 1938. A revision of the recent Bathycrinidae // Acta Universitatis Lundensis. N.F. Vol.34. No.10. P.1–30.
- Gislén T. 1947. The echinoderms collected by the “Skagerak” Expedition in the eastern Atlantic. II. Crinoidea // Göteborgs Kungl. Vetenskaps och Vitterhets Samhälles Handlingar. Ser.B. Vol.5. No.10. P.1–9.
- Gislén T. 1951. Crinoidea with a survey of the bathymetric distribution on the deep-sea crinoids // Rep. Swed. Deep-Sea Exped. 2. Zoology. Vol.4. P.49–55.
- Grieg J.A. 1932. Echinodermata // Rep. Sci. Res. “Michael Sars” North Atlantic deep-sea exped. 1910. Vol.3. No.2. P.1–47.
- Harvey R., Gage J.D., Billett D.S.M., Clark A.M., Paterson G.L.J. 1988. Echinoderms of the Rockall Trough and adjacent areas. 3. Additional records // Bull. Brit. Mus. (Nat. Hist.). Zoology. Vol.54. No.4. P.153–198.
- Hemery L.G., Roux M., Améziane N., Eléaume M. 2013. High-resolution crinoid phyletic inter-relationships derived from molecular data // Cahiers de biologie marine. Vol.54. No.4. P.511–523.
- Hess H., Messing C.G. (W.I. Ausich, coord. author). 2011. Treatise on Invertebrate Paleontology. Pt.T. Echinodermata 2. Revised. Crinoidea. Vol.3. Lawrence. Kansas: The Univ. Kansas Paleontol. Inst. 261 p.
- John D. 1937. Three new crinoids from the Indian Ocean // Ann. Mag. Nat. Hist. Vol.10. No.20. P.161–173.
- Koehler R. 1909. Échinodermes provenant des campagnes du yacht “Princesse Alice” // Rés. Camp. Sci. acc. par le Prince Albert Ier de Monaco. Vol.34. P.1–317.
- Koehler R., Vaney M.C. 1910. Note préliminaire sur les Crinoïdes du Travailleur et du Talisman // Bull. Mus. Nat. Hist. Nat. Vol.16. P.26–32.
- Koltun V.M. 1964. [On investigation of the benthic fauna of the Greenland Sea and central part of the Arctic] // Trudy Arkticheskogo i Antarkticheskogo Nauchno-Issledovatel'skogo Instituta [Transact. Arct. Antarct. Sci. Res. Inst.]. Vol.259. P.13–78 [in Russian].
- Macurda D.B., Meyer D.L. 1976. The morphology and life habits of the abyssal crinoid *Bathycrinus aldrichianus* Wyville Thomson and its paleontological implications // J. Paleontology. Vol.50. P.647–667.
- Macurda D.B., Roux M. 1981. The skeletal morphology of the isocrinid crinoids *Annaerinus wyvillethomsoni* and *Diplocrinus maclearanus* // Contrib. Mus. Paleont. Univ. Michigan. Vol.25. No.9. P.169–219.
- Marshall J., Guille A., Roux M. et al. 1981. Les types de Crinoïdes (Échinodermes) du Muséum National d’Histoire Naturelle de Paris // Bull. Mus. Nat. Hist. Nat. Paris. 4-e série. Vol.3. Sect.A. No.3. P.713–729.

- Marshall J., Rowe F.W.E. 1981. The crinoids of Madagascar // Bull. Mus. Nat. Hist. Nat. Paris. 4-e série. Vol.3. Sect.A. No 2. P.379–413.
- McKnight D.G. 1977. Additions to the New Zealand crinoid fauna // New Zealand Oceanogr. Inst. Rec. Vol.3. No 11. P.93–137.
- Messing C.G. 1978. *Pentametrocrinus atlanticus* (Perrier) (Crinoidea: Echinodermata): a review // J. Nat. Hist. Vol.12. P.699–708.
- Messing C.G. 2007. The Crinoid Fauna (Echinodermata: Crinoidea) of Palau // Pacific Science. Vol.61. No.1. P.91–111.
- Messing C.G., Dearborn J.H. 1990. Marine flora and fauna of the northeastern United States. Echinodermata: Crinoidea // NOAA Techn. Rep. Nat. Mar. Fish. Service. Vol.91. P.1–33.
- Messing C., Hansson H. 2012. *Pentametrocrinus* AH Clark, 1908 // Messing C. World List of Crinoidea. Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=123362> on 2012-11-24
- Messing C.G., White C.M. 2001. A revision of the Zenometridae (new rank) (Echinodermata, Crinoidea, Comatulidina) // Zool. Scripta. Vol.30. No.3. P.159–180.
- Mironov A.N. 2000. [New taxa of the stalked crinoids of the suborder *Bourgueticrinina*] // Zool. Zhurn. Vol.79. No.6. P.712–728 [in Russian, with English summary].
- Mironov A.N. 2008. Stalked crinoids of the family Bathycrinidae (Echinodermata) from the eastern Pacific // Invertebrate Zoology. Vol.5. No.2. P.133–153.
- Mironov A.N., Sorokina O.A. 1998. [Sea lilies of the order Hyocrinida (Echinodermata, Crinoidea)] // Zoologicheskie Issledovania. No.2. Moscow: Zoological Museum MGU. 117 p. [in Russian, with English summary].
- Murray J., Hjort J. 1912. The depths of the ocean. A general account of the modern science of oceanography based largely on the scientific researches of the Norwegian steamer “Michael Sars” in the North Atlantic. London: McMillan and Co. 821 p.
- Perrier E. 1875. Révision de la collection de Stellérides du Muséum d’Histoire Naturelle de Paris. Paris: C. Reinwald et Cie. 384 p.
- Perrier E. 1886. Les explorations sous-marines. Paris: Librairie Hachette et C. 352 p.
- Perrier E. 1893. Traité de Zoologie. Vol.1. No.2. Paris: Masson. 864 p.
- Piepenburg D., Chernova N.V., von Dorrien C.F., Gutt J., Neyelov A.V., Rachor E., Saldanha L., Schmid M.K. 1996. Megabenthic communities in the waters around Svalbard // Polar Biology. Vol.16. P.431–446.
- Rasmussen H.W., Sieverts-Doreck H. 1978. Articulata Classification // Moore R.C., Teichert C. (eds.). Treatise on Invertebrate Paleontology. Part T. Echinodermata 2 (3). Boulder, Colorado and Lawrence. Kansas: Geological Society of America and University of Kansas Press. P.813–928.
- Roux M. 1976. Découverte dans le Golfe de Gascogne de deux espèces actuelles du genre Cénozoïque *Conocrinus* (Échinodermes, Crinoïdes pédonculés) // C. R. Acad. Sci. Paris. Vol.D283. P.757–760.
- Roux M. 1977. Les Bourgueticrinina (Crinoidea) recueillis par la “Thalassa” dans le golfe de Gascogne: anatomie comparée des pédoncules et systématique // Bull. Mus. Nat. Hist. Nat. Paris. Ser. 3. Vol.426. P.25–84.
- Roux M. 1980a. Les articulations du pédoncule des Hyocrinidae (Echinodermes, Crinoïdes pédonculés): intérêt systématique et conséquences // Bull. Mus. Nat. Hist. Nat. Paris. 4e série. Vol.2. Sect.A. No.1. P.31–57.
- Roux M. 1980b. Les crinoïdes pédonculés (Échinodermes) photographiés sur les dorsales océaniques de l’Atlantique et du Pacifique. Implications biogéographiques // C. R. Acad. Sci. Paris. Vol.D291. No.11. P.901–904.
- Roux M. 1985. Les crinoïdes pédonculés (Échinodermes) de l’Atlantique N.E.: Inventaire, écologie et biogéographie // L. Laubier, C. Monniot (eds.). Peuplements profonds du Golfe de Gascogne. Brest: IFREMER. P.479–489.
- Roux M. 1990. Découverte d’une nouvelle espèce du genre *Ptilocrinus* (Echinodermes, Crinoïdes pédonculés) au large de Terre-Neuve // Canad. J. Zool. Vol.68. No.6. P.1132–1136.
- Roux M., Messing Ch.G., Améziiane N. 2002. Artificial keys to the genera of living stalked crinoids (Echinodermata) // Bull. Mar. Sci. Vol.70. No.3. P.799–830.
- Roux M., Bohn J.M. 2010. Revision of the genus *Gephyrocrinus* Koehler & Bather, 1902 (Echinodermata, Crinoidea, Hyocrinidae) // Zoosystema. Vol.32. No.3. P.425–437.
- Roux M., Lambert P. 2011. Two new species of stalked crinoids from the northeastern Pacific in the genera *Gephyrocrinus* and *Ptilocrinus* (Echinodermata, Crinoidea, Hyocrinidae). Effects of ontogeny and variability on hyocrinid taxonomy // Zootaxa. Vol.2825. P.1–54.
- Smith S. 1889. Lists of dredging stations in North American waters from 1867 to 1887 // Rep. U.S. Comm. Fish. 1886. P.869–1017.
- Speel J.A., Dearborn J.H. 1983. Comatulid crinoids from R/V “Eltanin” Cruises in the Southern Ocean // Antarctic Res. Ser. Biol. Antarct. Seas XIII. Vol.38. No.1. P.1–60.
- Thomson C.W. 1875. Les abîmes de la mer. Récit des expéditions de dragage des vaisseaux de S.M. Le Porcupine et Le Lightning pendant les étés de 1868, 1869 et 1870. Paris: Librairie Hachette et C. 454 p.

Responsible editors

A.V. Gebruk, H. Thiel, M. Thurston