

Revised list of Old World types and names of
Corticariinae (Coleoptera: Latridiidae) described by
Mannerheim and Motschulsky, 1843–1867

Исправленный список типов и названия
Corticariinae (Coleoptera: Latridiidae), описанных
Маннергеймом и Мочульским в 1843–1867
из Старого Света

Colin Johnson
КОЛИН ДЖОНСОН

17 Peaknaze Close, Glossop, Derbys., SK13 6UN, England. E-mail: Colin.Johnson17@virgin.net

KEY WORDS. Revised list, Latridiidae, Corticariinae, old world types and names, Mannerheim, Motschulsky, 1843–1867, lectotype designations, new status, new synonyms.

КЛЮЧЕВЫЕ СЛОВА. Список, типы и названия, Latridiidae, Corticariinae, Старый Свет, Маннергейм, Мочульский, 1843–1867, обозначения лектотипов, новый статус, новый синоним.

ABSTRACT. Types and names of all Mannerheim and Motschulsky old world (Eurasia, Africa, India and Sri Lanka) species of Corticariinae (Latridiidae) from 1843 to 1867, are revised and listed, together with new Lectotype designations and changes in nomenclature. Lectotype of *Coricaria fuscotestacea* Motschulsky, 1861, *C. axillaris* Mannerheim, 1844, *C. badia* Mannerheim, 1844, *C. baicalica* Mannerheim, 1844, *C. boreophila* Motschulsky, 1867, *C. convexipennis* Motschulsky, 1861, *C. crenicollis* Mannerheim, 1844, *C. cylindrica* Mannerheim, 1844, *C. flavifrons* Motschulsky, 1867, *C. gracilis* Mannerheim, 1844, *C. laticollis* Mannerheim, 1844, *C. melanophthalma* Mannerheim, 1844, *C. parvula* Mannerheim, 1844, *C. pharaonis* Motschulsky, 1867, *C. piligera* Mannerheim, 1844, *C. psammeticha* Motschulsky, 1867, *C. punctatissima* Motschulsky, 1867, *C. rubripes* Mannerheim, 1844, *C. setosa* Motschulsky, 1867, *C. subrugosa* Motschulsky, 1867, *C. subtilis* Mannerheim, 1844, *C. transversicollis* Motschulsky, 1867, *C. tincta* Mannerheim, 1844, *C. truncatella* Mannerheim, 1844, *C. unicariculata* Motschulsky, 1867, *C. validipes* Motschulsky, 1867, *Latridius formicetorum* Mannerheim, 1843, *Melanophthalma angulicollis* Motschulsky, 1866, *M. cylindricollis* Motschulsky, 1866, *M. gibbula* Motschulsky, 1866, *M. retroculis* Motschulsky, 1866 and *Mineauxia villigera* Motschulsky, 1867 designated. *Corticicara corpulenta* (Motschulsky, 1866), **stat.n.** from Sri Lanka and *Melanophthalma (Cortilena) picina* Motschulsky, 1866, **stat.n.** from South Africa., given as valid species; *Melanophthalma capicola* Belon, 1898 is **syn.n.** of *M. (C.) picina* Motschulsky, 1866.

РЕЗЮМЕ. Приведён список типов и названий видовой группы Corticariinae (Latridiidae), описанных Маннергеймом и Мочульским в 1843–1867 гг из Старого Света (Евразия, Африка, Индия и Шри Ланка). Обозначены лектотипы *Coricaria fuscotestacea* Motschulsky, 1861, *C. axillaris* Mannerheim, 1844, *C. badia* Mannerheim, 1844, *C. baicalica* Mannerheim, 1844, *C. boreophila* Motschulsky, 1867, *C. convexipennis* Motschulsky, 1861, *C. crenicollis* Mannerheim, 1844, *C. cylindrica* Mannerheim, 1844, *C. flavifrons* Motschulsky, 1867, *C. gracilis* Mannerheim, 1844, *C. laticollis* Mannerheim, 1844, *C. melanophthalma* Mannerheim, 1844, *C. parvula* Mannerheim, 1844, *C. pharaonis* Motschulsky, 1867, *C. piligera* Mannerheim, 1844, *C. psammeticha* Motschulsky, 1867, *C. punctatissima* Motschulsky, 1867, *C. rubripes* Mannerheim, 1844, *C. setosa* Motschulsky, 1867, *C. subrugosa* Motschulsky, 1867, *C. subtilis* Mannerheim, 1844, *C. transversicollis* Motschulsky, 1867, *C. tincta* Mannerheim, 1844, *C. truncatella* Mannerheim, 1844, *C. unicariculata* Motschulsky, 1867, *C. validipes* Motschulsky, 1867, *Latridius formicetorum* Mannerheim, 1843, *Melanophthalma angulicollis* Motschulsky, 1866, *M. cylindricollis* Motschulsky, 1866, *M. gibbula* Motschulsky, 1866, *M. retroculis* Motschulsky, 1866 и *Mineauxia villigera* Motschulsky, 1867. *Corticicara corpulenta* (Motschulsky, 1866), **stat.n.** из Шри Ланки и *Melanophthalma (Cortilena) picina* Motschulsky, 1866, **stat.n.** из Южной Африки, приводятся как валидные виды; *Melanophthalma capicola* Belon, 1898, **syn.n.** — как синоним *M. (C.) picina* Motschulsky, 1866.

Introduction

Count Carl Gustaf von Mannerheim (1797–1854) was the first coleopterist to attempt a monograph of the small beetles of the genera *Corticaria* Marsham, 1802 and *Lathridius* Illiger, 1801 (now *Latridius* Herbst, 1793) which together made up the family Latridiidae [Mannerheim, 1844], an interest which seems to have been awakened by his first discovery of a beetle in this family [Mannerheim, 1843]. He published additional papers on Alaskan beetles and also collaborated with other coleopterists of his day, especially Colonel Victor Ivanovitch de Motschulsky (1810–1871) who was also developing an interest in similar groups of beetles from his own studies and travels in both the old and new world [Motschulsky, 1849, 1861, 1866, 1867]. Consequently, many species of Latridiidae were described by these two great workers, so that their collections, rich in type specimens, in the University Zoological Museums of Helsinki (Finland) and Moscow (Russia) respectively, are of great importance in enabling us to interpret the many species which they described within this difficult family. The advent of new work which pioneered use of the male genitalia of Latridiidae as an important and hitherto overlooked character [Strand, 1937], laid the basis for modern studies whereby such characters provide the all important features that enable us to identify species with much greater accuracy and confidence than our predecessors. Since most of the Mannerheim and Motschulsky names and types had never been subjected to modern treatment, I resolved, nearly forty years ago, to start a revision of all the Old World Corticariinae names published by these two great coleopterists. The Mannerheim collection in Helsinki was much easier and quicker for me to study with postal loans from the museum than Motschulsky's in Moscow, because of differences in geographical position and political governance. I attempted to study all such species in both collections, and to carefully clean (many Motschulsky specimens were originally hidden by mould), dissect, remount and unambiguously label (although some specific names are now slightly outdated) where appropriate, all type material. As a result of my studies, many erroneous interpretations of catalogued species [Hetschko, 1926; Winkler, 1926] were found together with many new synonyms, all of which are given in the new Palaearctic Catalogue [Johnson, 2007]. Recent revisional work on the difficult *Melanophthalma taurica* species-group [Rücker & Johnson, 2007] made use of some of these new interpretations. Mannerheim described 39 species from Europe, Asia Minor, Caucasus, Middle Asia and Siberia. The bulk of Motschulsky's 41 species were described from Eurasia and North Africa (27), with others from South Africa (5), India and Sri Lanka (9). He also described two genera, the well-known and ubiquitous *Melanophthalma* [Motschulsky, 1866] and *Cortilena* [Motschulsky, 1867], now considered a subgenus of it.

List of Species and Types

Species are listed in alphabetical order for ease of reference and given a chronological number [in square brackets] to demonstrate priority of names for nomenclatural purposes. Type localities are given in full in their original published form, which is in Latin for Mannerheim, French for Motschulsky, together with their modern equivalents. International borders changed after World War 2, especially those between Finland and Russia, so it is essential to note that Villnäs (= Villnas), Mannerheim's ancestral home, still lies in Finland about 25 km NW of Turku / Åbo. Vyborg (= Viborg, Viipuri) is now the name of a north-west Russian city and province, Vozrozhdenie (= Kavantholm) being Mannerheim's manor from later years and 20 km NNE from Vyborg, both in the Karelian Isthmus. I have assumed that when there is only a single specimen in a collection, it represents the holotype, in cases where the original descriptions lack variation in measurements and localities. Multiple 'type' specimens are therefore syntypes from which I have selected Lectotypes, some of which have already been validated in earlier papers [Johnson 1972, 1986], but the bulk of them (15 Mannerheim names, 17 of Motschulsky) are designated below. Label limits are indicated by diagonal lines — // . The location of specimens is indicated by the symbols (ZMH) for Helsinki and (ZMM) for Moscow.

Names lacking types have some additional data summarized under 'Other material'. Only three species are without type material: *Corticaria illaesa* Mannerheim, 1844, *C. subacuminata* Mannerheim, 1844 and *C. umbilicifera* Mannerheim 1844. In the interests of nomenclatural stability, interpretation of these names follows traditional usage or synonymy, which is based on recent re-descriptions [Rücker, 1983], or correctly identified 'Other material' in the describers' collections.

Three new changes in nomenclature are: *Corticaria corpulenta* (Motschulsky, 1866) from Sri Lanka, **stat.n.**; *Melanophthalma (Cortilena) picina* Motschulsky, 1866 from South Africa, **stat.n.**; *M. capicola* Belon, 1898 is **syn.n.** of *M. (C.) picina* Motschulsky, 1866.

abietorum Motschulsky, 1867: 71 (*Corticaria*) [71]

TYPE LOCALITY: '...St. Pétersbourg...' = St. Petersburg, Russia.

TYPE MATERIAL: Holotype ♂: /type/, /15 Aug.59 fichtengr./, /*Corticaria abietorum* Motsch. Petropol/ (ZMM).

IDENTITY: synonym of *Corticaria longicornis* (Herbst, 1784) [Johnson, 2007].

adustipennis Motschulsky: 1867: 57 (*Corticaria*) [62]

TYPE LOCALITY: 'Cap de bonne Espérance' = Cape of Good Hope, South Africa.

TYPE MATERIAL: Holotype ♀: /Type/, /*Corticaria adustipennis* Motch. Cap.b.Sp./ (ZMM).

IDENTITY: confirmed synonym of *Corticaria serrata* (Paykull, 1798) [Johnson, 2007].

algerina Motschulsky, 1866: 273
(*Melanophthalma*) [46]

TYPE LOCALITY: ‘...en Algérie, en Egypte et en Corse’ = Algeria, Egypt and the French island of Corsica.

TYPE MATERIAL: Lectotype ♀ [Johnson, 1986]: /Algeria/, /Type/, /*Melanophthalma algerina* Motch. Algerie/ (ZMM). Paralectotypes: 1, /*Melanophthalma algerina* Motch. Algerie/ (ZMM); 1, /Alexandra, Meeres ufer/ (ZMM); 1, /Corsica/ (ZMM).

IDENTITY: valid species [Johnson, 2007].

angulicollis Motschulsky, 1866: 276
(*Melanophthalma*) [47]

TYPE LOCALITY: ‘Des Indes orientales’ = eastern India.

TYPE MATERIAL: Lectotype ♂ (by present designation) and paralectotype ♂: /type/, /*Melanophthalma angulicollis* Motch., Ind.or./ (ZMM).

IDENTITY: valid species [Johnson, 2007].

angulosa Motschulsky, 1849: 90
(*Corticaria*) [40]

TYPE LOCALITY: ‘...Russie méridionale’ = southern Russia.

TYPE MATERIAL: Lectotype ♂ [Johnson, 1986] and paralectotypes: 25, /*angulosa* m. Tuhugueff/, /*Melanophthalma angulosa* Motsch. Rus. mer./ (ZMM), includes 3 misidentified specimens: 2 ♀ *Melanophthalma transversalis* (Gyllenhal) and 1 ♀ *Corticaria similata* (Gyllenhal).

IDENTITY: synonym of *Melanophthalma taurica* (Mannerheim, 1844) [Johnson, 2007].

armata Mannerheim, 1844: 36
(*Corticaria*) [18]

TYPE LOCALITY: ‘Sibiria Transbaicalensi D. Motschulsky, in caju Musaeo individuum descriptum asservatur’ = Transbaikal area of eastern Siberia, Russia.

TYPE MATERIAL: Holotype ♂: /Turkinsk, Transbaikal/, /*Corticaria armata* Mann. Transbaicalia/ (ZMM).

IDENTITY: valid species [Johnson, 2007].

attenuata Motschulsky, 1867: 67
(*Corticaria*) [67]

TYPE LOCALITY: ‘Madère’ = Madeira.

TYPE MATERIAL: Holotype ♀: /Madeira/, /type/, /*Corticaria attenuata* Mot. Madeira/ (ZMM).

IDENTITY: synonym of *Corticaria fulva* (Comolli, 1837) [Johnson, 2007].

axillaris Mannerheim, 1844: 30
(*Corticaria*) [12]

TYPE LOCALITIES: ‘Rossia meridionali et ad Caucasum D. Motschulsky’ = southern Russia and Caucasus.

TYPE MATERIAL: Lectotype ♀ (by present designation) and paralectotype ♀: /Piatigorsk/, /type/, /*Corticaria axillaris* Motsch. Caucasus/ (ZMM).

IDENTITY: confirmed synonym of *Corticaria serrata* (Paykull, 1798) [Johnson, 2007].

badia Mannerheim, 1844: 25
(*Corticaria*) [7]

TYPE LOCALITIES: ‘Austria ad Linz; etiam in Sibiria et Caucaso D. Motschulsky’ = Linz in Austria; also in Siberia and Caucasus of Russia.

TYPE MATERIAL: Lectotype (by present designation) ♂: /type/, /Linz/, /*Corticaria badia* Megle, Austria/ (ZMM).

Paralectotype ♂, pinned: /Linz/, /*Corticaria badia* Megl./ (ZMM). IDENTITY: confirmed synonym of *Corticaria impressa* (Olivier, 1790) [Johnson, 2007].

baicalica Mannerheim, 1844: 42
(*Corticaria*) [23]

TYPE LOCALITY: ‘...ad lacum Baical Sibiriae D. Motschulsky’ = Lake Baikal, Siberia.

TYPE MATERIAL: Lectotype (by present designation) ♀ and paralectotypes: 2, /type/, /Turkinsk/, /*Corticaria baicalica* Motch. Sib. or./ (ZMM) — Turkinsk was a fort on the lake. Both paralectotypes are misidentified specimens: ♂ *Corticaria armata* Mannerheim and ♀ *Corticaria ferruginea* Marsham.

IDENTITY: confirmed synonym of *Corticaria rubripes* Mannerheim, 1844 [Johnson, 2007].

boreophila Motschulsky, 1867: 65
(*Corticaria*) [65]

(Note: Motschulsky avoided creating a primary homonym of *Corticaria borealis* Wollaston, 1855 by changing his original manuscript name of *borealis* to *boreophila*).

TYPE LOCALITIES: ‘... Nord de la Sibérie orientale et le Kamtschatka’ = north eastern Siberia and Kamtschatka, Russia.

TYPE MATERIAL: Lectotype (by present designation) ♂: /Kamt./, /type/, /*Corticaria borealis* Motch. Sib.or.bor./ (ZMM). Paralectotypes: 4, /Kamtch./, /type/, /*Corticaria borealis* Motsch. Sib.or.bor./ (ZMM).

IDENTITY: synonym of *Corticaria ferruginea* Marsham, 1802 [Johnson, 2007].

brevicollis Mannerheim, 1844: 52
(*Corticaria*) [28]

TYPE LOCALITIES: ‘...Styria, Helvetia et Gallia D. Motschulsky; in Italia D. Villa’ = Austria, Switzerland, France and Italy.

TYPE MATERIAL: Lectotype [Johnson, 1972] ♂: /Gratz/, /*brevicollis* Chevr./, /coll. Mannerheim/ (ZMH). Paralectotypes: ♀ /Geneva/, /type/, /*Melanophthalma brevicollis* Chevr. Helv. Geneva/ (ZMM) = *Melanophthalma transversalis* (Gyllenhal) (as *curticollis* Mannerheim) or *M. suturalis* Mannerheim; 2 ♀ /Paris/, /*Melanophthalma brevicollis* Chevr./ (ZMM) = *Melanophthalma suturalis* (Mannerheim) and *Corticaria gibbosa* (Herbst).

IDENTITY: confirmed synonym of *Corticaria minuta* (Fabricius, 1792) [Johnson, 2007].

campicola Mannerheim, 1844: 26
(*Corticaria*) [8]

TYPE LOCALITY: ‘Rossia meridionali ad Sareptam D. Motschulsky’ = Sarepta in southern Russia.

TYPE MATERIAL: Holotype ♂: /Sarepta/, /type/, /*Corticaria campicola* Motch. Rus. M. or. Sarepta/ (ZMM).

IDENTITY: confirmed synonym of *Corticaria impressa* (Olivier, 1790) [Johnson, 2007].

capensis Motschulsky, 1867: 51
(*Corticaria*) [59]

TYPE LOCALITY: ‘Du Cap de bonne Espérance’ = Cape of Good Hope, South Africa.

TYPE MATERIAL: Holotype ♀: /type/, /*Corticaria capensis* Motch. Cap.b.Sp./ (ZMM).

IDENTITY: confirmed synonym of *Corticaria pubescens* (Gyllenhal, 1827) [Johnson, 2007].

carinulata Motschulsky, 1867: 84
(*Corticaria*) [80]

TYPE LOCALITY: 'Egypte' = Egypt.
TYPE MATERIAL: Holotype ♀: /*Corticaria carinulata* Motch. Aegypt/ (ZMM).
IDENTITY: synonym of *Corticarina curta* (Wollaston, 1854) [Johnson, 2007].

ciliata Motschulsky, 1867: 55
(*Corticaria*) [61]

TYPE LOCALITY: '... de Madère' = Madeira.
TYPE MATERIAL: Holotype ♀: /*Corticaria ciliata* Motch. Madeira/, /type/, /Madeira/ (ZMM).
IDENTITY: synonym of *Corticaria fulva* (Comolli, 1837) [Johnson, 2007].

concinnum Mannerheim, 1844: 27
(*Corticaria*) [10]

TYPE LOCALITY: 'Sibiria occidentali D. Motschulsky' = western Siberia, Russia.
TYPE MATERIAL: Holotype ♂: /Omsk/, /*Corticaria concinnum* Mann. Sib. occ./ (ZMM) – Omsk is city in western Siberia..
IDENTITY: synonym of *Corticaria pubescens* (Gyllenhal, 1827) [Johnson, 2007].

convexipennis Motschulsky, 1861: 128
(*Corticaria*) [42]

TYPE LOCALITY: 'Montagnes de Nura-Ellia' = Nuwara-Eliya, Sri Lanka.
TYPE MATERIAL: Lectotype (by present designation) ♀: /type/, /*Corticaria convexipennis* Motch., Ind.or. Ceyl. Mt. Nura/, 1 female (ZMM). Paralectotype ♀: /*Corticaria convexipennis* Motch. Ceylon/ (ZMM).
IDENTITY: valid species, *Corticarina convexipennis* (Motschulsky, 1861).

corpulenta Motschulsky, 1866: 287
(*Melanophthalma*) [51]

= *inflata* Motschulsky, 1861 [44] not Rosenhauer, 1856
TYPE LOCALITY: 'Ceylan. Colombo' = Sri Lanka.
TYPE MATERIAL: Holotype ♀: /type/, /*inflata*/, /*Melanophthalma corpulenta* Mot., I.o. Ceyl. Colomb./ (ZMM).
IDENTITY: valid species, *Corticarina corpulenta* (Motschulsky, 1866), **stat.n.**

crenicollis Mannerheim, 1844: 37
(*Corticaria*) [20]

TYPE LOCALITIES: 'Finlandia orientali ad Kavantholm rarissime; in Gallia D. Motschulsky' = north-west Russia and France.
TYPE MATERIAL: Lectotype (by present designation) ♀: /Kavanth./, /Mannerheim/, /Spec. typ./, /Mus. Zool. H: fors Spec. typ. No 459 *Corticaria crenicollis* J. Sbg [sic]/ (ZMH).
IDENTITY: valid species, status confirmed [Johnson, 2007].

crocata Mannerheim, 1844: 53
(*Corticaria*) [30]

TYPE LOCALITY: 'Saxonia D. Motschulsky' = Saxony, Germany.
TYPE MATERIAL: Holotype ♀: /type/, /Saxonia/, /*Melanophthalma crocata* Motch. Saxonia/ (ZMM).
IDENTITY: synonym of *Melanophthalma transversalis* (Gyllenhal, 1827) [Johnson, 2007].

curticollis Mannerheim, 1844: 47
(*Corticaria*) [26]

TYPE LOCALITIES: '...Tauria D. Steven; in Armenia D. Motschulsky' = Ukraine and Armenia.
TYPE MATERIAL: Lectotype ♀ [Johnson, 1972]: /Steven/, /Tauria/, /Coll. Mannerh./ (ZMH). Paralectotypes: ♀ /Steven/, /Tauria/, /84/, /Coll. Mannerh./ (ZMH); ♀ /Dukulfa/ *Melanophthalma curticollis* Mann. Armenia/ (ZMM) is misidentified *Melanophthalma suturalis* (Mannerheim) [Johnson, 1986].
IDENTITY: synonym of *Melanophthalma transversalis* (Gyllenhal, 1827) [Johnson, 2007].

cylindrica Mannerheim, 1844: 35
(*Corticaria*) [17]

TYPE LOCALITIES: 'Suecia ad Holmiam D. Boheman; ad Kavantholm Gubernii Wiburgensis...., in Sibiria orientali D. Motschulsky; Finland ad Ylaene...D. Sahlberg et in Saxonia D. Märkel; Mus. D. Motschulsky' – Sweden, north-west Russia, eastern Siberia, Finland and Germany.
TYPE MATERIAL: Lectotype (by present designation) ♂: /type/, /Saxonia/, /*Corticaria cylindrica* Kunze Saxonia/ (ZMM).
IDENTITY: confirmed synonym of *Corticaria umbilicata* (Beck, 1817) [Johnson, 2007].

cylindricollis Motschulsky, 1866: 289
(*Melanophthalma*) [54]

(NB. Invalidly listed in synonymy (Mannerheim, 1844))
TYPE LOCALITIES: 'De l'Europe méridionale, en Italie, en France, et en Illyrie' = southern Europe, Italy, France and Slovenia.
TYPE MATERIAL: Lectotype (by present designation) ♀: /type/, /Ital/, /*Methophthalma cylindricollis* Motsch. Ital. Illyr. Gallia/ (ZMM). Paralectotypes: 1, /...Fontainebleau/ (ZMM); 4, /Paris/ (ZMM); 7, /Laybach/ type/ (ZMM) including misidentified ♀ of *Corticarina similata* (Gyllenhal).
IDENTITY: confirmed synonym of *Corticarina gibbosa* (Herbst, 1793) [Johnson, 2007].

cylindripennis Motschulsky, 1867: 68
(*Corticaria*) [68]

TYPE LOCALITY: 'Des environs de Kissingen en Bavarie' = Bavaria, Germany.
TYPE MATERIAL: Holotype ♀: /Kissingen/, /type/, /*Corticaria cylindripennis* Motch. Germania/ (ZMM).
IDENTITY: confirmed synonym of *Corticaria umbilicata* (Beck, 1817) [Johnson, 2007].

diluta Mannerheim, 1844, 20
(*Corticaria*) [4]

TYPE LOCALITY: 'Sibiria occidentali D. Motschulsky' = western Siberia, Russia.
TYPE MATERIAL: Holotype ♀: /type/, /Al Legastorna Gouv. Tomsk/, /Pour cet individu seul j'ai conserve votre nom *Cortic. diluta*/ *Corticaria diluti* mihi, Gouv. Tomsk/ (ZMM).
IDENTITY: synonym of *Corticaria pubescens* (Gyllenhal, 1827) [Johnson, 2007].

flavifrons Motschulsky, 1867: 66
(*Corticaria*) [66]

TYPE LOCALITY: '... de Madère' = Madeira.
TYPE MATERIAL: Lectotype (by present designation) ♂ and paralectotype ♀: /type/, /Madeira/, /*Corticaria flavifrons* Motch. Madeira/ (ZMM).

IDENTITY: synonym of *Corticaria serrata* (Paykull, 1798) [Johnson, 2007].

formicetorum Mannerheim, 1843: 85
(*Latridius*) [1].

TYPE LOCALITIES: 'Ad Yxpä prope Wiburgum..., ad Kavantholm hoc anno semel captus' = north-west Russia.

TYPE MATERIAL: Lectotype (by present designation) ♀: /Kavanth./, /Mannerh./, /cum *Formica rufa*./, /Mus Zool. H: fors Spec typ. No 462 *Corticaria formicetorum* Mann/ (ZMH). Paralectotypes: 1, /Kavanth./, /Mannerh./, /cum *Formica rufa*./, /Mus Zool. H: fors Spec typ. No 463 *Corticaria formicetorum* Mann/ (ZMH); 1, /*Corticaria formicetorum* Mannerh. Kavantholm./, /Type/, /*Corticaria formicetorum* Mannh. Finlandia/ (ZMM).

IDENTITY: confirmed synonym of *Corticaria longicollis* (Zetterstedt, 1838) [Johnson, 2007].

fuscipennis Mannerheim, 1844: 62
(*Corticaria*) [37]

TYPE LOCALITY: '...Italia boreali D. Prof. Kunze. Mus. Motschulsky;' = Nice, France.

TYPE MATERIAL: Holotype ♀: /*Latrid. ruficollis* Kunze, Nizza./, /type/, /*Corticaria fuscipennis* Mann. Nizza/ (ZMM).

IDENTITY: valid species, *Melanophthalma (Cortilena) fuscipennis* (Mannerheim, 1844) [Johnson, 2007].

fuscotestacea Motschulsky, 1861: 128
(*Coricaria*) [41]

TYPE LOCALITY: 'Montagnes de Nura-Ellia' = Nuwara-Eliya, Sri Lanka.

TYPE MATERIAL: Lectotype (by present designation) ♂ and paralectotypes: ♀ /type/, /*Melanophthalma fuscotestacea* Motch., I.o.Ceyl., Mt. N.E./ (ZMM); 1, /*Melanophthalma fuscotestacea* Motch. Ceylon/ (ZMM).

IDENTITY: synonym of *Corticaria gibbosa* (Herbst, 1793) [Johnson, 2007].

gibbula Motschulsky, 1866: 287
(*Melanophthalma*) [52]

TYPE LOCALITY: 'Indes orientales' = eastern India, including Sri Lanka.

TYPE MATERIAL: Lectotype (by present designation) ♀ and paralectotypes: 5, /type/, /*Melanophthalma gibbula* Motch., Ceylon/ (ZMM); 2, /*Melanophthalma gibbula* Motch., Ceylon/ (ZMM); 6, without original labels (ZMM).

IDENTITY: synonym of *Corticaria gibbosa* (Herbst, 1793) [Johnson, 2007].

gracilis Mannerheim, 1844: 32
(*Corticaria*) [14]

TYPE LOCALITY: 'Sibiria orientali D. Motschulsky' = eastern Siberia of Russia.

TYPE MATERIAL: Lectotype (by present designation) ♂ and paralectotypes: 4, /Turkinsk au Navalfotz/, /*Corticaria gracilis* Motch./ (ZMM); 2, /Turkinsk au Navalfotz/, /*Corticaria gracilis* Mot. Sib. or./ (ZMM); 2, /Sib. orient/, /type/, /*Corticaria gracilis* m. Sibir. Orient Gurbra/ (ZMM), 1 is misplaced *Corticaria umbilicata* (Beck).

IDENTITY: confirmed as valid species [Johnson, 2007].

hexagona Motschulsky, 1866: 279
(*Melanophthalma*) [48]

TYPE LOCALITY: 'Des Indes orientales' = eastern India.

TYPE MATERIAL: Holotype (antennae and right elytron only!): /type/, /*Melanophthalma hexagona* Motch., Ind.or./ (ZMM).

IDENTITY: synonym of *Melanophthalma angulicollis* Motschulsky, 1866 [Johnson, 2007].

hortensis Mannerheim, 1844: 53
(*Corticaria*) [29]

TYPE LOCALITY: 'Lithuania D. Motschulsky = Lithuania.
TYPE MATERIAL: Lectotype (Johnson, 1986) ♂ and paralectotypes: 2, /type/, /Lithuania/, /*Melanophthalma hortensis* Motsch. Lithuania/ (ZMM); ♂, 3, /Lithuan./, /*Melanophthalma hortensis* mihi Lithuania/ (ZMM).

IDENTITY: synonym of *Melanophthalma transversalis* (Gyllenhal, 1827) [Johnson, 2007].

illaesa Mannerheim, 1844: 33
(*Corticaria*) [15]

TYPE LOCALITIES: 'Tauria D. Steven et in Caucaso D.Motschulsky' = Ukraine and Caucasus.

TYPE MATERIAL: Syntypes are presumed lost since they are absent from both the Mannerheim (ZMH) and Motschulsky collections (ZMM).

OTHER MATERIAL: 1, 'Nurha' (? - illegible) (ZMM).
IDENTITY: valid species [Rücker, 1983], status confirmed [Johnson, 2007].

inflata Motschulsky, 1861: 129
(*Carticaria*) [44]

not Rosenhauer, 1856 (*Corticaria*)

TYPE LOCALITY: 'Ceylan. Colombo' = Sri Lanka.

TYPE MATERIAL: Holotype ♀: /type/, /*inflata*./, /*Melanophthalma corpulenta* Mot., I.o. Ceyl. Colomb./ (ZMM).

IDENTITY: synonym of *Corticaria corpulenta* (Motschulsky, 1866) – see above.

interstitialis Mannerheim, 1844: 21
(*Corticaria*) [5]

TYPE LOCALITY: 'Laponiae paroeciam Rouvaniemi D. Fellman' = Finnish Lapland.

TYPE MATERIAL: Holotype ♂: /Lapponia/, /Fellm./, /Spec. typ./Mus. Zool. H: fors Spec typ. No 451 *Corticaria interstitialis* Mann./ (ZMH).

IDENTITY: confirmed as valid species [Johnson, 2007].

intricata Mannerheim, 1844: 20
(*Corticaria*) [3]

TYPE LOCALITY: 'Sibiria D. Motschulsky' = Siberia, Russia.

Holotype ♂: /type/, /h. Uda Kul. Bad/, /*Corticaria intricata* Mann. Sibiria/ (ZMM).

IDENTITY: synonym of *Corticaria pubescens* (Gyllenhal, 1827) [Johnson, 2007].

lacerata Mannerheim, 1844: 38
(*Corticaria*) [21]

TYPE LOCALITY: 'Finlandia...ad Kavantholm capta' = north-west Russia.

TYPE MATERIAL: Holotype ♂: /Kavanth./, /Mannerheim/, /Spec. typ./Zool. Mus. H: fors Spec. typ. No. 460 *Corticaria lacerata* Mann./ (ZMH).

IDENTITY: confirmed synonym of *Corticaria crenicollis* Mannerheim, 1844 [Johnson, 2007].

lateritia Mannerheim, 1844: 24
(*Corticaria*) [24]

TYPE LOCALITY: 'In Finlandia ad Kavantholm....' = north west Russia.

TYPE MATERIAL: Holotype ♀: /Kavanth./, /Mannerheim./, /Spec. typ./, /*Corticaria lateritia* Mannerh./, /Mus. Zool. H: fors Spec. typ. No. 458 *Corticaria lateritia* Mann./ (ZMH).

IDENTITY: valid species, status confirmed [Johnson, 2007].

laticollis Mannerheim, 1844: 29
(*Corticaria*) [11]

TYPE LOCALITIES: 'Suecia; ... D. Gyllenhal...; etiam in Polonia a D. Motschulsky capta' = Sweden and Polonia in Siberia.

TYPE MATERIAL: Lectotype (by present designation) ♂: /Gyllh./, /Sub nomine *L. elongatus* minsa/, /Mus. Zool. H: fors Spec. typ. No 453 *Corticaria laticollis* Mann./ (ZMH). Paralectotype: ♀ /Polonia/, /type/, /C'est ma *C. laticollis*./, /*Corticaria laticollis* Mann, Polonia, Sibiria/ (ZMM).

IDENTITY: confirmed synonym of *Corticaria serrata* (Paykull, 1798) [Johnson, 2007].

limbicollis Motschulsky, 1867: 64
(*Corticaria*) [64]

TYPE LOCALITY: '...Irkoutsk' = Irkutsk, east Siberia, Russia.

TYPE MATERIAL: Holotype ♀: /Irkutsk./, /type/, /*Corticaria limbicollis* m. Irkutsk/ (ZMM).

IDENTITY: synonym of *Corticaria rubripes* Mannerheim, 1844 [Johnson, 2007].

maura Motschulsky, 1866: 271
(*Melanophthalma*) [45]

TYPE LOCALITIES: 'Styria et Russie méridionale' = Styria, Austria and southern Russia.

TYPE MATERIAL: Lectotype [Johnson, 1986] ♀: /Styria/, /type/, /*Melanophthalma maura* Motch./ (ZMM). Paralectotypes: ♂, 2, /maura m. Tschugueff/, /*Melanophthalma maura* Motsch. Rus. mer./ (ZMM).

IDENTITY: valid species, status confirmed [Johnson, 2007].

melanophthalma Mannerheim, 1844: 30
(*Corticaria*) [13]

TYPE LOCALITY: 'Willnaes in Finlandia...' = Finland.

TYPE MATERIAL: Lectotype (by present designation) ♂: /Villnas/, /Mannerh./, /Mus. Zool. H: fors Spec. typ. No 454 *Corticaria melanophthalma* Mann./ (ZMH). Paralectotypes: 1, with same labels but / No 455/ (ZMH); 1, /Spec. typ./, /No 457/ is misidentified *Corticaria longicollis* Zetterstedt) (ZMH).

IDENTITY: synonym of *Corticaria serrata* (Paykull, 1798) [Johnson, 2007].

pallens Mannerheim, 1844: 58
(*Corticaria*) [34]

TYPE LOCALITY: '...campis Kirgisorum D. Motschulsky' = Kazakhstan.

TYPE MATERIAL: Lectotype [Johnson, 1972] ♀: /K. Tettion/, /pallens./, /Coll. Mannerh./ (ZMH). Paralectotypes: 2, /K. Tekim./, /type/, /*Cortilena pallens* Motsch. Des. Kirg./ (ZMM); 2, /K. Tekim/ (ZMM), ♂ misidentified *Corticarina truncatella* (Mannerheim); 2, /K. Tekim/, /*Cortilena pallens* mihi K. Tekim/ (ZMM), ♀ misidentified *Corticarina minuta* (Fabricius).

IDENTITY: confirmed as valid species, *Melanophthalma* (*Cortilena*) *pallens* (Mannerheim, 1844) [Johnson, 2007].

parvicollis Mannerheim, 1844: 62
(*Corticaria*) [38]

TYPE LOCALITY: '...Daghestan D. Motschulsky' = Caucasus region of southern Russia.

TYPE MATERIAL: Lectotype [Johnson, 1972] ♀: /Caucasus/, /parvicollis./, /Coll. Mannerh./ (ZMH). Paralectotypes: ♂, ♀, /Dortin./, /type/, /*Melanophthalma parvicollis* Manh. Daghestan/ (ZMM).

IDENTITY: valid species, *Melanophthalma parvicollis* (Mannerheim, 1844) [Johnson, 2007].

parvula Mannerheim, 1844: 54
(*Corticaria*) [31]

TYPE LOCALITIES: '...Austria D. Motschulsky; in Italia D. Villa' = Austria and Italy.

TYPE MATERIAL: Lectotype (by present designation) ♀: /Linz./, /type/, /*Corticaria parvula* Schuep. Austria/ (ZMM). Paralectotype: ♀, /Linz./, foreparts missing (ZMM).

IDENTITY: valid species, *Corticarina parvula* (Mannerheim, 1844) [Johnson, 2007].

pharaonis Motschulsky, 1867: 74
(*Corticaria*) [72]

TYPE LOCALITY: '...Egypte.' = Egypt.

TYPE MATERIAL: Lectotype (by present designation) ♂ and paralectotypes: 3, /type/, /*Corticaria pharaonis* m. Aegyptus/ (ZMM); 1, /*Corticaria pharaonis* Motch. Aegypt./ (ZMM). IDENTITY: valid species [Johnson, 2007].

picina Motschulsky 1866: 285
(*Melanophthalma*) [50]

TYPE LOCALITY: 'Du Cap de Bonne-Espérance' = Cape of Good Hope, South Africa.

TYPE MATERIAL: Holotype ♂: /type/, /*Melanophthalma picina* Mots. Cap. b. Sp.[blue label]/ (ZMM) (Specimen also labelled as a male Lectotypus, det. H.W. Ruecker, 1980).

IDENTITY: valid species, *Melanophthalma* (*Cortilena*) *picina* Motschulsky, 1866, **stat. nov.** The name takes priority over *M. capicola* Belon, 1898, from South Africa, **syn. nov.**

picipennis Mannerheim, 1844: 63
(*Corticaria*) [39]

TYPE LOCALITY: 'Daghestan D. Motschulsky' = Caucasus region of southern Russia.

TYPE MATERIAL: Holotype ♂: /Timishar, Schura./, /*Corticaria picipennis* mihi Daghestan/ (ZMM).

IDENTITY: synonym of *Melanophthalma* (*Cortilena*) *fuscipennis* (Mannerheim, 1844) [Johnson, 2007].

piligera Mannerheim, 1844: 19
(*Corticaria*) [2]

TYPE LOCALITIES: 'Lithuania, Austria, Helvetia, Sibiria et Caucaso. Mus. D. Motschulsky' = Lithuania, Austria, Switzerland, Siberia and Caucasus of Russia.

TYPE MATERIAL: Lectotype (by present designation) ♂ and paralectotypes: ♀, /...Verona - Genf./, /*Corticaria piligera* Mann. Helvetia/ (ZMM); 2 ♀, /Lithuania./, /type/, (ZMM) probably *Corticaria pubescens* (Gyllenhal).

IDENTITY: synonym of *Corticaria punctulata* Marsham, 1802 [Johnson, 2007].

pilitecta Motschulsky, 1867: 83
(*Corticaria*) [79]

TYPE LOCALITY: '...Egypte.' = Egypt.

TYPE MATERIAL: Holotype ♀: /*Corticaria pilitecta* Motch. Aegypt/ (ZMM).

IDENTITY: synonym of *Corticarina curta* (Wollaston, 1854) [Johnson, 2007].

pilosa Motschulsky, 1867: 48
(*Corticaria*) [56]

TYPE LOCALITY: 'Sur les bords de la mer à Alexandrie en Egypte' = Egypt.

TYPE MATERIAL: Holotype ♀: /*Corticaria pilosa* Motch. Aegypt/ (ZMM).

IDENTITY: confirmed synonym of *Corticaria illaesa* Mannerheim, 1844 [Johnson, 2007].

psammeticha Motschulsky, 1867: 80
(*Corticaria*) [77]

TYPE LOCALITY: 'Egypte' = Egypt.

TYPE MATERIAL: Lectotype (by present designation) ♂ and paralectotypes: 3, /type/, / m. Aegyptus/ (ZMM).

IDENTITY: valid species, *Migneauxia psammeticha* (Motschulsky, 1867) [Johnson, 2007].

punctatissima Motschulsky, 1867: 69
(*Corticaria*) [69]

TYPE LOCALITY: 'De la Tauride' = Ukraine.

Lectotype (by present designation) ♀: /Tauria/, /type/, /*Corticaria punctatissima* Motsch. Tauria/ (ZMM). Paralectotypes: 7, /Sta.. Krim/, /type/, /*Corticaria punctatissima* Motsch. Tauria/ (ZMM).

IDENTITY: synonym of *Corticaria illaesa* Mannerheim, 1844 [Johnson, 2007].

quadricollis Motschulsky, 1867: 59
(*Corticaria*) [63]

TYPE LOCALITY: 'Cap de Bonne-Espérance' = Cape of Good Hope, South Africa.

TYPE MATERIAL: Holotype ♀: /type/, /*Corticaria quadricollis* Motch. Cap.b.Sp./ (ZMM).

IDENTITY: confirmed synonym of *Corticaria serrata* (Paykull, 1798) [Johnson, 2007].

quadrimaculata Mannerheim, 1844: 33
(*Corticaria*) [16]

TYPE LOCALITY: 'Georgia ad Tiflin D. Motschulsky' = Georgia.

TYPE MATERIAL: Holotype ♀: /Tiflis/, /type/, /*Corticaria quadrimaculata* Motch. Georgia/ (ZMM). IDENTITY: valid species [Johnson, 2007].

retroculis Motschulsky, 1866: 280
(*Melanophthalma*) [49]

TYPE LOCALITY: 'Des Indes orientales' = eastern India.

TYPE MATERIAL: Lectotype (by present designation) ♂ and paralectotype: ♂, /M/, /type/, /*Melanophthalma retroculis* Mots., Ind.or./ (ZMM); ♀, /*Melanophthalma retroculis* Motch., Ind.or./ (ZMM). The male paralectotype is a different and probably undescribed species, whilst the female could belong to either.

IDENTITY: valid species.

rubripes Mannerheim, 1844: 41
(*Corticaria*) [22]

TYPE LOCALITIES: 'Petropoli et circa Wiburgum in Finlandia...; ...Lapponia D. Sahlberg' = St. Petersburg, north-west Russia and Finnish Lapland.

TYPE MATERIAL: Lectotype (by present designation) ♂: /Viborg/, /Mannerh./, /Mus. Zool. H: fors. Spec. typ. No.461 *Corticaria rubripes* Mann./ (ZMH).

IDENTITY: valid species [Johnson, 2007].

rutila Motschulsky, 1866: 288
(*Melanophthalma*) [53]

TYPE LOCALITY: 'Egypte...' = Egypt.

TYPE MATERIAL: Lectotype [Johnson, 1986] ♀ and paralectotype 1, damaged: /*Melanophthalma rutila* Mot. Aegyptus/ (ZMM).

IDENTITY: synonym of *Melanophthalma sericea* (Mannerheim, 1844) [Johnson, 2007].

saginata Mannerheim, 1844: 24
(*Corticaria*) [6]

TYPE LOCALITIES: '...Finlandia...; ...D. Sahlberg,...Kavantholm Gubernii Wiburgensis...' = Finland and north-west Russia.

TYPE MATERIAL: Holotype ♀: /Kavanth/, /Mannerh./, /*Corticaria saginata* Mannerh./, /Mus. Zool. H: fors Spec. typ. No 452 *Corticaria saginata* Mann/ (ZMH).

IDENTITY: valid species [Johnson, 2007].

sericea Mannerheim, 1844: 61
(*Corticaria*) [36]

TYPE LOCALITY: '...Daghestan D. Motschulsky' = Caucasus of Russia.

TYPE MATERIAL: Lectotype (Johnson, 1986) ♀: /Timirhan Schura/, /type/, /*Melanophthalma sericea* Motch. Daghestan (ZMM). Paralectotype: ♂, /Daghestan/, /*Melanophthalma sericea* mihi Caucasus Daghestan/ (ZMM).

IDENTITY: valid species, *Melanophthalma sericea* (Mannerheim, 1844) [Johnson, 2007].

setosa Motschulsky, 1867: 49
(*Corticaria*) [58]

TYPE LOCALITY: '...sur les bords de la mer d'Azov à Tonkaïa' = Sea of Azov, Ukraine.

TYPE MATERIAL: Lectotype (by present designation) ♀ and paralectotypes: ♀, ♂, /typeLitt. M.Azov/, /*Corticaria setosa* Mots. Taurie. Azov./ (ZMM), misidentified *Corticaria illaesa* Mannerheim; 16, including 2 ♂, 1 ♀, /Ufer des Azovischen Meeres/, /*Corticaria setosa* Motch. Tauria Asov M./ (ZMM).

IDENTITY: synonym of *Corticaria quadrimaculata* Mannerheim, 1844 [Johnson, 2007].

stigma Motschulsky, 1867: 79
(*Corticaria*) [76]

TYPE LOCALITY: 'De la Styrie' = Styria, Austria.

TYPE MATERIAL: Holotype ♀: /Styria/, /type/, /*Corticaria stigma* Motchh. Styria/ (ZMM).

IDENTITY: confirmed synonym of *Corticaria longicollis* (Zetterstedt, 1838) (Johnson, 2007).

subacuminata Mannerheim, 1844: 46 (*Corticaria*) [25]

TYPE LOCALITY: 'Petropoli rarissime capta' = St. Petersburg, Russia.

TYPE MATERIAL: absent from both the Mannerheim (ZMH) and Motschulsky (ZMM) collections.

OTHER MATERIAL: 30, Tunkinsk, L.Baical (ZMM), including 1 misidentified *Corticarina minuta* (Fabricius); 11, Perm (ZMM); 1, Daur. B. (ZMM); 7, Hamar Daban (ZMM).

IDENTITY: confirmed synonym of *Corticaria ferruginea* Marsham, 1802, on this other material [Johnson, 2007].

subpicea Motschulsky, 1867: 70
(*Corticaria*) [70]

TYPE LOCALITY: 'Tauride' = Ukraine.

TYPE MATERIAL: Holotype ♂: /Tauria/, /type/, /*Corticaria subpicea* Motch. Tauria/ (ZMM).

IDENTITY: synonym of *Corticaria obscura* C. N. F. Brisout de Barneville, 1863 [Johnson, 2007].

subrugosa Motschulsky, 1867: 82
(*Corticaria*) [78]

TYPE LOCALITY: 'Egypte' = Egypt.

TYPE MATERIAL: Lectotype (by present designation) ♂ and paralectotypes: 3, /type/, /*Corticaria subrugosa* m. Aegyptus/ (ZMM).

IDENTITY: synonym of *Migneauxia psammeticha* (Motschulsky, 1867) [Johnson, 2007].

subtilis Mannerheim, 1844: 57
(*Corticaria*) [32]

TYPE LOCALITIES: 'In Sudermannia Sueciae et ad Willnaesin Finlandia australi...' = Sweden and Finland.

TYPE MATERIAL: Lectotype (by present designation) ♂: /Viborg/, /Mannerh./, /*Subtilis* Mannerh./, /Mus. Zool. H: fors Spec. typ. No. 464 *Corticaria subtilis* Mann./ (ZMH).

IDENTITY: confirmed synonym of *Corticarina similata* (Gyllenhal, 1827) [Johnson, 2007].

suturalis Mannerheim, 1844: 58
(*Corticaria*) [33]

TYPE LOCALITIES: '...in campis Kirgisorum et in Rossia meridionali ad Gurjev D. Motschulsky' = Kazakhstan and southern Russia.

TYPE MATERIAL: Lectotype [Johnson, 1986] ♂: /type/, /K. Tinion, Gurjew/, /*Corticaria suturalis* mihi Camp. Kirg. & Gurjev/ (ZMM). Paralectotype: ♀, /type/, /Gurjev/, /*Corticaria suturalis* Motch. Rus. m. or. Gurj./ (ZMM).

IDENTITY: valid species, *Melanophthalma suturalis* (Mannerheim, 1844) [Johnson, 2007].

taurica Mannerheim, 1844: 51
(*Corticaria*) [27]

TYPE LOCALITY: '...in Tauria D. Steven' = Ukraine.

TYPE MATERIAL: Lectotype (Johnson, 1972) ♀: /Steven/, /Tauria/, /82/, /Coll. Mannerh./ (ZMH).

IDENTITY: valid species, *Melanophthalma taurica* (Mannerheim, 1844) [Johnson, 2007].

tenuicornis Motschulsky, 1867: 77
(*Corticaria*) [75]

TYPE LOCALITY: 'Du Cap de Bon. Esp.' = Cape of Good Hope, South Africa.

TYPE MATERIAL: Holotype ♂: /type/, /*Corticaria tenuicornis* Motch. Cap.b.Sp./ (ZMM).

IDENTITY: synonym of *Corticaria serrata* (Paykull, 1798) [Johnson, 2007].

tincta Mannerheim, 1844: 26
(*Corticaria*) [9]

TYPE LOCALITIES: 'Daghestan et in Sibiria Transbailcalensi D. Motschulsky' = Caucasus of Russia and Transbailcalia in eastern Siberia.

TYPE MATERIAL: Lectotype (by present designation) ♂: /type/, /Timirhan Schura/, /*Corticaria tincta* Mann./ (ZMM). Paralectotypes: ♂, /Timirhan Schura/, /*Corticaria tincta* Manh Transcauc/ (ZMM); ♂, /H. Uda Kulscka Bad/, /*Corticaria tincta* Mannerh./, /*Corticaria tincta* Mann. Sibir. orient./ (ZMM), misidentified *C. armata* Mannerheim.

IDENTITY: synonym of *Corticaria pubescens* (Gyllenhal, 1827) [Johnson, 2007].

transversicollis Motschulsky, 1867: 76
(*Corticaria*) [73]

TYPE LOCALITY: '...Alexandriè, en Egypte' = Egypt.

TYPE MATERIAL: Lectotype (by present designation) ♀ and paralectotypes: ♀, /type/, /*Corticaria transversicollis* Motch. Aegyptus/ (ZMM); ♀, /*Corticaria transversicollis* Motch. Aegypt/ (ZMM).

IDENTITY: synonym of *Corticaria pharaonis* Motschulsky, 1867 [Johnson, 2007].

truncatella Mannerheim, 1844: 59
(*Corticaria*) [35]

TYPE LOCALITIES: '...Rossia meridionali ad fl. Ural prope Gurjev, in Sibiria orientali; ut et in Helvetia Italiaque D. Motschulsky' = southern Russia at foot of Urals near Gurjev, in eastern Siberia and in Switzerland (Ticino).

TYPE MATERIAL: Lectotype (by present designation) ♂ and paralectotypes: 3, /type/, /*Latridius truncatellus* mihi Helvetia/ (ZMM); 4, /Gurjev/ (ZMM); 15, /K. Uda, Kulsika Bad/, /type/, /*Corticaria truncatella* Motch. Sib. or. Rus. m. or./ (ZMM).

IDENTITY: valid species, *Corticarina truncatella* (Mannerheim, 1844) [Johnson, 2007].

umbilicifera Mannerheim, 1844: 37
(*Corticaria*) [19]

TYPE LOCALITIES: 'Rossia boreali ad Petropolin D. Motschulsky; in Bavaria rarius DD. Beck et Westerhauser' = St. Petersburg, Russia and Bavaria, Germany.

TYPE MATERIAL: syntypes are absent from both the Mannerheim (ZMH) and Motschulsky (ZMM) collections.

OTHER MATERIAL: 1, Aachen, 'Anglia' [sic] (ZMM).

IDENTITY: confirmed synonym of *Corticaria umbilicata* (Beck, 1817) on this other material [Johnson, 2007].

umbripennis Motschulsky, 1861: 129
(*Corticaria*) [43]

TYPE LOCALITY: 'Montagnes de Nura-Ellia' = Nuwara-Eliya, Sri Lanka.

TYPE MATERIAL: Holotype ♀: /Type/, /*Melanophthalma umbripennis* Motch., I.o. Ceyl. Mt. N.E./ (ZMM).

IDENTITY: synonym of *Corticarina convexipennis* (Motschulsky, 1861), **syn. n.**

unicarinulata Motschulsky, 1867: 76
(*Corticaria*) [74]

TYPE LOCALITIES: '...Madère, ...Allegmagne, ...Géorgie rosse' = Madeira, Germany and Georgia.

TYPE MATERIAL: Lectotype (by present designation) ♀: /type/, /Madeira/, /*Corticaria uncarinulata* Mots. Madeira/ (ZMM). Paralectotypes: 1, /Aachen/ (ZMM); 2, /Tiflis/, /*Corticaria uncarinulata* Motch. Transcauc/ (ZMM).

IDENTITY: synonym of *Corticaria fulva* (Comolli, 1837) [Johnson, 2007].

validipes Motschulsky, 1867: 54
(*Corticaria*) [60]

TYPE LOCALITIES: ‘...Russie méridionale à Tschougoueff, et aussi à St. Pétersbourg’ = southern Russia and St. Petersburg, Russia.

TYPE MATERIAL: Lectotype (by present designation) ♂ and paralectotypes: 2, /type/, /*Corticaria validipes* Motch. Rus.mer/ (ZMM).

IDENTITY: confirmed synonym of *Corticaria impressa* (Olivier, 1790) [Johnson, 2007].

villigera Motschulsky, 1867: 40
(*Migneauxia*) [55]

TYPE LOCALITIES: ‘...la Crimée et le Caucase’ = Crimea, Ukraine and Russian Caucasus.

TYPE MATERIAL: Lectotype (by present designation) ♀: /type/, /*Tauria*/, /*Migneauxia villigera* Motsch. Tauria/ (ZMM).

IDENTITY: confirmed synonym of *Migneauxia crassiuscula* (Aubé, 1850) [Johnson, 2007].

villosa Motschulsky, 1867
(*Corticaria*) [57]

TYPE LOCALITY: ‘Des bords du Golfe Persique en Mésopotamie’ = Persian Gulf of Iraq.

TYPE MATERIAL: Holotype ♀: /typ/, /M/, /*Corticaria villosa* Motch. Mesopotamia/ (ZMM).

IDENTITY: confirmed synonym of *Corticaria illaesa* Mannerheim, 1844 [Johnson, 2007].

ACKNOWLEDGEMENTS. The writer is deeply indebted to: old friends Dr. Hans Silfverberg and Prof. Olof Biström, University Zoological Museum, Helsinki, Finland, who kindly loaned material (often twice!) from the Mannerheim and other collections, which provided the trigger for this work, as well as helping with information on present day names of localities along the Finno-Russian border; colleagues Dr. Nikolai B. Nikitsky and his predecessor Dr. S. Kelejnikova, University Zoological Museum, Moscow, Russia, whose considerable commitment and help with loans from the Motschulsky collection over many years made the present work possible; the late (died 2008) Dr. Nicole Berti, Muséum National d’Histoire Naturelle, Paris, France, for help or loans from the Charles and Henri Brisout de Barneville and Belon collections; the late (died 1980) Dr. Andreas Strand, Oslo, for gifts of many rare species to the Manchester Museum during my time there as a member of staff; the late (died 1980) Baron Alexis de Porochin (= Poronen), Stockport, England, an excellent Finnish and British Coleopterist who was an unending source of valuable information and specimens from Finland; Tony A. Allen, London, England, who encouraged my interest in this family many years ago.

References

- Hetschko A. 1926. Lathridiidae // W. Junk W & S. Schenkling S. (eds.). Coleopterorum Catalogus. Pars 85. Berlin: W. Junk. S.86.
- Johnson C. 1972. Studies on the genera *Corticarina* Reitter and *Melanophthalma* Motschulsky (Col., Lathridiidae) // Nouvelle Revue d’Entomologie. Ser.II. T.2. P.185–199.
- Johnson C. 1986. Notes on some Palaearctic *Melanophthalma* Motschulsky (Coleoptera: Latridiidae), with special reference to *transversalis* auctt. // Entomologist’s Gazette. Vol.37. P.117–125.
- Johnson C. 2007. Latridiidae // I. Löbl & A. Smetana (eds.). Catalogue of Palaearctic Coleoptera. Stenstrup: Apollo Books. Vol. 4. P.74–80, 635–648
- Mannerheim C.G. von. 1843. Mémoire sur la recolté d’Insectes Coléoptères, faite en 1842 // Bulletin de la Société Impériale des Naturalistes de Moscou. T.16. P.70–87.
- Mannerheim C.G. von. 1844. Versuch einer monographischen Darstellung der Käfergattungen *Corticaria* und *Lathridius* // Zeitschrift für die Entomologie. Bd.5. S.1–112.
- Motschulsky V. de. 1849. Coléoptères reçus d’un voyage de M. Handshuh dans le Midi de l’Espagne, énumérés et suivis de notes // Bulletin de la Société Impériale des Naturalistes de Moscou. T.22. P.52–163.
- Motschulsky V. de. 1861. Essai d’un catalogue des insectes de l’île Ceylan // Bulletin de la Société Impériale des Naturalistes de Moscou. T.34. P.95–155.
- Motschulsky V. de. 1866. Énumération des espèces de Coléoptères rapportés de ses voyages // Bulletin de la Société Impériale des Naturalistes de Moscou. T.39 [1865]. P.225–290.
- Motschulsky V. de. 1867. Énumération des espèces de Coléoptères rapportés de ses voyages // Bulletin de la Société Impériale des Naturalistes de Moscou. T.4. P.39–103.
- Rücker H.W. 1983. Különbözö Csápú Bogarak VI. — Diversicornia VI. Bunkócsápú Bogarak VII. — Clavicornia VII // Budapest: Akadémiai Kiadó. Magyarországi Állatvilága. Fauna Hungariae. Vol.158. P.1–68.
- Rücker H.W. & Johnson C. 2007. Revision of *Melanophthalma taurica* (Mannerheim, 1844) species-group and description of three new species (Coleoptera: Latridiidae) // Lathridiidae. Ht.5. P.11–24.
- Strand A. 1937. Bemerkungen über nordische *Corticaria*- und *Corticarina*-Arten, nebst Beschreibung von zwei neuen Arten // Norsk Entomologisk Tidsskrift. Bd.5. P.7–23.
- Winkler A. 1926. Lathridiidae // Catalogus Coleopterorum regionis palaearticae (1924–32). Wien: Albert Winkler. P.736–743.