

Additions and corrections to the new Catalogue of Palaearctic  
Cerambycidae (Coleoptera) edited by I. Löbl and A. Smetana, 2010.  
Part II

Дополнения и исправления к новому каталогу палеарктических  
Cerambycidae (Coleoptera) под редакцией I. Löbl и A. Smetana, 2010.  
Часть II

M.L. Danilevsky  
М.Л. Данилевский

A. N. Severtzov Institute of Ecology and Evolution, Russian Academy of Sciences, Leninsky prospect 33, Moscow 119071, Russia. E-mail: danilevskym1@rambler.ru, danilevsky@cerambycidae.net

Институт Проблем Экологии и Эволюции им. А.Н. Северцова РАН, Ленинский пр. 33, Москва 119071, Россия.

KEY WORDS: Cerambycidae, taxonomy, new combinations, new synonyms, new names, Palearctic.

КЛЮЧЕВЫЕ СЛОВА: Cerambycidae, таксономия, новые сочетания, новые названия, Палеарктика.

ABSTRACT. About seventy misprints, wrong combinations, wrong geographical records, wrong references, wrong status of certain names, wrong synonyms, wrong authorships and dates of certain names, wrong original combinations, wrong spelling of several names and so on are fixed. Sometimes unavailable names were published as available. Missing names, geographical data and references are added. *Molorchus sterbai* Adlbauer, 1988 from Libanon is accepted as valid name. New synonyms are established: *Diastocera* Dejean, 1835 = *Analeptes* Gistel, 1848, **syn.n.**; *Leptura rufoannulata* (Pic, 1933) = *Leptura fisheriana* Gressitt, 1938, **syn.n.**; *Molorchus sterbai* Adlbauer, 1988 = *Glaphyra azri* Sama, Rapuzzi & Kairouz, 2010, **syn.n.**

РЕЗЮМЕ. Исправлено около семидесяти разнообразных ошибок: опечаток, неправильных сочетаний, синонимов, ошибочных географических указаний, ошибочно указанных дат публикации названий и их авторов, ошибочных оригинальных сочетаний, написаний и т.д. Указаны несколько непригодных названий, опубликованных как пригодные. Добавлены пропущенные названия, публикации и географические данные. *Molorchus sterbai* Adlbauer, 1988 из Ливана принят в качестве валидного названия. Предложены новые синонимы: *Diastocera* Dejean, 1835 = *Analeptes* Gistel, 1848, **syn.n.**; *Leptura rufoannulata* (Pic, 1933) = *Leptura fisheriana* Gressitt, 1938, **syn.n.**; *Molorchus sterbai* Adlbauer, 1988 = *Glaphyra azri* Sama, Rapuzzi & Kairouz, 2010, **syn.n.**

The new Catalogue of Palaearctic Cerambycidae (Coleoptera) edited by I. Löbl and A. Smetana [2010] was prepared with the participation of 11 authors: K. Adlbauer, M.L. Danilevsky, A. Drumont, L. Hubweber, Z. Komiya,

I. Löbl, J. Morati, P. Rapuzzi, G. Sama, A. Smetana and A. Weigel. The system of the authorship was rather complicated. It was partly arranged in taxonomy way and partly in geographical. So, such groups as Prioninae or Dorcadionini had own authors, while other authors were responsible for a certain geographical areas. The taxa of Western Palaearctic, for example, were arranged by G. Sama and I. Löbl, while M.L. Danilevsky and A. Smetana were responsible for the taxa from Russia and countries of the former USSR, and Mongolia. So, the authorship of all Cerambycidae genera, which are distributed in West Europe and in Russia was in fact mutual (with the exception of Dorcadionini). Each author had his own point of view on the system of each subfamily and each genus. So, the editors had to create a generally accepted system on the base of several different positions. As a result I. Löbl and A. Smetana took part in the arrangement of nearly each genus.

The necessity to join together several rather different texts (certain portions were presented to the editors in last moment before the publication) was the primary cause of many mistakes and misprints. Several names are published contemporary in different genera, the status of others was wrongly interpreted, many names were wrongly spelled or with wrong authors and dates, several geographical records were so strange, that could be simply misprints, many references were wrong, several publications were referred with two different names and so on. The main purpose of the present publication is to fix direct mistakes.

The references to the present article include only the publications absent in the references to the Catalog. The references inside the text of the present article to the publications included in the references to the Catalog have same letters after the number of the year as in the Catalog.

## 1. PAGE 47

PRINTED (TWO TIMES):

*Poecilium alni elburzense* Holzschuh, 1977

MUST BE:

*Poecilium alni elburzense* Holzschuh, 1977

## 2. PAGES 85–252

All names proposed by Holzschuh [1995] are published in the Catalog as “Holzschuh, 1965”

PRINTED:

- p.85 — *mellina* Holzschuh, 1965: 5 A: YUN  
 p.85 — *perforans* Holzschuh, 1965: 6 A: YUN  
 p.85 — *rufobrunnea* Holzschuh, 1965: 6 A: SCH  
 p.98 — *tenebraria* Holzschuh, 1965: 10 (*Anoploderomorpha*) A: YUN  
 p.109 — *congesta* Holzschuh, 1965: 11 A: YUN  
 p.109 — *eucera* Holzschuh, 1965: 12 A: YUN  
 p.117 — **genus *Thrangalia* Holzschuh, 1965: 10** type species *Thrangalia diaboliella* Holzschuh, 1995  
*diaboliella* Holzschuh, 1965: 11 A: YUN  
 p.124 — *wewalkai* Holzschuh, 1965: 9 A: TR  
*wittmeri* Holzschuh, 1965: 9 A: TR  
 p.131 — *palligera* Holzschuh, 1965: 8 A: YUN  
 p.143 — **genus *Pufujia* Holzschuh, 1965: 16** type species *Pufujia luteosignata* Pu, 1991  
 p.153 — *eximium* Holzschuh, 1965: 35 A: YUN  
 p.156 — *fallaciosum* Holzschuh, 1965: 19 A: YUN **ORR**  
 p.162 — *consona* Holzschuh, 1965: 18 A: NP SD  
 p.163 — *sausai* Holzschuh, 1965: 31 A: YUN  
 p.163 — *atricornis* Holzschuh, 1965: 33 A: BT  
 p.163 — *giganteus* Holzschuh, 1965: 34 A: YUN  
 p.163 — *modicatus* Holzschuh, 1965: 33 A: YUN  
 p.164 — *lucens* Holzschuh, 1965: 28 A: SCH  
 p.172 — *jendeki* Holzschuh, 1965: 41 A: YUN  
 p.173 — *parilis* Holzschuh, 1965: 39 A: YUN  
 p.173 — *sausai* Holzschuh, 1965: 40 A: YUN  
 p.174 — *decolorata* Holzschuh, 1965: 38 A: YUN  
 p.178 — *decora* Holzschuh, 1965: 39 A: YUN  
 p.186 — *arenbergeri* Holzschuh, 1965: 14 E: IT (Sardegna)  
 p.190 — *unanimis* Holzschuh, 1965: 26 A: SCH  
 p.193 — *fumigatum* Holzschuh, 1965: 23 A: YUN  
 p.202 — *clarinus* Holzschuh, 1965: 23 A: YUN  
 p.204 — *aethiops* Holzschuh, 1965: 25 (*Euchlanis*) A: YUN  
 p.249 — *maceki* Holzschuh, 1965: 41 A: TR  
 p.252 — *scrobicolle morulum* Holzschuh, 1965: 42 A: TR
- MUST BE:
- p.85 — *mellina* Holzschuh, 1995: 5 A: YUN  
 p.85 — *perforans* Holzschuh, 1995: 6 A: YUN  
 p.85 — *rufobrunnea* Holzschuh, 1995: 6 A: SCH  
 p.98 — *tenebraria* Holzschuh, 1995: 10 (*Anoploderomorpha*) A: YUN  
 p.109 — *congesta* Holzschuh, 1995: 11 A: YUN  
 p.109 — *eucera* Holzschuh, 1995: 12 A: YUN  
 p.117 — **genus *Thrangalia* Holzschuh, 1995: 10** type species *Thrangalia diaboliella* Holzschuh, 1995  
*diaboliella* Holzschuh, 1995: 11 A: YUN  
 p.124 — *wewalkai* Holzschuh, 1995: 9 A: TR  
*wittmeri* Holzschuh, 1995: 9 A: TR  
 p.131 — *palligera* Holzschuh, 1995: 8 A: YUN  
 p.143 — **genus *Pufujia* Holzschuh, 1995: 16** type species *Nortia luteosignata* Pu, 1991  
 p.153 — *eximium* Holzschuh, 1995: 35 A: YUN  
 p.156 — *fallaciosum* Holzschuh, 1995: 19 A: YUN **ORR**  
 p.162 — *consona* Holzschuh, 1995: 18 A: NP SD  
 p.163 — *sausai* Holzschuh, 1995: 31 A: YUN

- p.163 — *atricornis* Holzschuh, 1995: 33 A: BT  
 p.163 — *giganteus* Holzschuh, 1995: 34 A: YUN  
 p.163 — *modicatus* Holzschuh, 1995: 33 A: YUN  
 p.164 — *lucens* Holzschuh, 1995: 28 A: SCH  
 p.172 — *jendeki* Holzschuh, 1995: 41 A: YUN  
 p.173 — *parilis* Holzschuh, 1995: 39 A: YUN  
 p.173 — *sausai* Holzschuh, 1995: 40 A: YUN  
 p.174 — *decolorata* Holzschuh, 1995: 38 A: YUN  
 p.178 — *decora* Holzschuh, 1995: 39 A: YUN  
 p.186 — *arenbergeri* Holzschuh, 1995: 14 E: IT (Sardegna)  
 p.190 — *unanimis* Holzschuh, 1995: 26 A: SCH  
 p.193 — *fumigatum* Holzschuh, 1995: 23 A: YUN  
 p.202 — *clarinus* Holzschuh, 1995: 23 A: YUN  
 p.204 — *aethiops* Holzschuh, 1995: 25 (*Euchlanis*) A: YUN  
 p.249 — *maceki* Holzschuh, 1995: 41 A: TR  
 p.252 — *scrobicolle morulum* Holzschuh, 1995: 42 A: TR

## 3. PAGE 92

PRINTED:

*besikanus* Fairmaire, 1855: 318 (*Prionus*) E: AL BU GR MC MD TR YU A: CY TR  
*batelkai* Sláma, 1996: 75 (*Prionus*)  
*tangerianus* Sláma, 1996: 76 (*Prionus*)

NOTE:

*Prionus tangerianus* Slama, 1996 was described from Morocco on the base of a single old male. But newly collected males are also known. Sama [1998] mentioned a male (from coll. of A. Drumont) labelled “Maroc, Moyen Atlas, VI.1996”.

Another male is known with the label: “NE Marocco, Atlas Mts., Houria vill., 6.7.1951, Lorenc coll.” — see: [http://www.cerambycidae.cz/beetlespages/Mesoprio\\_tangerianus.htm](http://www.cerambycidae.cz/beetlespages/Mesoprio_tangerianus.htm)

Sama [1998] declared: “I regard collecting labels of these specimens quite suspect; it is extremely unlikely that professional entomologists such as Antoine, Rungs, Kocher and many others never recorded this species, all the more that it is very easily attracted to light.” Drumont [2010 — personal message] maintained Sama’s opinion, that all labels of *Mesoprius* for Africa were false.

## 4. PAGE 96 AND 109

PRINTED:

- (p. 96)  
**genus *Acanthoptura* Fairmaire, 1894a: 224** type species *Acanthoptura spinipennis* Fairmaire, 1894  
*denticollis* Holzschuh, 1993a: 17 A: SCH  
*impressicollis* Pic, 1920f: 117 (*Leptura*) A: SCH YUN  
*pallescens* Holzschuh, 1993a: 20 (*Parastrangalis*) A: GAN SHA  
 and (p. 109)  
**genus *Parastrangalis* Ganglbauer, 1889a: 57** type species *Leptura potanini* Ganglbauer, 1889  
 ...  
*palleago* Holzschuh, 1998: 25 A: HUB  
*palpalis* Holzschuh, 1991c: 29 A: SCH  
 ...  
 MUST BE  
 (p. 96)  
**genus *Acanthoptura* Fairmaire, 1894a: 224** type species *Acanthoptura spinipennis* Fairmaire, 1894  
*denticollis* Holzschuh, 1993a: 17 A: SCH  
*impressicollis* Pic, 1920f: 117 (*Leptura*) A: SCH YUN  
 and  
 (p. 109):  
**genus *Parastrangalis* Ganglbauer, 1889a: 57** type species *Leptura potanini* Ganglbauer, 1889  
 ...

*palleago* Holzschuh, 1998: 25 A: HUB  
*pallescens* Holzschuh, 1993a: 20 A: GAN SHA  
*palpalis* Holzschuh, 1991c: 29 A: SCH

...

NOTE:

*Parastrangalis pallescens* Holzschuh, 1993a is quite a normal *Parastrangalis*. Its transfer to *Acanthoptura* was just a nonsense (and not reflected in the «New Acts»).

5. PAGE 96

PRINTED:

*tabacicolor erythropus* Gebler, 1841b: 612 (*Leptura*) A: ES  
 FE JA KZ MG NT WS

MUST BE:

*tabacicolor erythropus* Gebler, 1841b: 612 (*Leptura*) A: ES  
 FE JA KZ MG NC NE SC WS

6. PAGE 96

PRINTED:

*fusca* Matsushita, 1930: 24

NOTE:

The name must be excluded from the Catalog as unavailable. It was introduced as *Alosterna tabacicolor* var. *fusca* Matsushita, 1930 (Mt. Kurodake, Hokkaido) together with *Alosterna tabacicolor* var. *bivittis*: Matsushita, 1930 (Mt. Kurodake, Hokkaido) — two variations from one locality, so “its author expressly gave it infrasubspecific rank” according to the Article 45.6.4. of ICZN.

7. PAGE 96

PRINTED:

*tabacicolor tabacicolor* DeGeer, 1775: 139 (*Leptura*) E: AL  
 AU BE BH BU BY CR CT CZ DE EN FI FR GB GE GR  
 HU IR IT KZ LA LS LT LU MC MD NE NL NR NT PL  
 RO SK SL SP ST SV SZ TR UK YU A: JA KZ NE SC TR  
 WS

MUST BE:

*tabacicolor tabacicolor* DeGeer, 1775: 139 (*Leptura*) E: AL  
 AU BE BH BU BY CR CT CZ DE EN FI FR GB GE GR  
 HU IR IT KZ LA LS LT LU MC MD NE NL NR NT PL  
 RO SK SL SP ST SV SZ TR UK YU A: KZ TR WS

8. PAGE 98

PRINTED:

*rufipes rufipes* Schaller, 1783: 296 (*Leptura*) E: AB AR AU  
 BH BU BY CR CT CZ EN FR GE GB GG GR HU IT LA  
 LT MD NT PL RO SK SL SP ST SV SZ YU UK A: ES  
 IN KZ

MUST BE:

*rufipes rufipes* Schaller, 1783: 296 (*Leptura*) E: AB AR AU  
 BH BU BY CR CT CZ EN FR GE GB GG GR HU IT LA  
 LT MD NT PL RO SK SL SP ST SV SZ YU UK A: ES  
 IN KZ TR

9. PAGE 99 AND 112

PRINTED (p. 99):

**genus *Corennys* Bates, 1884: 224** type species *Corennys sericata* Bates, 1884  
*Pseudocorennys* Pic, 1952d: 47 type species *Pyrocalymma diversicornis* Pic, 1947 (= *Pyrocalymma conspicua* Gahan, 1906  
*brevipennis prescutellaris* Pic, 1947c: 17 (*Pyrocalymma*) A:  
 CH **ORR**

*caduca* Holzschuh, 1998: 28 A: HUB  
*cardinalis* Fairmaire, 1887a: 131 (*Ephies*) A: YUN  
*circellaris* Holzschuh, 1992: 12 A: SCH

*conspicua* Gahan, 1906a: 89 (*Pyrocalymma*) A: BT HAI  
 HEB SCH SHA XIZ YUN **ORR**

*diversicornis* Pic, 1947c: 17 (*Pyrocalymma*)

*sensitiva* Holzschuh, 1998: 29 A: YUN

*sanguinea* Kano, 1933a: 271 A: HAI TAI

*sericata* Bates, 1884: 225 A: HEB JA NE SC

*taiwana* Hayashi, 1963d: 130 A: TAI

AND (p. 112)

**genus *Pyrocorenys* N. Ohbayashi & Niisato, 2009: 160**  
 type species *Pyrocalymna latipennis* Pic, 1927  
*latipennis latipennis* Pic, 1927b: 26 (*Pyrocalymna*) A: YUN  
**ORR**

*brevipennis* Pic, 1946: 17 (*Pyrocalymna*) [no such pages in the referenses!]

*latipennis taiwanensis* Hayashi, 1969a: 61 (*Formosopyrrhona*) A: TAI

MUST BE (p.99):

**genus *Corennys* Bates, 1884: 224** type species *Corennys sericata* Bates, 1884

*Pseudocorennys* Pic, 1952d: 47 type species *Pyrocalymma diversicornis* Pic, 1947 (= *Pyrocalymma conspicua* Gahan, 1906

*caduca* Holzschuh, 1998: 28 A: HUB

*cardinalis* Fairmaire, 1887a: 131 (*Ephies*) A: YUN

*circellaris* Holzschuh, 1992: 12 A: SCH

*conspicua* Gahan, 1906a: 89 (*Pyrocalymma*) A: BT HAI  
 HEB SCH SHA XIZ YUN **ORR**

*diversicornis* Pic, 1947c: 17 (*Pyrocalymma*)

*sensitiva* Holzschuh, 1998: 29 A: YUN

*sanguinea* Kano, 1933a: 271 A: HAI TAI

*sericata* Bates, 1884: 225 A: HEB JA NE SC

*taiwana* Hayashi, 1963d: 130 A: TAI

and (p. 112)

**genus *Pyrocorenys* N. Ohbayashi & Niisato, 2009: 160**  
 type species *Pyrocalymna latipennis* Pic, 1927  
*latipennis latipennis* Pic, 1927b: 26 (*Pyrocalymna*) A: YUN  
**ORR**

*brevipennis* Pic, 1947c: 17 (*Pyrocalymna*)

*latipennis taiwanensis* Hayashi, 1969a: 61 (*Formosopyrrhona*) A: TAI

NOTE:

According to Ohbayashi & Niisato [2009: 161]: “*Pyrocorenys latipennis prescutellaris* Pic, 1946” (described as “*Pyrocalymna brevipennis* var. *prescutellaris* Pic, 1946: 17”) is known only from North Vietnam.

10. PAGES 102–103

PRINTED: (p. 102)

**genus *Judolia* Mulsant, 1863: 496** type species *Leptura sexmaculata* Linnaeus, 1758

*Judolia* Pic, 1891b: 12 type species *Leptura sexmaculata* Linnaeus, 1758

*japonica* Tamanuki, 1942: 179 (*Strangalia*) A: JA

*parallelopipeda* Motschulsky, 1860b: 146 (*Grammoptera*)  
 E: NT A: ES FE JA MG NC SC WS “Korea”

*abbreviata* Motschulsky, 1875: 143 (*Grammoptera*)

*multidisjuncta* Pic, 1914c: 5

*shirarakensis* Matsumura, 1911a: 137 (*Leptura*)

and (p. 103)

genus *Judolia*, nomen dubium

*rufimembris* Pic, 1917g: 3 (*Leptura*) A: “Siberia or.”

MUST BE (p. 102):

**genus *Judolia* Mulsant, 1863: 496** type species *Leptura sexmaculata* Linnaeus, 1758

*Judolia* Pic, 1891b: 12 type species *Leptura sexmaculata* Linnaeus, 1758



*japonica* Tamanuki, 1942: 179 (*Strangalia*) A: JA  
*parallelopipeda* Motschulsky, 1860b: 146 (*Grammoptera*)  
 E: NT A: ES FE JA MG NC SC WS “Korea”  
*abbreviata* Motschulsky, 1875: 143 (*Grammoptera*)  
*multidisjuncta* Pic, 1914c: 5  
*rufimembris* Pic, 1917g: 3 (*Leptura*)  
*shirarakensis* Matsumura, 1911a: 137 (*Leptura*)

## NOTE:

The type investigation of *Leptura (Judolia) sexmaculata* var. *rufimembris* Pic, 1917 preserved in Pic’s collection in Paris allows to identify the half-colored specimen (a female) as *Judolia parallelopipeda* (Motschulsky, 1860).

## 11. PAGE 104

## PRINTED:

*duodecimguttata duodecimguttata* Fabricius, 1801b: 353 A:  
 ES FE FUJ HEI HEN JA JIL KZ MG NC NMO QIN SC  
 SCH SHX WS ZHE  
*bisbijuncta* Pic, 1904d: 14  
*kapfereri* Pic, 1912j: 89  
*mediojuncta* Pic, 1902d: 10 (*Strangalia*)  
*mediosemijuncta* Pic, 1927e: 13 (*Strangalia*)  
*subobliterata* Pic, 1927e: 10 (*Strangalia*)  
*duodecimguttata rufoannulata* Pic, 1933b: 26 (*Strangalia*)  
 A: SCH  
*fisheriana* Gressitt, 1938a: 45 A: FUJ HUB SCH

## MUST BE:

*duodecimguttata* Fabricius, 1801b: 353 A: ES FE FUJ HEI  
 HEN JA JIL KZ MG NC NMO QIN SC SCH SHX WS  
 ZHE  
*bisbijuncta* Pic, 1904d: 14  
*kapfereri* Pic, 1912j: 89  
*mediojuncta* Pic, 1902d: 10 (*Strangalia*)  
*mediosemijuncta* Pic, 1927e: 13 (*Strangalia*)  
*subobliterata* Pic, 1927e: 10 (*Strangalia*)  
 ...

*rufoannulata* Pic, 1933b: 26 (*Strangalia*) A: FUJ HUB SCH  
*fisheriana* Gressitt, 1938a: 45

## NOTE:

According to Heyrovsky [1934] *Leptura duodecimguttata* ssp. *rufoannulata* (Pic, 1933) is in fact a good species.

A comparison of the original description of *Strangalia duodecimguttata* var. *rufoannulata* Pic, 1933 (redescribe as *Leptura rufoannulata* by Heyrovsky, 1934) show its identity to *Leptura fisheriana* Gressitt, 1938, so *Leptura rufoannulata* (Pic, 1933) = *Leptura fisheriana* Gressitt, 1938, **syn.n.**

## 12. PAGE 104

## PRINTED:

*gradatula* Holzschuh, 2006a: 216 A: SHA

## MUST BE:

*gradatula* Holzschuh, 2006a: 216 A: GAN SCH SHA  
 According to the original description.

## 13. PAGE 108

## PRINTED:

*erraticus erraticus* Dalman, 1817a: 490 (*Leptura*) E: AB AL  
 AR AU BH BU BY CR CT CZ FR GE GG GR HU IT MC  
 MD PL RO SK SL SP ST SZ TR UK YU A: IN SY TR

## MUST BE:

*erraticus erraticus* Dalman, 1817a: 490 (*Leptura*) E: AB AL  
 AR AU BH BU BY CR CT CZ FR GE GG GR HU IT KZ  
 MC MD PL RO SK SL SP ST SZ TR UK YU A: IN KZ  
 SY TR

## NOTE:

See: Kostin [1973] and Zhuravlev [1914].

## 14. PAGE 111

## PRINTED:

**genus *Pseudovadonia* Lobanov, Danilevsky & Murzin, 1981: 787** type species *Leptura livida* Fabricius, 1777  
*livida desbrochersi* Pic, 1891k: xvi (*Vadonia*) E: AB AR GG  
 A: TR  
*corallipes* Reitter, 1894f: 247 (*Leptura*)  
*livida livida* Fabricius, 1777: 233 (*Leptura*) E: AL AR AU  
 BE BU BH BY CR CT CZ DE EN FR GB GE GG GR HU  
 IR IT KZ LA LT MC MD NE NL NT PL PT RO SK SL  
SP ST SZ TR UK YU A: ES IS KZ LE TR SY WS XIN  
*bicarinata* N. Arnold, 1869: 137 (*Grammoptera*)  
*pastinacae* Panzer, 1795: 275 (*Leptura*)  
*pecta* K. Daniel & L. Daniel, 1891: 38 (*Leptura*)

## MUST BE:

**genus *Pseudovadonia* Lobanov, Danilevsky & Murzin, 1981: 787** type species *Leptura livida* Fabricius, 1777  
*livida bicarinata* N. Arnold, 1869: 137 (*Grammoptera*) E:  
 AB AR BY CT EN GG KZ LA LT NT PL ST UK A: ES  
 IR KZ KI TR WS XIN  
*livida desbrochersi* Pic, 1891k: xvi (*Vadonia*) E: AB AR GG  
 A: TR  
*corallipes* Reitter, 1894f: 247 (*Leptura*)  
*livida livida* Fabricius, 1777: 233 (*Leptura*) E: AL AU BE  
 BU BH CR CZ DE FR GB GE GR HU IR MC MD NE NL  
 ?PT RO SK SL ?SP SZ TR UK YU A: ?IS ?LE ?TR ?SY  
*pastinacae* Panzer, 1795: 275 (*Leptura*)  
*livida pecta* K. Daniel & J. Daniel, 1891: 38 (*Leptura*) E: IT

## NOTE:

*P. l. livida* (Fabricius, 1777), described from near Kiel (Germany), is characterized by strongly erect straight dorsal pronotal setae (see “Galleri” www.cerambycidae.net). Such form of pronotal pubescence can be observed in most populations from West Europe (available specimens are from: France, Germany, Czechia, Hungary, Moldavia, West Ukraine — Transcarpathia, Bulgaria, Greece), as well as from West Turkey (Antalia).

*P. l. bicarinata* (N. Arnold, 1869), described from near Mogilev (East Belorussia) is characterized by obliquely erect dorsal pronotal setae (see “Gallery” www.cerambycidae.net). Such form of pronotal pubescence can be observed all over Russia, in most of Ukraine territory, in Baltic countries, in Poland, in Transcaucasia with neighbour regions of Turkey, in Kazakhstan and Kirgizia. “*Leptura l. var. bicarinata* (N. Arnold, 1869)” was already accepted as a taxon for European Russia [K. Daniel & J. Daniel, 1891].

The type locality of *P. livida pecta* (K. Daniel & J. Daniel, 1891) was not definitely mentioned in the original description, neither holotype was designated. The authors called the corresponding form as “Bozener Form” and specially described specimens from near “Bozen” — now Bolzano in North Italy (Trentino — Alto Adige). But they included in the area of their “*Leptura livida* var. *pecta*”: Piedmont (Italy), Digne (France), Lugano (Switzerland), as well as Spain, “Kleinasien”, “Kaukasus” and Siberia (“Irkutsk”), so the lectotype from near Bolzano is necessary to be designated for the fixation of the taxon. Specimens from North Italy (available specimens are from Bolzano and Trento – coll. of M. Egger; Fanano near Modena – author’s collection) are characterized by strongly recumbent dorsal pronotal setae (see “Galleri” www.cerambycidae.net). Such form of pronotal pubescence is not known to me in any other area. It seems to be an endemic of North Italy. The specimens from Central and South Italy have obliquely erect dorsal pronotal setae and so similar to *P. l. bicarinata* and must be described as another subspecies,

as well as populations from Iberian Peninsula and Near East must be also described as new subspecies. The record of the species for Iran was published by Gfeller [1972]: “Chalus (Now Shar) Mazandaran”

15. PAGE 113

PRINTED:

*bifasciata bifasciata* O. F. Müller, 1776: 93 (*Leptura*) E: AB AL AR AU BE BH BU BY CR CT CZ EN FR GE GG GR HU IT LA LS LT LU MC MD NL NT PL PT RO SK SL SP ST SZ TR UK YU A: ES IN IQ KZ TR WS XIN

*albarracina* Wagner, 1927a: 45 (*Leptura*)

*cruciata* Olivier, 1795: 7 (*Leptura*)

*ferruginipes* Pic, 1895d: 76 (*Strangalia*)

*immaculata* Pic, 1889b: 55 (*Strangalia*)

*lanceolata* Mulsant & Rey, 1863: 177 (*Strangalia*)

*nigriventris* Pic, 1891b: 15 (*Strangalia*)

*sedakovi* Mannerheim, 1852b: 307 (*Stenura*)

*ustulata* Laicharting, 1784: 157 [HN] (*Leptura*)

*bifasciata limbiventris* Reitter, 1898a: 21 (*Strangalia*) E: GG A: TR

*bifasciata nigrosuturalis* Reitter, 1895a: 88 (*Strangalia*) A: LE SY TR

*hybridula* Reitter, 1901h: 188 (*Strangalia*) E: PT SP

*atriventris* Pic, 1905a: 8 (*Leptura*) [DA]

*atronotata* Pic, 1918d: 5

*intermedia* Holzschuh, 2006a: 219 E: GR

MUST BE:

*bifasciata bifasciata* O. F. Müller, 1776: 93 (*Leptura*) E: AB AL AR AU BE BH BU BY CR CT CZ EN FR GE GG GR HU IT LA LS LT LU MC MD NL NT PL PT RO SK SL SP ST SZ TR UK YU A: ES IN IQ KZ TR WS XIN

*albarracina* Wagner, 1927a: 45 (*Leptura*)

*cruciata* Olivier, 1795: 7 (*Leptura*)

*immaculata* Pic, 1889b: 55 (*Strangalia*)

*nigriventris* Pic, 1891b: 15 (*Strangalia*)

*sedakovi* Mannerheim, 1852b: 307 (*Stenura*)

*ustulata* Laicharting, 1784: 157 [HN] (*Leptura*)

*bifasciata ferruginipes* Pic, 1895d: 76 (*Strangalia*) A: TR

*bifasciata intermedia* Holzschuh, 2006a: 219 E: GR BG

*bifasciata lanceolata* Mulsant & Rey, 1863: 177 (*Strangalia*) E: FR SP

*bifasciata limbiventris* Reitter, 1898a: 21 (*Strangalia*) E: GG A: TR

*bifasciata nigrosuturalis* Reitter, 1895a: 88 (*Strangalia*) A: LE SY TR

*hybridula* Reitter, 1901h: 188 (*Strangalia*) E: PT SP

*atriventris* Pic, 1905a: 8 (*Leptura*) [DA]

*atronotata* Pic, 1918d: 5

NOTE:

A lot of specimens of *Stenurella bifasciata intermedia* Holzschuh, 2006a (including many females, which are not described yet) were collected by A. Napolov in Greece from South Peloponnesus (Mani Peninsula) to Struma valley in Bulgaria in May–June 2010. Relatively large pronotal punctation of *S. bifasciata intermedia* Holz. (the main distinguishing character of the species according to the original description) is really a little larger than in specimens from Central Europe, but just same as in many south populations from Italy to Caucasus. Females (see “Galleri” in www.cerambycidae.net) of *S. b. intermedia* Holz. are very similar to *S. b. bifasciata*, but black elytral design is a little reduced.

The taxon described as *Strangalia lanceolata* Mulsant & Rey, 1863 from «L’Espagne» on the base of females with elytra widely darkened along suture is a well formed Iberian subspe-

cies *S. bifasciata* ssp. *lanceolata* (Mulsant & Rey, 1863). *S. b. lanceolata* penetrates in South France. Two females with the label: «France, Pyrénées Orientales, Prades, 24–30.6.1986, Schimmel leg.» are preserved in my collection.

Populations, which contain specimens with pale-orange legs, represent a well delimited subspecies described as *Strangalia bifasciata* var. *ferruginipes* Pic, 1895 from «Bitlis». *Stenurella bifasciata* ssp. *ferruginipes* (Pic, 1895) is represented in my materials from Mardin (Hop Geçidi) and Bitlis (Tatvan environs).

16. PAGE 116

PRINTED:

*semirufa* Kraatz, 1880b: 376 (*Leptura*)

MUST BE:

*semirufula* Kraatz, 1880b: 376 (*Leptura*)

NOTE:

According to the original description.

17. PAGE 116

PRINTED:

*variicornis* Dalman, 1817a: 482 (*Leptura*) E: CT NT PL UK A: ES FE KZ MG NC SC WS

MUST BE:

*variicornis* Dalman, 1817a: 482 (*Leptura*) E: BY CT EN LA LT NT PL UK A: ES FE JP KZ MG NC NE NO SC WS

18. PAGE 116

PRINTED:

genus *Stictoleptura* nomen dubium

*silbermanni* Lefebvre, 1835: 303 (*Leptura*) A: “Syrie Mont Liban”

NOTE:

According to Sama et al. [2010] *Leptura silbermanni* Lefebvre, 1835 was the first name for *Stictoleptura heydeni* (Ganglbauer, 1888); it was proposed to be regarded as “nomen oblitum” following the Article 23.9.2 of ICZN. But the obligated condition of that article is the existence of 25 publications with the protected name by at least 10 authors for the last 50 years. Without such a condition the name *Stictoleptura silbermanni* (Lefebvre, 1835) must be accepted as valid. In fact the type of *Leptura silbermanni* Lefebvre, 1835 is not known, so the real nature of the described taxon rests doubtful, and for the stability of the current nomenclature it is better to leave it as “nomen dubium”.

19. PAGE 117

PRINTED:

*bipunctata adusta* Kraatz, 1859: 97 (*Leptura*) E: HU MC RO SK SL

*litigiosa* Mulsant, 1863: 564

*bipunctata bipunctata* Fabricius, 1781: 245 (*Leptura*) E: CT ST A: KZ WS

*beckeri* Pic, 1941d: 14 (*Leptura*)

*fischeri* Zubkov, 1829: 167 (*Leptura*)

*laterimaculata* Motschulsky, 1875: 142 (*Leptura*)

*sareptana* Pic, 1941d: 15

*bipunctata mulsantiana* Plavilstshikov, 1936: 341 (*Leptura*) E: MD RO ST UK

*bipunctata puchneri* Holzschuh, 2007: 183 E: ST UK

*bipunctata steveni* Sperk, 1835: 158 (*Leptura*) E: HU MD SK SL UK

*bilitigiosa* Pic, 1941d: 15 [DA]

*globicollis* Desbrochers des Loges, 1870c: 127 (*Leptura*) [DA]

*rufonotata* Pic, 1926d: 10 (*Leptura*)

## MUST BE:

- bipunctata adusta* Kraatz, 1859: 97 (*Leptura*) E: HU MC RO SK SL  
*litigiosa* Mulsant, 1863: 564  
*rufonotata* Pic, 1926d: 10 (*Leptura*)  
*bipunctata bipunctata* Fabricius, 1781: 245 (*Leptura*) E: CT ST A: KZ WS  
*beckeri* Pic, 1941d: 14 (*Leptura*)  
*fischeri* Zubkov, 1829: 168 (*Leptura*)  
*sareptana* Pic, 1941d: 15  
*bipunctata globicollis* Desbrochers des Loges, 1870c: 127 (*Leptura*) E: MD RO ST UK  
*mulsantiana* Plavilstshikov, 1936: 341 (*Leptura*)  
*bilitigiosa* Pic, 1941d: 15  
*bipunctata laterimaculata* Motschulsky, 1875: 141 (*Leptura*) E: UK (Crimea)  
*puchneri* Holzschuh, 2007: 183  
*bipunctata steveni* Sperk, 1835: 158 (*Leptura*) E: HU MD SK SL ST UK

## NOTE:

*Leptura laterimaculata* Motschulsky, 1875 was described from Crimea (“Tauride”) on the base of a male with black elytra, each with small lateral yellow spots. The holotype (see “Galleri” in www.cerambycidae.net) of the taxon (head, prothorax and several legs are absent) is preserved in Zoological Museum of Moscow University. It is undoubtedly a form of *Vadonia bipunctata* (because of typical elytral design and numerous erect setae on hind femur), which was recently described as *V. puchneri* Holzschuh, 2007. So, *Leptura laterimaculata* Motschulsky, 1875 = *Vadonia puchneri* Holzschuh, 2007, **syn.n.**

*Leptura (Vadonia) bipunctata* var. *rufonotata* Pic, 1926d was described from “Hongrie”

*Leptura globicollis* Desbrochers des Loges, 1870c was described from “Kustendjé (Turquie)” [Constanța, Romania], so very close to Izmail — the the type locality of *Leptura (Vadonia) bipunctata mulsantiana* Plavilstshikov, 1936, so *Leptura globicollis* Desbrochers des Loges, 1870 = *L. (Vadonia) bipunctata mulsantiana* Plavilstshikov, 1936.

*Leptura steveni* Sperk, 1835 was described from Podolia (north-west Ukraine). *Vadonai bipunctata steveni* Sperk, 1835 is distributed eastwards to North Caucasus. The specimens of the subspecies have very rough pronotal punctation similar to *V. unipunctata* and *V. b. laterimaculata* Motschulsky, 1875. They differs from the later by usual elytral design: pale elytral color is often very dark and elytral apices are often contrastly black. Males of *V. b. steveni* Sperk, 1835 (as well as in *V. b. globicollis* Desbrochers des Loges, 1870) have sometimes only one spine in hind tibia. Such structure can be also observed in all other subspecies of *V. bipunctata*, but very rare.

## 20. PAGE 117

## PRINTED:

*bitlisiensis* Chevrolat, 1882: 59 E: AR A: TR

## MUST BE:

*bitlisiensis* Chevrolat, 1882: 59 E: AR AB A: TR

## NOTE:

The species was recorded for Nakhichevan Republic of Azerbajdzhan by Plavilstshikov [1948].

## 21. PAGE 117

## PRINTED:

*cribricollis* Pic, 1889b: 20 [mispaginated: 5]

## MUST BE:

*cribricollis* Pic, 1889b: 20 [mispaginated: 5] (*Leptura*)

## 22. PAGE 118

## PRINTED:

*unipunctata unipunctata* Fabricius, 1787: 157 (*Leptura*) [NP]  
 E: AB AR AU BH BU CR CT CZ FR GG GR HU IT KZ MD PL RO SB SK SL SP ST TR UK A: KZ TR  
*obscuripilosa* Pic, 1892q: lxxxiv  
*pilosa* Forster, 1771: 44 (*Leptura*) [NO]

## MUST BE:

*unipunctata unipunctata* Fabricius, 1787: 157 (*Leptura*) [NP]  
 E: AB AR AU BH BU CR CT CZ FR GG GR HU IT KZ MD PL RO SB SK SL SP ST TR UK A: KZ TR  
*obscuripilosa* Pic, 1892q: lxxxiv  
*pilosa* Forster, 1771: 44 (*Leptura*) [NO]  
*uninstigmata* Pic, 1891b: 9 (*Leptura*)

## NOTE:

A taxonomy Act for the protection of *Leptura unipunctata* Fabricius, 1787 is not published in the Catalog. According to Sama [2008]: *Leptura unipunctata* Fabricius, 1787 (“nomen protectum”) = *Leptura pilosa* Forster, 1771 (“nomen oblitum”). But 25 publications by 10 authors for the last 50 years [ICZN Art. 23.9.1.2] were not listed, so the action can not be regarded as valid.

Here is the list of 25 publications, where the name “*unipunctata* Fabricius, 1787” was used as valid: Bartenev, 2004, 2009; Danilevsky, 2009; Danilevsky & Miroshnikov, 1985; Esjunin & Kozminykh, 1992; Zagaikevich, 1991, 1979; Isaev, 2007; Isaev et al., 2004; Isaev & Magdeev, 2003; Kaliuzhnaia et al., 2000; Kostin, 1968, 1973; Kryzhanovsky, 1974; Lobanov et al., 1981; Magdeev, 1996, 2003; Martynov & Pisarenko, 2004; Mikhailov, 1999; Miroshnikov, 1998; Naumov, 1994; Negrobov et al., 2005; Pisarenko, 1999; Plavilstshikov, 1965; Romadina, 1954; Shapovalov et al., 2006.

## 23. PAGE 119

## PRINTED:

*dentipes* Mulsant, 1842a: 209

## MUST BE:

*dentipes* Mulsant, 1842a: 209 (*Toxotus*)

## 24. PAGE 124

## PRINTED:

*carneola* Schrank, 1798: 698 (*Leptura*)

## MUST BE:

*carneola* Schrank, 1798: 696 (*Leptura*)

## 25. PAGE 124

## PRINTED:

*minuta* Gebler, 1832: 69 (*Pachyta*) A: ES FE GUX HEB HEI HEN JA JIL JIX LIA NC NIN NMO SC SHA SHN SHX ZHE  
*criocerina* Bates, 1873: 194 (*Acmaeops*)  
*japonica* Pic, 1907d: 20 (*Acmaeops*)

## MUST BE:

*minuta criocerina* Bates, 1873: 194 (*Acmaeops*) A: JA  
*japonica* Pic, 1907d: 20 (*Acmaeops*)  
*minuta minuta* Gebler, 1832: 69 (*Pachyta*) A: ES FE GUX HEB HEI HEN JIL JIX LIA NC NIN NMO SC SHA SHN SHX ZHE

## NOTE:

The synonyms of *Pachyta minuta* Gebler, 1832 (described from Transbaicalia) and *Acmaeops criocerina* Bates, 1873 (described from Japan) were generally accepted by all recent Japan publications, but continental and Japan populations are rather different. First of all most of Japan specimens have partly or totally red abdomen, while such form is



unknown in Russia (from Baikal to Kamchatka and Sakhalin). It is known from South Korea as rare aberration. So, at least subspecies rank of two names must be accepted. But in fact *Dinoptera minuta* (Gebler, 1832) and *Dinoptera criocerina* (Bates, 1873) must be different species, because *D. minuta* penetrates far northwards in the continent (to about Kamchatka Peninsula), but *D. criocerina* is known from South Japan only (absent in Hokkaido).

26. PAGE 133

PRINTED:

*nubecula* Bergsträsser, 1778: 25 (*Cerambyx*)

MUST BE:

*nubecula* Bergsträsser, 1778: 26 (*Cerambyx*)

27. PAGE 134

PRINTED:

*meridianus* Linnaeus, 1758: 398 (*Leptura*) E: AL AU BE BH BU BY CR CT CZ DE EN FI FR GB GE GR HU IR IT LA LS LT MC MD NL NR NT PL RO SK SL SP ST SV SZ TR UK YU A: ES KZ NC SC WS

MUST BE:

*meridianus* Linnaeus, 1758: 398 (*Leptura*) E: AL AU BE BH BU BY CR CT CZ DE EN FI FR GB GE GR HU IR IT LA LS LT MC MD NL NR NT PL RO SK SL SP ST SV SZ TR UK YU A: ES KZ WS

NOTE:

Several old records of *Stenocorus meridianus* for Korea were connected with *S. amurensis*.

The records of *S. meridianus* for Gansu and Shaanxi [Hua, 2002] were quite doubtful and adequately not accepted in the Catalog.

28. PAGE 135

PRINTED:

*ambustum* Heyden, 1877a: 394

NOTE:

The name was introduced as *Rhamnusium bicolor* var. *ambustum* Heyden, 1877 among other variations from same locality: "Drei Linden" bei Soden, so it must be excluded as unavailable.

29. PAGE 137

PRINTED:

*ferus* Mulsant, 1839: 64 (*Criocephalus*) E: AB AR AU BE BH BU BY CR CT CZ DE EN FR GB GE GG GR HU IR IT LA LT LU MA MC MD NL NT PL PT RO SK SL SP ST SV SZ UK N: AG MO MR TU A: ES CY FE IS JO KZ TR WS

MUST BE:

*ferus* Mulsant, 1839: 64 (*Criocephalus*) E: AB AR AU BE BH BU BY CR CT CZ DE EN FR GB GE GG GR HU IR IT LA LT LU MA MC MD NL NT PL PT RO SK SL SP ST SV SZ UK N: AG MO MR TU A: ES CY FE IS JO KZ NO NE TR WS

30. PAGE 143

PRINTED:

genus *Pufujia* Holzschuh, 1965: 16 type species *Pufujia luteosignata* Pu, 1991

MUST BE:

genus *Pufujia* Holzschuh, 1995: 16 type species *Nortia luteosignata* Pu, 1991

31. PAGE 153

PRINTED:

*elbursense* Holzschuh, 1977a: 128

MUST BE:

*elbursense* Holzschuh, 1977a: 128

32. PAGE 155

PRINTED:

*macropus* Germar, 1824: 514 (*Callidium*) E: AB AR AU BH BU BY CR CT CZ GE GG HU LA MD PL RO SK SL ST SZ TR UK YU A: IN TR

MUST BE:

*macropus* Germar, 1824: 514 (*Callidium*) E: AB AR AU BH BU BY CR CT CZ GE GG HU LA LT MD PL RO SK SL ST SZ TR UK YU A: IN TR

NOTE:

*Ropalopus macropus* was recorded for Lithuania by Inokaitis [2004, 2009].

33. PAGE 156

PRINTED:

*johannis johannis* Baeckmann, 1922: 24 A: KI XIN

MUST BE:

*johannis johannis* Baeckmann, 1922: 24 A: KI

NOTE:

*Turanium johannis johannis* Baeckmann, 1922 (described from Talas Ridge) was recorded for Xinjiang (Sinkiang) by Hua [2002]. The species is impossible for China. The record could be connected with *T. badenkoi* Danilevsky, 2001e described from the north slope of Zailiysky Alatau, or with a new species.

34. PAGE 156

PRINTED:

tribe Callidiopini Lacordaire, 1868

genus *Ceresium* Newman, 1842d: 322 type species *Ceresium varipilum* Newman, 1842

*Diatomocephala* Blanchard, 1853: 266 type species *Diatomocephala maculicollis* Blanchard, 1853

*Paraceresium* Matsushita, 1932: 71 type species *Paraceresium saipanicum* Matsushita, 1932

*Pneumida* J. Thomson, 1864a: 191 type species *Pneumida argenteofasciata* J. Thomson, 1864

NOTE:

According to Holzschuh [1995: 16] *Pneumida* J. Thomson, 1864a is a valid name in Cerambycini.

35. PAGE 157

PRINTED:

genus *Parasalpinia* Hayashi, 1962c: 4 type species *Parasalpinia kojimai* Hayashi, 1962

*kojimai* Hayashi, 1962c: 4 A: JA (Ryukyus) TAI

*laosensis* Gressitt & Rondon, 1970: 102 (*Salpinia*) A: YUN

**ORR**

genus *Salpinia* Pascoe, 1869: 536 type species *Salpinia diluta* Pascoe, 1869

*laosensis* Gressitt & Rondon, 1970: 102 A: HAI YUN **ORR**

MUST BE:

genus *Parasalpinia* Hayashi, 1962c: 4 type species *Parasalpinia kojimai* Hayashi, 1962

*kojimai* Hayashi, 1962c: 4 A: JA (Ryukyus) TAI

*laosensis* Gressitt & Rondon, 1970: 102 (*Salpinia*) A: HAI

**YUN ORR**

NOTE:

Genus *Salpinia* Pascoe, 1869 absent in Palaearctic Region. According to Niisato [2002]: "*Salpinia laosensis* agrees with the characteristics of *Parasalpinia* and should be transferred to that genus."

## 36. PAGE 159

## PRINTED:

*cerdo cerdo* Linnaeus, 1758: 392 **E**: AB AL AR AU BE BH BU BY CR CT CZ FR GBi GE GG GR HU IR IT LA LU MA MC MD NL PL PT RO SK SL SP ST SZ TR UK YU N: MO **A**: IN IQ IS JO LE SY TR

## MUST BE:

*cerdo cerdo* Linnaeus, 1758: 392 **E**: AB AL AR AU BE BH BU BY CR CT CZ FR GBi GE GG GR HU IR IT LA LU MA MC MD NL PL RO SK SL ST SV SZ TR UK YU N: MO **A**: IN IQ IS JO LE SY TR

## 37. PAGE 159

## PRINTED:

*manderstjaer-nae* Mulsant & Godart, 1855a: 180

## MUST BE:

*manderstjernae* Mulsant & Godart, 1855b: 280 [= 1855a: 180]

## 38. PAGE 167

## PRINTED:

*gratiosus gratiosus* Marseul, 1868: 203 (*Clytus*) **A**: IS LE SY *gratiosus sparsus* Reitter, 1886: 67 (*Clytus*) **A**: TR

## MUST BE:

*gratiosus gratiosus* Marseul, 1868: 203 (*Clytus*) **A**: IS LE *gratiosus sparsus* Reitter, 1886: 67 (*Clytus*) **E**: GR **A**: SY TR

## NOTE:

According to Sama et al. [2010]: “La forme type, connue du Liban, est remplacée au Sud de la Turquie, en Syrie et dans l’île de Rhodes: environs de Arhagelos (espèce nouvelle pour la faune de Grèce et de l’Europe) par la sous-espèce *C. gratiosus sparsus* (Reitter, 1886) à coloration élytrale entièrement noire.”

## 39. PAGE 167

## PRINTED:

*latofasciatus* Motschulsky, 1861b: 41 (*Clytus*) **A**: ES FE FUJ GAN HEB HEI HEN JIL LIA MG NC NMO SC SHA SHN SHX ZHE

*chasanensis* Tsherepanov, 1982a: 175

*motschulskyi* Ganglbauer, 1887a: 135 (*Clytanthus*)

## MUST BE:

*motschulskyi* Ganglbauer, 1887a: 135 (*Clytus*) [RN] **A**: ES FE FUJ GAN HEB HEI HEN JIL LIA MG NC NMO SC SHA SHN SHX ZHE

*chasanensis* Tsherepanov, 1982a: 175

*latofasciatus* Motschulsky, 1861b: 41 (*Clytus*) [HN]

## NOTE:

*Clytus latofasciatus* Motschulsky, 1861b is a junior homonym of *Clytus latofasciatus* Fischer von Waldhein, 1832. See Art. 58.12 about different connecting vowels in compound words.

## 40. PAGE 169

## PRINTED:

*varius varius* O. F. Müller, 1766: 188 (*Leptura*) **E**: AB AL AR AU BH BU BY CR CT CZ FR GBi GE GG GR HU IT LS LT MA MC MD NL PL RO SK SL SP ST SZ TR UK YU **A**: JIA KZ TR WS

## MUST BE:

*varius varius* O. F. Müller, 1766: 188 (*Leptura*) **E**: AB AL AR AU BH BU BY CR CT CZ FR GBi GE GG GR HU IT LS LT MA MC MD NL PL RO SK SL SP ST SZ TR UK YU **A**: KZ TR WS

## 41. PAGE 178

## PRINTED:

**genus *Rhaphuma* Pascoe, 1858: 240** type species *Rhaphuma placida* Pascoe, 1858

*Arcyophorus* Gemminger, 1872: 2938 type species *Arcyophorus histrio* Chevrolat, 1863

*Arcyophorus* Chevrolat, 1863b: 287 type species *Arcyophorus histrio* Chevrolat, 1863

*Raphuma* J. Thomson, 1864: 192 type species *Clytus quadricolor* Laporte & Gory, 1836

## MUST BE:

**genus *Rhaphuma* Pascoe, 1858: 240 [RN]** type species *Clytus quadricolor* Laporte & Gory, 1836

*Arcyophorus* Gemminger, 1872: 2938 [unjustified emendation]

*Arcyophorus* Chevrolat, 1863b: 287 type species *Arcyophorus histrio* Chevrolat, 1863

*Raphuma* J. Thomson, 1861: 221 [unjustified emendation]

*Rhaphium* A. White, 1855: 289 [HN] type species *Clytus quadricolor* Laporte & Gory, 1836

## 42. PAGE 180

## PRINTED:

**genus *Sclethrus* Newman, 1842a: 247** type species *Ibidion amoenum* Gory, 1833

## MUST BE:

**genus *Sclethrus* Newman, 1842a: 247** type species *Sclethrus newmany* Chevrolat, 1863

*Neocollyroides* Schultze, 1920:196 type species *Neocollyroides macgregory* Schultze, 1920

## NOTE:

According to Han & Niisato [2009].

## 43. PAGE 186

## PRINTED:

*campestris* Faldermann, 1835c: 435 (*Callidium*) **E**: CT MD PLi RO ST UK **A**: ANH ES FE GAN GUI HEB HEI HEN HUB HUN IN JA JIA JIL JIX KI KZ LIA MG NC NMO QIN SC SCH SHA SHN SHX TD TM UZ XIN XIZ YUN ZHE **ORR**

## MUST BE:

*campestris* Faldermann, 1835c: 435 (*Callidium*) **E**: AB AR CT CZ MD PLi RO SK ST UK **A**: ANH ES FE GAN GUI HEB HEI HEN HUB HUN IN JA JIA JIL JIX KI KZ LIA MG NC NMO QIN SC SCH SHA SHN SHX TD TM UZ XIN XIZ YUN ZHE NARi **ORR**

## NOTE:

The species was recorded for Canada [Grebennikov et al., 2010].

## 44. PAGE 190

Missing name: *Molorchus abieticola* Holzschuh, 2007: 218

Must be accepted as: *Molorchus (Coenoptera) abieticola* Holzschuh, 2007: 218 **E**: TR

## 45. PAGE 190

## PRINTED:

*tenuitarsis* Holzschuh, 1981: 97 (*Molorchus*) **A**: TR *sterbai* Adlbauer, 1988: 277 (*Molorchus*)

## MUST BE:

*sterbai* Adlbauer, 1988: 277 **A**: LE

*azri* Sama, Rapuzzi & Kairouz, 2010: 151 (*Glaphyra*)

*tenuitarsis* Holzschuh, 1981: 97 **A**: TR

## NOTE:

*Molorchus kiesenwetteri* ab. *sterbai* Heyrovsky, 1936 described from Lebanon was not available; it was validated as



“*Molorchus sterbai* Heyrovsky, 1936” by Adlbauer [1988] (without description — just a name in the list of his specimens — so, fits to the Article 13.1.2). The holotype of the name is a specimen of ab. *sterbai* from Heyrovsky’s collection, but not a specimen identified so by Adlbauer.

The taxon was described once more as *Glaphyra azri* Sama, Rapuzzi & Kairouz, 2010 also from Lebanon, so *Molorchus sterbai* Adlbauer, 1988 = *Glaphyra azri* Sama, Rapuzzi & Kairouz, 2010, **syn.n.**

46. PAGE 192

PRINTED:

**genus *Chinobrium* Gressitt, 1937c: 449** type species *Chinobrium mediofasciatum* Gressitt, 1937  
*aegrotum* Holzschuh, 1982a: 67 (*Obrium*) A: NP  
*mediofasciatum* Gressitt, 1937c: 449 A: JIX **ORR**  
*opacum* Holzschuh, 1984c: 348 (*Stenhomalus*) A: BT **ORR**

MUST BE:

The correct combination is:

*Obrium aegrotum* Holzschuh, 1982a: 67 A: NP

47. PAGE 195

PRINTED:

**genus *Allotraeus* Bates, 1877: 36** type species *Allotraeus sphaerioninus* Bates, 1877

...

**subgenus *Pseudallotraeus* Pic, 1923a: 13** type species *Pseudallotraeus rufescens* Pic, 1923

*rufescens* Pic, 1923a: 13 A: JA SC TAI

and

**genus *Nysina* Gahan, 1906a: 153** type species *Sphaerion orientale* A. White, 1853

*Neosphaerion* Schwarzer, 1925a: 21 type species *Neosphaerion asiaticum* Schwarzer, 1925

*Pseudallotraeus* Pic, 1923a: 13 type species *Pseudallotraeus rufescens* Pic, 1923

...

*rufescens* Pic, 1923a: 13 (*Pseudallotraeus*) A: JA TAI

*japonica* K. Ohbayashi, 1936a: 13

NOTE:

So, the name *Pseudallotraeus* Pic, 1923 was used twice in different genera with different rank. According to Niisato [2007]: *Pseudallotraeus* Pic, 1923 is a synonym of *Nysina* Gahan, 1906, which is a subgenus of *Allotraeus* Bates, 1877, so the second case of the Catalog is acceptable.

48. PAGE 196

PRINTED:

*halodendri ephippium* Steven & Dalman, 1817: 157 (*Cerambyx*) E: ST UK A: KZ

MUST BE:

*halodendri ephippium* Steven & Dalman, 1817: 157 (*Cerambyx*) E: AL BU ST UK KZ A: KZ

NOTE:

*Asias halodendri* was recorded for Albania [Muraj, 1960] and Bulgaria [Angelov, 1995].

49. PAGE 202

PRINTED:

*alpina alpina* Linnaeus, 1758: 392 (*Cerambyx*) E: AB AL AU AR BH BU BY CR CT CZ FR GE GG GR HU LS MC PL RO SK SL SP ST SV SZ UK YU A: TR

MUST BE:

*alpina alpina* Linnaeus, 1758: 392 (*Cerambyx*) E: AB AL AU AR BH BU BY CR CT CZ FR GE GG GR HU IT LS MC PL RO SK SL SP ST SZ UK YU A: TR

50. PAGE 207

PRINTED:

*Neacanthocinus* Dillon, 1956: 230 type species *Acanthocinus obsoletus* Olivier, 1837

MUST BE:

According to the original publication,  
*Neacanthocinus* Dillon, 1956: 230 type species *Acanthocinus obsoletus* Olivier, 1837

51. PAGE 215

PRINTED:

*kirbyi* Gyllenhal, 1817: 186 (*Saperda*) E: AB AL AR BH BU CR FR GG GR HU IT MC MD RO SK SP ST TR UK YU A: IN IS SY TM

MUST BE:

*kirbyi* Gyllenhal, 1817: 186 (*Saperda*) E: AB AL AR BH BU CR CT FR GG GR HU IT MC MD RO SK SP ST TR UK YU A: IN IS SY TM TR

NOTE:

The species is well known from Samara and Ulianovsk regions of Central Russia, as well as from Turkey.

52. PAGE 227

PRINTED:

*sokotrensis* Téocchi, Jiroux & Sudre, 2004: 22 (*Arasbosybrinus*) A: YE (Suqutra)

MUST BE:

*sokotrensis* Téocchi, Jiroux & Sudre, 2007: 22 (*Arasbosybrinus*) A: YE (Suqutra)

53. PAGE 239

PRINTED:

**genus *Diastocera* J. Thomson, 1857: 183** type species *Lamia tricincta* Duncan, 1835 (= *Lamia wallichi* Hope, 1831)  
*Thysia* J. Thomson, 1860: 96 type species *Lamia wallichi* Hope, 1831

*Thysiotus* J. Thomson, 1868: 201 [RN] type species *Lamia wallichi* Hope, 1831

MUST BE:

**genus *Thysia* J. Thomson, 1860: 96** type species *Lamia wallichi* Hope, 1831

*Thysiotus* J. Thomson, 1868: 201 [RN] type species *Lamia wallichi* Hope, 1831

NOTE:

The author of *Diastocera* is Dejean [1835: 342] with *Lamia trifasciata* Fabricius, 1775: 174 as type species (monotypy) — African genus. *Lamia trifasciata* Fabricius, 1775 is the type species of African *Analeptes* Gistel, 1848: 430

So, *Diastocera* Dejean, 1835: 342 = *Analeptes* Gistel, 1848: 430, **syn.n.** *Thysia* J. Thomson, 1860: 96 (type species *Lamia wallichi* Hope, 1831) is valid.

54. PAGE 239

PRINTED:

*savioi* Yen, 1933: 165

NOTE:

The reference to the publication by Yen [1933] absent in the References to the Catalog. According to Hua [2002] it was:

Yen Chia-hsien, 1932: A new species of Cerambycidae from Kwangsi. *Peking Natural History Bulletin* 7(2): 165–166.

55. PAGE 240

PRINTED:

*Tylophorus* Blessig, 1873: 213 type species *Tylophorus wulffusi* Blessig, 1873

- and  
*wulffiusi* Blessig, 1873: 215 (*Thylophorus*)
- MUST BE:  
*Thylophorus* Blessig, 1873: 213 type species *Thylophorus wulffiusi* Blessig, 1873
- and  
*wulffiusi* Blessig, 1873: 215 (*Thylophorus*)
- NOTE:  
According to the original description.
56. PAGE 242
- PRINTED:  
*fulvum fulvum* Scopoli, 1763: 53 (*Cerambyx*) E: AU CZ HU SK SL UK  
*freyi* Tippmann, 1958b: 158  
*frontale* Mulsant & Rey, 1863: 154  
*kulzeri* Tippmann, 1958b: 158  
*nigripenne* Fleischer, 1894: 121  
*sanguinolentum* Scopoli, 1772: 99 (*Prionus*)
- MUST BE:  
*fulvum fulvum* Scopoli, 1763: 53 (*Cerambyx*) E: AU CZ HU SK SL UK  
*freyi* Tippmann, 1958b: 158  
*frontale* Mulsant & Rey, 1863: 154  
*kulzeri* Tippmann, 1958b: 158  
*nigripenne* Fleischer, 1894: 121  
*rugipenne* Tippmann, 1958b: 155  
*sanguinolentum* Scopoli, 1772: 99 (*Prionus*)
57. PAGES 243, 246 AND 247
- PRINTED (p. 243):  
*arenarium marsicanum* Fracassi, 1905: 112 E: FR IT  
*florij* Breuning, 1942b: 126  
and (p.246)  
*etruscum etruscum* Rossi, 1790: 147 (*Lamia*) E: IT  
*apenninum* Depoli, 1926a: 25  
*apulium* Depoli, 1926a: 25  
*beieri* Pic, 1932d: 18  
*calabricum* Breuning, 1942b: 126  
*femoratum* Brullé, 1832: 259  
*florij* Breuning, 1942b: 126  
and (p. 247)  
*etruscum florij* Breuning, 1942b: 126 E: IT
- MUST BE (p. 247):  
*etruscum florij* Breuning, 1942b: 126 E: IT
- NOTE:  
According to Pesarini & Sabbadini [2007].
58. PAGES 245, 250
- PRINTED (p.245):  
*ciscaucasicum* Jakovlev, 1900b: 59 E: ST  
*borodini* Suvorov, 1915: 118  
and (p.250)  
*mokrzeckii* Jakovlev, 1902: 148 E: UK
- MUST BE (p.245):  
*ciscaucasicum abramovi* Lazarev, 2009: 14 E: ST (Taman)  
*ciscaucasicum ciscaucasicum* Jakovlev, 1900b: 59 E: ST  
*borodini* Suvorov, 1915: 118  
*ciscaucasicum mokrzeckii* Jakovlev, 1902: 148 E: UK (Krym)
59. PAGE 270
- PRINTED:  
**genus *Choeromorpha* Chevrolat, 1843: 613** type species *Choeromorpha pigra* Aurivillius, 1920  
**subgenus *Choeromorpha* Chevrolat, 1843: 613** type species *Choeromorpha pigra* Aurivillius, 1920

- MUST BE:  
**genus *Choeromorpha* Chevrolat, 1843: 613** type species *Choeromorpha pigra* Chevrolat, 1843  
**subgenus *Choeromorpha* Chevrolat, 1843: 613** type species *Choeromorpha pigra* Chevrolat, 1843
60. PAGE 279
- PRINTED:  
**genus *Cereopsius* Pascoe, 1862a: 344** type species *Cereopsius exoletus* Pascoe, 1862
- MUST BE:  
**genus *Cereopsius* Pascoe, 1857b: 105** type species *Cereopsius exoletus* Pascoe, 1857
61. PAGE 279
- PRINTED:  
*Falsapriona* Pic, 1925c: 3 type species *Falsapriona luteopubens* Pic, 1925
- MUST BE:  
*Falsapriona* Pic, 1925c: 3 type species *Falsapriona luteopubens* Pic, 1925
62. PAGE 282
- PRINTED:  
*Tibetobia szechenyana* Frivaldszky, 1892  
and  
*szechenyanus* Frivaldszky, 1892a: 119 (*Tibetobia*)
- MUST BE:  
*Tibetobia szechenyana* Frivaldszky, 1892  
and  
*szechenyianus* Frivaldszky, 1892a: 119 (*Tibetobia*)
63. PAGE 281
- PRINTED:  
**genus *Hoplothrix* Gahan, 1888a: 278** type species *Hoplothrix simplex* Gahan, 1888
- MUST BE:  
**genus *Hoplothrix* Gahan, 1888a: 278** type species *Hoplothrix simplex* Gahan, 1888
- NOTE:  
According to the original publication.
64. PAGE 284
- PRINTED:  
**genus *Paraepeotes* Breuning, 1938c: 182** type species *Paraepeotes breuningi* Pic, 1935  
*affinis* Breuning, 1938c: 183 A: UP **ORR**  
*albomaculatus* Gahan, 1888a: 272 (*Epepeotes*) A: BT "North India"  
*breuningi* Pic, 1935b: 16 (*Paraepeotes*) A: SCH XIZ **ORR**  
*guttatus* Guérin-Méneville, 1844: 242 (*Monohamus*) A: NP SCH "Himalaya" **ORR**  
*punctulatus* Westwood, 1848: 12 (*Monohammus*)  
*marmoratus* Pic, 1925a: 19 (*Monohammus*) A: YUN **ORR**  
*szetschuanicus* Breuning, 1969e: 188 A: SCH  
*westwoodi* Westwood, 1848: 12 (*Monohammus*) A: SD "Himalaya"
- MUST BE:  
**genus *Paraepeotes* Pic, 1935: 16** type species *Paraepeotes breuningi* Pic, 1935  
*Paraepeotes* Breuning, 1938c: 182 [unjustified emendation]  
*affinis* Breuning, 1938c: 183 (*Paraepeotes*) A: UP **ORR**  
*albomaculatus* Gahan, 1888a: 272 (*Epepeotes*) A: BT "North India"  
*breuningi* Pic, 1935b: 16 A: SCH XIZ **ORR**

*guttatus* Guérin-Méneville, 1844: 242 (*Monohammus*) A: NP SCH “Himalaya” **ORR**  
*punctulatus* Westwood, 1848: 12 (*Monohammus*)  
*marmoratus* Pic, 1925a: 19 (*Monohammus*) A: YUN **ORR**  
*szetschuanicus* Breuning, 1969e: 188 (*Parepepeotes*) A: SCH  
*westwoodi* Westwood, 1848: 12 (*Monohammus*) A: SD  
 “Himalaya”

65. PAGE 300

PRINTED:

*ressli* Demelt, 1963: 150 A: TR

NOTE:

The taxon was wrongly attributed to the subgenus *Obe-rea* s. str., but in fact it belongs to *Amaurostoma*.

66. PAGE 310 AND 311

PRINTED (p.310):

*diversiceps* Pic, 1931f: 259 A: BT NP SD YUN **ORR**

*lateraloides* Breuning, 1958f: 300

*rufoampliatius* Breuning, 1958f: 300

*subbicolor* Breuning, 1958f: 300

and (p.311)

*subbicolor* Breuning, 1958f: 300 [RN] A: NP **ORR**

*bicolor* Pic, 1929a: 30 [HN]

MUST BE (p. 310):

*diversiceps* Pic, 1931f: 259 A: BT NP SD YUN **ORR**

*lateraloides* Breuning, 1958f: 300

*rufoampliatius* Breuning, 1958f: 300

The second case is correct.

67. PAGE 819

PRINTED:

Pic M. 1890e: [new taxa]. *Bulletin de la Société Entomologique de France* **1889**: clxxvi–clxxvii.

MUST BE:

Pic M. 1890e: [new taxa]. *Bulletin de la Société Entomologique de France* **1889**: clxxv–clxxvi.

68. PAGE 801

PRINTED:

Müller J. 1907: Cerambycidae Dalmaciae.

MUST BE:

Müller J. 1907: Cerambycidae Dalmatiae.

69. PAGE 801

PRINTED:

Mulsant E. 1847d: Description de deux coléoptères nouveaux, constituant chacun une nouvelle coupe générique. *Annales des Sciences Physiques et Naturelles, d’Agriculture et d’Industrie de Lyon* **10**: 513–521, pl. 7.

MUST BE:

Mulsant E. 1847d: Description de deux coléoptères nouveaux, constituant chacun une nouvelle coupe générique. *Annales des Sciences Physiques et Naturelles, d’Agriculture et d’Industrie de Lyon* **10**: 515–521, pl. 7.

70. PAGE 883

PRINTED:

Waterhouse C. O. 1889: Coleoptera. Pp. 121–131. — Aitchison J. E. T.: The Zoology of the Afghan Delimitation Commission. *The Transactions of the Linnean Society of London* (2) **5 Zoology** [1888–1894]: 53–142, pls. 6–14.

MUST BE:

Waterhouse C. O. 1889: Coleoptera. Pp. 122–131. — Aitchison J. E. T.: The Zoology of the Afghan Delimitation Commission. *The Transactions of the Linnean Society of London* (2) **5 Zoology** [1888–1894]: 53–142, pls. 6–14.

ACKNOWLEDGEMENTS. I am very grateful to Alexey Gusakov and Andrey Ozerov (Zoological Museum of Moscow University), Andrey Lobanov (Zoological Institute, Sankt-Petersburg) for providing me with the opportunity to study museums’ materials. My special thanks to Karl Adlbauer, Alain Drumont, Martin Egger, Walter Grosser, Dan Heffern, Maxim Lazarev, Ivan Löbl, Tatsuya Niisato, Nobuo Ohbayashi, Sergey Saluk, Gianfranco Sama, Andrey Shapovalov, Ales Smetana, Petr Svacha, Szczepan Ziarko for valuable friendly consultations on many taxonomy problems, their own remarks and loaned specimens.

## References

- Bartenev A.F. 2004. Obzor vidov Zhukov-usachey (Coleoptera: Cerambycidae) fauny Ukrainy. [A review of the long-horned beetles species (Coleoptera: Cerambycidae) of the fauna of Ukraine] // *Izvestiya Kharkovskogo Entomologicheskogo Obshchestva* [The Kharkov Entomological Society Gazette]. Vol.11 [2003]. No.1–2. P.24–43. [in Russian].
- Bartenev A.F. 2009. Zhuki-usachi Levoberezhnoy Ukrainy i Kryma. Kharkov, Kharkovskiy Natsionalnyy Universitet. [Longicorn-beetles of Left-Bank Ukraine and Crimea. Kharkov: Kharkov National University]. 405p. [in Russian].
- Danilevsky M.L. 2009. Species Group Taxa of Longhorned Beetles (Coleoptera, Cerambycidae) Described by N. N. Plavilstshikov and Their Types Preserved in the Zoological Museum of the Moscow State University and in the Zoological Institute of the Russian Academy of Sciences, St. Petersburg // *Entomological Review*. Vol.89. No.6. P.689–720.
- Danilevsky M.L. & Miroshnikov A.I. 1985. Zhuki-drovoseki Kavkaza (Coleoptera, Cerambycidae). *Opredelitel*. Krasnodar: Kubanskiy Selskokhozyaistvennyy Institut. 417 p., 10 pls. [in Russian].
- Esjunin S.L. & Kozminykh V.O. 1992. Materialy k faune zhestkokrylykh zakaznika “Troitzkiy” // *Chlenistonogie okhranyaemykh territoriy Cheliabinskoy oblasti*. Sverdlovsk. UrO AN SSSR. [Materials on the Coleoptera fauna of “Troitzky” reservation // Arthropoda of preserved territories of Cheljabinsk region. Sverdlovsk. Ural Branch of Ac. of Sc of the USSR]. P.59–67. [in Russian].
- Gfeller W. 1972. Cerambycidae (Coleoptera) der Tuerkei. *Persienexpedition 1970 der Herren Dr. h.c. Wittmer und U. v. Bothmer // Mitteilungen der Entomologischen Gesellschaft Basel*. (N.F.) Bd.22. No.1. S.1–8.
- Grebennikov V.V., Gill B.D. & Vigneault R. 2010. *Trichoferus campestris* (Faldermann) (Coleoptera: Cerambycidae), an Asian wood-boring beetle recorded in North America // *Coleopterists Bulletin*. Vol.64. No.1. P.13–20.
- Han Ch. & Niisato T. 2009. Clytine Beetles of the genus *Sclethrus* Newman (Coleoptera, Cerambycidae) // *Special Bulletin of the Japanese Society of Coleopterology*, Tokyo. Vol.7. P.117–126.
- Hayashi M. & Villiers A. 1987. Revision of the Asian Lepturinae (Coleoptera: Cerambycidae) With special reference to the type specimens’ inspection. Part II // *Bulletin of Osaka Jonan Women’s Junior College*. Vol.22. P.1–20. Pls.1–3.
- Heyrovsky L. 1934. Druhý příspěvek ke známosti tribu Lepturini. (Col. Cerambycidae) // *Časopis Československé společnosti entomologické*. Vol.31. P.8–12.
- Inokaitis V. 2004. Naujos ir retos Lietuvos entomofaunos vabalu (Coleoptera) rusys, aptiktos 2000–2003 metais. New and rare for the Lithuanian fauna Coleoptera species found in 2000–2003 // *New and Rare for Lithuania Insect Species Records and Descriptions*. Vol.16. P.7–10.
- Inokaitis V. 2009. Rare and very rare for the Lithuanian fauna Coleoptera species found in 2004–2009 // *New and rare for Lithuania insect species. Records and description*. Vol.21. P.40–44.
- Isaev A.Yu. 2007. *Opredelitel zhestkokrulykh Srednego Povolzhya (chast III. Polyphaga, Phytophaga)*. Seriya “Priroda Ulyanovskoy oblasti”. [A key for beetles of Middle Volga (part III. Polyphaga,



- Phytophaga). Serie "Nature of Ulianovsk Region". No.14. Ulianovsk. Publishing house "Vektor-C". P.1–225 [in Russian].
- Isaev A.Yu., Egorov L.V. & Egorov K.A. 2004. Zhestkokrylye (Insecta, Coleoptera) lesostepi Srednego Povolzhya. Katalog. [Beetles (Insecta, Coleoptera) of forest-steppe of Middle Volga. Catalogue] Ulianovsk. 72 p. [in Russian].
- Isaev A.Yu., Magdeev D.V. 2003. Fauna Zhukov-usachey (Coleoptera, Cerambycidae) lesostepnoy zony Povolzhya // Issledovaniya v oblasti biologii i metodiki ee prepodavaniya. Mezhdunarodnyy sbornik nauchnykh trudov k 100-letiyu D.N. Florova, vyp. 3, ch. 1. [Fauna of Longicorn-beetles (Coleoptera, Cerambycidae) of forest-steppe zone of Volga valley // Investigations in the field of biology and methods of its teaching. Inter-institutes collection of transactions to the 100<sup>th</sup> anniversary of D.N. Flerov, 3(1)] Samara. P.290–300 [in Russian].
- Kaliuzhnaja N.S., Komarov E.V. & Cherezova L. B. 2000. Zhestkokrylye nasekomye Nizhnego Povolzhya. [Coleoptera of lower Volga river]. Volgograd. 204 p. [in Russian].
- Kostin I.A. 1968. Rasprostranenie drovosekov (Coleoptera, Cerambycidae) listvennykh porod v Kazakhstane // Nasekomye — vrediteli selskogo i lesnogo khozyaistva Kazakhstana. Trudy Instituta Zoologii, Akademiya nauk Kaz. SSR. [Distribution of Cerambycidae (Coleoptera) in the leaf-bearing woods in Kazakhstan // Pest Insects of Agriculture and Forests of Kazakhstan. Proceedings of the Institute of Zoology, Academy of Sciences of Kazakhstan SSR]. Vol.30. P.184–197 [in Russian].
- Kostin I.A. 1973. Zhuki-dendrofagi Kazakhstana (koroedy, drovoseki, zlatki). Alma-Ata: Akademiya Nauk Kazakhskoy SSR. 288 p.
- Kryzhanovskiy O.L. 1974. 35. Sem. Cerambycidae — drovoseki ili usachi // Nasekomye i kleshchi — vrediteli selskokhozyaistvennykh kultur. [35. Fam. Cerambycidae — timber beetle or longicorn beetles // Insects and ticks — pests of agricultural crops]. Leningrad: Nauka. P.139–157 [in Russian].
- Lobanov A. L., Danilevsky M. L. & Murzin S. V. 1981. Sistematicheskii spisok usachei (Coleoptera, Cerambycidae) fauny SSSR. I // Entomologicheskoe Obozrenie. Vol.60. P.784–803 [in Russian].
- Löbl I. & Smetana A. (eds.). 2010. Catalogue of Palaearctic Coleoptera. Vol.6. Chrysomeloidea. Stenstrup. Apollo Books. 924pp.
- Magdeev D.V. 1996. Zhuki-usachi (Cerambycidae, Coleoptera) // Flora i fauna zapovednikov. Vyp.61. Bespozvonochnye Zhigulevskogo zapovednika. Moscow. P.39–47 [in Russian].
- Magdeev D.V. 2003. Fauna Zhukov-usachey (Coleoptera, Cerambycidae) Samarskoy oblasti // Kraevedchieskie zapiski, XI. SamarSKIY oblastnoy istoriko-kraevedchieskiy muzey. [Fauna of Longicorn-Beetles (Coleoptera, Cerambycidae) of Samara Region. // Local history notes, XI. Samara Regional Local History Museum]. Samara: Fine Design. P.202–208 [in Russian].
- Martynov V.V. & Pisarenko T.A. 2004. Ecologo-faunisticheskii obzor Zhukov-usachey (Coleoptera: Cerambycidae) yugo-vostochnoy Ukrainy // Izvestiya Kharkovskogo entomologicheskogo obshchestva. [A review of the fauna and ecology of the longhorned beetles (Coleoptera: Cerambycidae) of southeast Ukraine // The Kharkov Entomological Society Gazette]. Vol.11.(2003). No.1–2. P.44–69 [in Russian].
- Mikhailov Yu.E. 1999. Nasekomye muzeya-zapovednika "Arkaim". Zhestkokrylye: vidovoy sostav i zametki po strukture populyatsiy. Prirodnye sistemy Yuzhnogo Urala. Cheliabinsk. Cheliabinskii gos. universitet. [Insects of "Arkaim" museum-natural reserve. Beetles: species composition and notes on the structure of populations // Natural systems of South Urals]. Cheljabinsk. Cheljabinsk University. P.221–248 [in Russian].
- Miroshnikov A.I. 1998. Novaya klassifikatsiya zhukov-drovosekov kompleksa Anoplodera tribu Lepturini (Coleoptera, Cerambycidae) fauny golarktiki. I // Entomologicheskoe Obozrenie. Vol.77. No.2. P.384–421.
- Muraj X. 1960. Inventarizimi i fam. Cerambycidae ne vendin tone. Quelques Cerambycides en Albanie // Bull Univ. Shtet. Tiranes. Ser. Shkencat nat. Vol.14. No.4. P.137–141.
- Naumov R.V. 1994. Zhuki-usachi (Coleoptera, Cerambycidae) Ulyanovskoy oblasti. Ser. "Priroda Ulyanovskoy oblasti". [Longicorn-beetles (Coleoptera, Cerambycidae) of Ulianovsk region. Ser. "The Nature of Ulianovsk Region"]. 5. Ulianovsk. P.43–46 [in Russian].
- Negrobov S.O., Tzurikov M.N., Logvinovskiy V.D., Fomichev A. I., Prokin A.A. & Gilmutdinov K.S. 2005. Otryad Coleoptera. — Kadastr bespozvonochnykh Voronezhskoy oblasti. [Order Coleoptera. Cadastre of Invertebrata of Voronezh region]. Voronezh. P.534–673 [in Russian].
- Niisato T. 2002. New locality of *Salpinia laosensis* (Coleoptera, Cerambycidae), with a note on its systematic position // Elytra. Vol.30. No.1. P.262.
- Niisto T. 2007. Subfamily Cerambycinae Latreille, 1804 // Ohbayashi N. & Niisato T. (eds). Longicorn beetles of Japan. Kanagawa: Tokai Univ. Press. P.424–512
- Ohbayashi N. & Niisato T. 2009. Review of the *Pyrocalimma* Generic-Group sensu Hayashi & Villiers, 1997, with descriptions of new Genus and two new species (Coleoptera, Cerambycidae, Lepturinae) // Longicornists, Special Bulletin of the Japanese Society of Coleopterology. No.7. Tokyo. P.139–167.
- Pisarenko T.A. 1999. Osoblivosti fauni rodini vysachiv (Coleoptera, Cerambycidae) promislivikh i zapovidnikh rayoniv pivdenno-skhidnoy Ukraini // Naukoviy Visnik. Suchasna ekologiya i problemi stalogo rozvitku suspilstva. Vip.97. Lvov. P.149–155 [in Ukrainian]
- Plavilstshikov N.N. 1965. 75. Sem. Cerambycidae — zhuki-drovoseki, usachi // Opredelitel nasekomykh Evropeyskoy chasti SSSR, t.2, zhestkokrylye i veerokrylye. [75-th Fam. Cerambycidae — Timber Beetles, Longicornes // A Key to Insects of the European Part of the USSR, v. 2, Coleoptera and Strepsiptera]. M.-L.: Nauka. P.389–419 [in Russian].
- Romadina K.G. 1954. Drevogryzushchie lichinki Zhukov-usachey (Cerambycidae) doliny reki Urala // Trudy Zoologicheskogo Instituta akademii nauk SSSR. [Woodboring Longicorn (Cerambycidae) beetle-larvae of Ural river valley // Archives of the Zoological Institute of the Academy of Sciences of the USSR]. Vol.16. P.211–228 [in Russian].
- Sama G., Rapuzzi P. & Kairouz A. 2010. Catalogue commenté des Cerambycidae du Liban. An annotated catalogue of the Cerambycidae of Lebanon (Insecta Coleoptera Cerambycidae) // Quaderno di Studi e Notizie di Storia Naturale della Romagna. Vol.30. P.131–201.
- Schultze W. 1920. Eights contribution to the Coleoptera fauna of the Philippines // Philippine Journal of Science. Vol.16. P.191–203. 2 pls.
- Shapovalov A.M., Nemkov V.A., Rusakov A.V. & Shovkun D.F. 2006. Zhuki-usachi (Coleoptera, Cerambycidae) Orenburgskoy oblasti. // Vestnik Orenburgskogo gosudarstvennogo universiteta. Orenburg: Izd-vo OGU. [Longicorn-beetles (Coleoptera, Cerambycidae) of Orenburg Region. — Vestnik of Orenburg State University]. Vol.4. No.54. P.105–109. [in Russian].
- Yen Chia-hsien 1932. A new species of Cerambycidae from Kwangsi // Peking Natural History Bulletin. Vol.7. No.2. P.165–166.
- Zagaikevich I.K. 1979. Nekotorye aspekty khorologii usachey (Coleoptera, Cerambycidae) fauny Ukrainy. — VII Mezhdunarodnyy simpozium po entomofaune Sredney Evropy. Materialy. Leningrad, 19–24 sentyabrya 1977. P.195–197.
- Zagaikevich I.K. 1991. Taksonomiya i ekologiya usachey. Kiev: Naukova Dumka. 178 p. [in Russian].
- Zhuravlev S.M. 1914. Materialy k faune zhukov Uralskoy oblasti // Trudy Russkogo Entomologicheskogo Obshchestva. Vol.41. No.3. P.33–61.