

## ACAROID MITES OF NORTHERN AND EASTERN ASIA (ACARI: ACAROIDEA)

P. B. Klimov<sup>1</sup>, A. V. Tolstikov<sup>2</sup><sup>1</sup> University of Michigan, Museum of Zoology, 1109 Geddes Ave., Ann Arbor, Michigan 48109-1079 USA; email: pklimov@umich.edu<sup>2</sup> Tyumen State University, Department of Zoology and Evolutionary Ecology of Animals, 10 Semakova str., Tyumen, 625003 Russia

ABSTRACT: Acaroid mites (superfamily Acaroidea) are economically and veterinary important pests. We give a list of 5 families 49 genera, and 180 species of acaroid mites recorded in the Northern and Eastern Asia (Asian part of Russia, South Korea, Japan, and China). One new genus, *Sphēcacarus* Klimov gen. n., is described. The following new synonymies and combinations are proposed: *Tortonia* Oudemans, 1911 (= *Hyohondania* Sasa, 1952, syn. nov.); *Sancassania polyphyllae* (Zachvatkin, 1941) (= *Sancassania berteanaeoursae* Samšičák, 1970 syn. nov.); *Mezorhizoglyphus* Kadzhaja, 1966 (= *Boletacarus* Volgin et Mironov, 1980 syn. nov.); *Tortonia kanoi* (Sasa, 1952) comb. nov. (from *Hyohondania*), *Tortonia jinyunensis* (Zhang et Li, 2003) comb. nov. (from *Sinosuidasia*).

KEY WORDS: Acaroidea, Siberia, Russian Far East, South Korea, China, Japan

## INTRODUCTION

Acaroid mites exploit a variety of patchy habitats. They can be found in nests of vertebrates (birds, mammals) and invertebrates (termites, beetles, bees, wasps, ants), fungal sporocarps, decomposing organic matter (such as carrion, various plant materials, animal manure and guano), under bark, in treeholes, and in seaweeds. A few species are aquatic or parasitic on gills of fresh-water crabs. Being serious pests of stored food and products, and a common component of house dust mites, acaroid mites are of economical, medical, and veterinary importance. Such taxa as *Acarus*, *Mycetoglyphus*, *Tyrophagus*, *Rhizoglyphus*, and *Sancassania* (= *Caloglyphus*) are damaging subterranean parts and young leaves of various crops, and are serious agricultural pests and subject to quarantine in many countries (Diaz et al. 2000; Fan and Zhang 2004, 2007; Kasuga and Amano 2006; Nakao 1991; Nakao and Kurosa 1988; Olsen 1983; Rojas and Klimov 2007; Shinkaji et al. 1986).

Some acaroid mites live continuously in a soil environment, disperse through soil and exploit various patchy habitats. Others use soil to disperse to patchy habitats as adults or, indirectly, as deutonymphs. For example, *Sapracarus tuberculatus* (Suidasiidae) lives in deep soil in the Holarctic and occasionally colonizes bird nests. Species of *Sancassania* (*Rhizoglyphoides*) inhabit fruiting bodies of mushrooms and only reproduce several weeks each year when mushrooms are available; they spend the remaining time inactive in soil waiting for the new mushroom season (Klimov and OConnor 2003). Some species of *Tyrophagus*, *Rhizoglyphus*, *Schwiebea*, and *Sancassania* are habitat generalists, and they are common in both soil and patchy habitats, indicating that they may

adopt both strategies. These species are usually the most serious agricultural and stored product pests, and they may cause respiratory diseases or allergic reactions to individuals who come into direct contact with these materials. They can easily extend their ranges via commercial trade and establish themselves in new countries and habitats.

The following list summarizes taxonomic information about mites of the superfamily Acaroidea found in Northern (Asian part of Russia) and Eastern Asia (Japan, China, South and North Korea). The list is based primarily on the senior author original investigations made in 1993–2001 in Central Siberia and the Russian Far East and literature records. We also include some literature records on *Tyrophagus*, *Rhizoglyphus*, *Sancassania* which can not be independently verified at this time. They may be useful as a starting point for future research. Here we note that the superfamily itself may not be non-monophyletic, Horsiindae may be artificial and not part of Acaridae (with *Sennertionyx* is close to *Cerophagus*, currently in Gaudiellidae), Acarinae plus Tyrophaginae plus *Sancassania* and related genera may be monophyletic, and the two genera with two setae on the chelicera (*Viedebantia* and an undescribed genus from *Lycoperdon*) may represent a family-level lineage (Klimov and OConnor 2008; unpubl.).

Abbreviations: ZIN = Zoological Institute of the Russian Academy of Sciences, Saint Petersburg.

## Superfam. ACAROIDEA Latreille, 1802

I. Fam. SCATOGLYPHIDAE Zachvatkin et Volgin, 1955

1. *Scatoglyphus* Berlese, 19131. *Scatoglyphus polytrematus* Berlese, 1913: China (Wang 1964)

- II. Fam. **GAUDIPELLIDAE** Atyeo, Baker et Delfinado, 1974
2. **Cerophagus** Oudemans, 1904
2. *Cerophagus granulatus* (Dujardin, 1849): Primorskiy Kray, ZIN
- III. Fam. **SUIDASIIDAE** Fain et Philips, 1978
3. **Suidasia** Oudemans, 1905 (= *Aphelenia* Oudemans, 1923; *Chibidania* Sasa, 1952)
3. *Suidasia nesbitti* Hughes, 1948 (= *Chibidania tokyoensis* Sasa, 1952): Russian Far East; Japan (Sasa 1952); China (Olsen 1983; Wang 1964), incl. Hong Kong (Olsen 1983); South Korea (Chu et al. 1967; Lee and Choi 1980)
4. *Suidasia pontifica* Oudemans, 1905 (= *Suidasia medianensis* Oudemans, 1924; *Suidasia insectorum* Fox, 1950): Japan (Noge et al. 2005; Oshima 1980); China (Fan et al. 2010; Wang 1964), incl. Hong Kong (Olsen 1983) and Taiwan (Olsen 1983); South Korea (Chu et al. 1967; Jung et al. 2010b)
4. **Sinosuidasia** Jiang, 1996 (differs from *Tortonia* by presence of male paraanal suckers)
5. *Sinosuidasia orientalis* Jiang, 1996: China (Jiang, 1996)
5. **Tortonia** Oudemans, 1911 (= *Ebertia* Oudemans, 1924 (OConnor, pers. comm.); *Neottiglyphus* Volgin, 1974 (OConnor, pers. comm.); *Hyohondania* Sasa, 1952, **syn. nov.**)
6. *Tortonia intermedia* (Oudemans, 1902): "Siberia, basin of Nel'kan river" (Zachvatkin 1941); Yakutia or Khabarovskiy Kray
7. *Tortonia kanoi* (Sasa, 1952) **comb. nov.** (from *Hyohondania*): Japan (Sasa 1952)
8. *Tortonia jinyunensis* (Zhang at Li, 2003) **comb. nov.** (from *Sinosuidasia*): China (Zhang and Li 2002)
9. *Tortonia* sp1. n.: Khabarovskiy Kray
10. *Tortonia* sp2. n.: Primorskiy Kray
11. *Tortonia* sp3. n.: Sakhalin
- IV. Fam. **LARDOGLYPHIDAE** Oudemans, 1927
6. **Lardoglyphus** Oudemans, 1927
12. *Lardoglyphus angelinae* Olsen, 1982: China: Hong Kong (Olsen 1983); USA: California (introduced) (Olsen 1982; Olsen et al. 1987)
13. *Lardoglyphus kanoi* (Sasa et Asanuma, 1951) (= *Hoshikadania kanoi* Sasa et Asanuma, 1951): Japan (Noge et al. 2005; Olsen 1983; Sasa and Asanuma 1951); South Korea (Lee and Choi 1980); China (Fan et al. 2010), incl. Hong Kong (Olsen 1983)
14. *Lardoglyphus zacheri* Oudemans, 1927: Japan (this paper); China, incl. Hong Kong (Fan et al. 2010; Olsen 1983)
15. *Lardoglyphus* sp. n. aff. *zacheri*: ZIN, 1 HDN, 1 F Primorskiy Kray; probably *Lardoglyphus zacheri* in this ref. (Tareev 1970)
7. **Sinolardoglyphus** Jiang, 1991
16. *Sinolardoglyphus nanchangensis* Jiang, 1991: China: Jiangxi (Jiang 1991a)
- V. Fam. **ACARIDAE** Latreille, 1802
- Subfam. **HORSTIINAE** Fain, 1984
8. **Horstia** Oudemans, 1905
17. *Horstia ornata* (Oudemans, 1900): Japan (Oudemans 1900; Zachvatkin 1941) (our data)
18. *Horstia helenae* (Oudemans, 1902): Japan (Zachvatkin 1941) (probably from a historical mainland colony of Japan)
9. **Sennertionyx** Zachvatkin, 1941
19. *Sennertionyx manicati* (Giard, 1890): Primorskiy Kray
10. **Cerophagopsis** Zachvatkin, 1941
20. *Cerophagopsis skorikovi* Zachvatkin, 1941 (= *Cerophagus skorikovi* (Zachvatkin, 1941): Japan (Kurosa 2000; Zachvatkin 1941)
21. *Cerophagopsis indicus* (Potter et Olsen, 1987): China: Taiwan (Fan et al. 2010)
22. *Cerophagopsis* sp. n.: Japan
- Subfam. **ACARINAE** Latreille, 1802
11. **Acarus** Linnaeus, 1758
23. *Acarus bomiensis* Wang, 1982: China (Wang 1982)
24. *Acarus chaetoxysilos* Griffiths, 1970: Japan (Griffiths 1970)
25. *Acarus farris* (Oudemans, 1905) (?= *A. lushanensis* Jiang, 1992; China: Jiangxi OConnor, pers. comm.): Primorskiy Kray, Irkutskaya Oblast' (our data); China (Fan et al. 2010; Jiang 1992)
26. *Acarus fengxianensis* Wang, 1985: China (Wang 1985)
27. *Acarus gracilis* Hughes, 1957: China (Fan et al. 2010)
28. *Acarus immobilis* Griffiths, 1964: Primorskiy Kray, Irkutskaya Oblast' (our data); China (Fan et al. 2010)
29. *Acarus mirabilis* Volgin, 1965: China (Fan et al. 2010)
30. *Acarus siro* Linnaeus, 1758: Primorskiy Kray (Tareev 1970) (may be misidentification); China (Fan et al. 2010) (some records may be misidentifications)
- Subfam. **TYROPHAGINAE** Oudemans, 1923
12. **Bembidioglyphus** Klimov, 1998
31. *Bembidioglyphus acinacisetosus* Klimov, 1998: Kuril Islands: Paramushir (Klimov 1998d)

- 13. *Aleuroglyphus*** Zachvatkin, 1941  
 32. *Aleuroglyphus ovatus* (Troupeau, 1879): Primorskiy Kray (Tareev 1970); Japan (Oshima 1980); South Korea (Lee and Choi 1980); China (Fan et al. 2010)  
 33. *Aleuroglyphus chinensis* Jiang, 1994: China (Jiang 1994)  
 34. *Aleuroglyphus formosanus* Tseng, 1972: China: Taiwan (Tseng 1989)
- 14. *Garsaultia*** Oudemans, 1916  
 35. *Garsaultia* sp.: Japan (Kurosa 2000)
- 15. *Kuzinia*** Zachvatkin, 1941  
 36. *Kuzinia laevis* (Dujardin, 1849): Primorskiy Kray, Irkutskaya Oblast', Kuril Islands (Klimov 1998d); Japan (Kurosa 2000)  
 37. *Kuzinia* sp. n.: Primorskiy Kray, ex *Lygaeus sjostedti*
- 16. *Lackerbaueria*** Zachvatkin, 1941  
 38. *Lackerbaueria cribratissima* Zachvatkin, 1941: Sakhalin, Kuril Islands (Klimov 1997)
- 17. *Mycetoglyphus*** Oudemans, 1932  
 39. *Mycetoglyphus fungivorus* Oudemans, 1932: Primorskiy Kray (Tareev 1970), Irkutskaya Oblast' (Zachvatkin 1953b), Tomskaya Oblast' (Sorokin 1953); Japan (Nakao and Kurosa 1988); China (Fan et al. 2010)
- 18. *Forcellinia*** Oudemans, 1924  
 40. *Forcellinia diamesa* Zachvatkin, 1941: Primorskiy Kray  
 41. *Forcellinia wasmanni* (Moniez, 1892): Primorskiy Kray (Tareev 1970), Kamchatka Kuril, Islands (Klimov 1997)
- 19. *Schulzea*** Zachvatkin, 1941  
 42. *Schulzea caucasica* Zachvatkin, 1941: Irkutskaya oblast', Primorskiy Kray, Sakhalin (Klimov 1997)  
 43. *Schulzea pamirensis* Zachvatkin, 1941: "Pamir"
- 20. *Sphecacarus*** Klimov gen. n.  
 44. *Sphecacarus psenuli* Vitzthum, 1931 (= *Tyroglyphus psenuli* Vitzthum, 1931, *Lackerbaueria psenuli* (Vitzthum, 1931)): Sakhalin (our data); Japan (Kurosa 2000)  
 45. *Sphecacarus* sp. n.: Primorskiy Kray, ex *Carinostigmus rhinocerus*
- 21. *Tyrolichus*** Oudemans, 1923  
 46. *Tyrolichus casei* Oudemans, 1923: Primorskiy Kray (Tareev 1970), Irkutskaya Oblast' (Zachvatkin 1953b); China (Fan et al. 2010)
- 22. *Tyrophagus*** Oudemans, 1924  
 47. *Tyrophagus brevicrinatus* Robertson, 1959: China (Fan et al. 2010)  
 48. *Tyrophagus ganjiangensis* Jiang, 1993: China (Jiang, 1993c)  
 49. *Tyrophagus fanetzhangorum* Klimov et OConnor, 2010: China (intercepted in New Zealand) (Fan and Zhang 2007); Japan (intercepted in New Zealand) (Fan and Zhang 2007)  
 50. *Tyrophagus formicetorum* Volgin, 1948: Irkutskaya oblast', ZIN  
 51. *Tyrophagus jingdezhenensis* Jiang, 1993: China: Jiangxi (Jiang 1993b)  
 52. *Tyrophagus kentinus* Tseng, 1973: China: Taiwan (Tseng 1989)  
 53. *Tyrophagus bambusae* Tseng, 1973: China: Taiwan (Tseng 1989)  
 54. *Tyrophagus longior* (Gervias, 1844): Primorskiy Kray (Tareev 1970), Irkutskaya Oblast' (Zachvatkin 1953b), Novosibirskaya Oblast' (Zachvatkin 1953b); Japan (Noguchi et al. 1998); China (Fan et al. 2010), incl. Taiwan (Tseng 1989)  
 55. *Tyrophagus mimlongior* Jiang, 1993: China: Jiangxi (Jiang 1993a)  
 56. *Tyrophagus mixtus* Volgin, 1948: Tomskaya Oblast' (Volgin 1975)  
 57. *Tyrophagus neiswanderi* Johnston et Bruce, 1965: Japan (Kurosa and Nakao 1993; Nakao, 1991); China (Fan et al. 2010; Jiang 1989)  
 58. *Tyrophagus perniciosus* Zachvatkin, 1941: Primorskiy Kray (Tareev 1970), Novosibirskaya Oblast' (Zachvatkin 1953b), Altayskiy Kray (Zachvatkin 1941); Japan (Nakao 1991; Nakao and Kurosa 1988); China (Fan et al. 2010)  
 59. *Tyrophagus putrescentiae* (Schrank, 1781) (= *T. communis* Fan et Zhang, 2007. Probable synonyms (may be also *T. fanetzhangorum*): *Tyroglyphus muscae* Sasaki, 1921; *Tyroglyphus longior* var. *taiwanensis* Sugimoto, 1938 (Klimov and OConnor, 2009a, b)): Primorskiy Kray (Zachvatkin 1941), Irkutskaya Oblast' (our data), Novosibirskaya Oblast' (Zachvatkin 1953b); South Korea (Jeong et al. 2010; Kim et al. 2003; Lee and Choi 1980; Ree and Lee 1997); Japan (Fan and Zhang 2007; Okabe et al. 2001; Oshima 1980); China, incl. Hong Kong and Taiwan (Fan et al. 2010; Fan and Zhang 2007; Olsen 1983) (records published before 2007 need re-verification to distinguish from *T. fanetzhangorum*)  
 60. *Tyrophagus silvester* Zachvatkin, 1941 (= *Tyrophagus humerosus silvester* Zachvatkin, 1941): Primorskiy Kray (Tareev 1970), Irkutskaya Oblast' (Zachvatkin 1953b) (our data), Novosibirskaya Oblast' (Zachvatkin 1953b)



61. *Tyrophagus similis* Volgin, 1949: Korea (Jung et al. 2010a); Japan (Nakao and Kurosa 1988); China (Fan et al. 2010)
62. *Tyrophagus tropicus* Robertson, 1959: China (Fan et al. 2010; Wang 1985)
63. *Tyrophagus vanheurni* Oudemans, 1924 (= *T. palmarum* sensu Roberson (Fan and Zhang, 2007); ? *T. parvulus* Volgin, 1949 (Volgin, 1975)): "European and Siberian part of the [former] USSR" (as *parvulus*) (Volgin 1966b); China (Jiang 1989), incl. Taiwan (Tseng 1989)
- 23. Tyroborus** Oudemans, 1923
64. *Tyroborus lini* Oudemans, 1923: Japan (Tomita et al. 2003); China (Fan et al. 2010; Fan and Zhang 2006; Jiang 1989)
- 24. Paulacarellus** Fain, 1977
65. *Paulacarellus faini* Klimov, 2001: Primorskiy Kray (Klimov 2000 (2001)b)
- Subfam. **RHIZOGLYPHINAE** Oudemans, 1923
- 25. Australhypopus** Fain et Friend, 1984
66. *Australhypopus formosani* (Phillipsen et Coppel, 1977) (= *Acotyledon formosani* Phillipsen et Coppel, 1977): termite culture brought to USA from Japan (Phillipsen and Coppel 1977)
67. *Australhypopus* sp.: China (Wang et al. 2002)
- 26. gen. n.** (ex *Lycoperdon* spp.)
68. gen. n., sp. n. Primorskiy Kray (our data); Japan (Fig. 27) (Kurosa 2000)
- 27. Viedebanttia** Oudemans, 1929
69. *Viedebanttia* sp. n. (Primorskiy Kray, ex *Silpha peforata*)
70. *Viedebanttia egorovi* Klimov, 1998: Primorskiy Kray, Irkutskaya Oblast' (our data); South Korea (Klimov 1998e)
71. *Viedebanttia macrocnemis* (Zachvatkin, 1941): Primorskiy Kray (Tareev 1970)
- 28. Apiacarus** Volgin, 1974
72. *Apiacarus inflatus* Volgin, 1974: ZIN, Irkutskaya Oblast' (Note: *ve* on the middle of lateral edges of propodosoma shield, *aa* present in this genus)
- 29. Carabidobius** Volgin, 1953
73. *Carabidobius* sp. n.: Primorskiy Kray
- 30. Acotyledon** Oudemans, 1903
74. *Acotyledon paradoxa* Oudemans, 1903: Primorskiy Kray (Tareev 1970); China (Fan et al. 2010; Zou and Wang 1991)
75. *Acotyledon corporis* Ishikawa et Morikawa, 1964: Japan (Ishikawa and Morikawa 1964)
- 31. Neoacotyledon** Samšičák, 1982
76. *Neoacotyledon rhizoglyphoides* (Zachvatkin, 1937): Primorskiy Kray (Tareev 1970), Tomskaya Oblast' (Sorokin 1953), Novosibirskaya Oblast' (Zachvatkin 1941); China (Fan et al. 2010), including Taiwan (Tseng 1989); records need to be verified following incorrect interpretations of this species by Hughes (Hughes 1976) and Türk and Türk (Türk and Türk 1957)
77. *Neoacotyledon sokolovi* (Zachvatkin, 1940): Primorskiy Kray (Tareev 1970), Tomskaya Oblast' (Sorokin 1953), Novosibirskaya Oblast' (Zachvatkin 1941)
- 32. Cosmoglyphus** Oudemans, 1932
78. *Cosmoglyphus absoloni* (Samšičák, 1961): South China (Samšičák 1961), Turkmenistan
79. *Cosmoglyphus batsylevi* (Zachvatkin, 1941) (= *Acotyledon batsylevi* Zachvatkin, 1941): China (Fan et al. 2010)
80. *Cosmoglyphus chinensis* Samšičák, 1966: China (Samšičák 1966)
81. *Cosmoglyphus hosoyai* (Kugoh, 1957) (= *Murodania hosoyai*): Japan (Kugoh 1957)
82. *Cosmoglyphus hughesi* Samšičák, 1966: Japan (Noge et al. 2005; Ryono et al. 2001), China (Fan et al. 2010)
83. *Cosmoglyphus krameri* (Berlese, 1881): South Korea (our data); China: Taiwan (Tseng 1989)
84. *Cosmoglyphus kunshanensis* Zou et Wang, 1991 (= *Sancassania kunshanensis*): China: Jiangsu (Zou and Wang 1991) (see notes for *Cosmoglyphus volgini*)
85. *Cosmoglyphus lishihmeii* (Samšičák, 1961): China (Samšičák 1961) (see notes for *Cosmoglyphus volgini*)
86. *Cosmoglyphus oudemansi* (Zachvatkin, 1937): China (Fan et al. 2010)
87. *Cosmoglyphus redikorzevi* (Zachvatkin, 1937): Irkutskaya Oblast' (Zachvatkin 1937)
88. *Cosmoglyphus volgini* (Zachvatkin, 1941): Primorskiy Kray (Tareev 1970) (HDNs similar but do not match exactly "*Caloglyphus feytaudi*" sensu Türk and Türk (Türk and Türk 1957) non Oudemans, 1928 and *Sancassania kunshanensis*. *C. volgini*, *C. lishihmeii* and *C. kunshanensis* differ from typical *Cosmoglyphus* by genu IV not distinctly longer than femur (distinctly longer in *Cosmoglyphus*), however, hind borders of coxal fields II are close to fore edges of coxal fields III and separated by small, thin, equal on all its length space, and small, shifted to the anterior attachment organ as in true *Cosmoglyphus*. These character states suggests for close affinities of the group including

- C. volgini* with *Australhyopopus*. Adults of “*C. feytaudi*” sensu Türk and Türk and *C. kunshanensis* are more similar to *Sancassania* by filiform *scx*, some tarsal setae dilated at tips (not figured in *kunshanensis*), and anus shifted anteriorly (not so much in *kunshanensis*)
33. *Sancassania* Oudemans, 1916 (= *Caloglyphus* Berlese, 1923, *Isoglyphus* Zachvatkin, 1937; *Mycetosancassania* Klimov, 2000, *Rhizoglyphoides* Volgin, 1978) (Klimov and OConnor, 2003)
89. *Sancassania caroli* (ChannaBasavanna et Krishna Rao, 1982): China: Jiangxi (Tu et al. 2003); India (Channabasavanna et al. 1981)
90. *Sancassania chelone* Oudemans, 1916 (= *Caloglyphus moniezi* Zachvatkin, 1937): Primorskiy Kray (Klimov 1997), Sakhalin, the Republic of Buryatia; Japan (our data)
91. *Sancassania coprophila* Mahunka, 1968: China: Fujian (Zou and Wang 1991); Hungary (Fan et al. 2010)
92. *Sancassania fujianensis* (Zou, Wang et Chang 1987): China (Zhang et al. 1997; Zou et al. 1987), morphologically and biologically very close to *S. chelone*
93. *Sancassania geotruporum* (Zachvatkin, 1941): Sakhalin, Kuril Islands: Kunashir (Klimov 1997); Japan (Klimov 1997; Kurosa 2000)
94. *Sancassania grifolapholiotae* (Klimov, 2000) (= *Mycetosancassania grifolapholiotae*): Irkutskaya oblast', Khabarovskiy Kray (Klimov 2000a; Klimov and OConnor 2003)
95. *Sancassania mandzhur* (Zachvatkin, 1940): Primorskiy Kray (Tareev 1970); China (garlic from Harbin intercepted in USSR) (Zachvatkin 1940)
96. *Sancassania mironovi* Klimov et OConnor, 2003: Irkutskaya oblast', Khabarovskiy Kray (Klimov and OConnor 2003)
97. *Sancassania phyllophagiana* (Oseto et Mayo, 1975): known from USA; record from South Korea (Ree et al. 1997) is probably a misidentification
98. *Sancassania polyphyllae* (Zachvatkin, 1941) (= *Sancassania berteanaeursae* Samšičák, 1970 **syn. nov.**, types of *polyphyllae* studied): Georgia (Zachvatkin 1941); Turkey (our data); China (Samšičák 1957); probably records of “*Caloglyphus polyphyllae*” from Japan (Kuwahara et al. 1989; Leal et al. 1989; Noge et al. 2005) belong to *C. polyphyllae* sensu Türk and Türk 1957 non Zachvatkin
99. *Sancassania rodionovi* (Zachvatkin, 1935) (= *S. berlesei*, auct. non Michael): China, incl. Taiwan (Fan et al. 2010); Primorskiy Kray (Tareev 1970), Irkutskaya Oblast' (our data), Altayskiy Kray (Zachvatkin 1953a); South Korea (Lee and Choi 1980); China (Fan et al. 2010; Timms et al. 1981): incl. Taiwan (Tseng 1989)
100. *Sancassania* sp1. n.: Primorskiy Kray, ex *Lasiopsis golovjankoi*
101. *Sancassania* sp2. n.: South Korea, ex *Rhombonyx testaceipes*
102. *Sancassania shanghaiensis*: China (Zou and Wang 1989); Japan (Noge et al. 2005; Sakata et al. 2001) (close to *S. sphaerogaster*)
103. *Sancassania sphaerogaster* (Zachvatkin, 1937) (= *Caloglyphus michaeli* non Oudemans, 1924 sensu Türk and Türk 1957; *Caloglyphus mycophagus* sensu Hughes, 1976 non Mégnin, 1874; *Caloglyphus striatus* Klimov, 1996): Primorskiy Kray (Klimov 1996b; Tareev 1970); South Korea; Japan (our data); China, including Taiwan (Fan et al. 2010)
104. *Sancassania spinatarsus* (Herman, 1804): Primorskiy Kray (Klimov 1997); Japan (Kurosa 2000; Noge et al. 2005)
105. *Sancassania rodriguezii* Samšičák, 1980: Japan (Mori et al. 1998; Noge et al. 2005)
- Sancassania karnatakaensis* (Krishna Rao et Ranganath, 1982) (excluded): known from India only (Krishna Rao et al. 1982). Recorded for China (Fan et al. 2010) with reference to (Zou et al. 1987), however, this work only compares *fujianensis* with *karnatakaensis*.
34. *Acarotalpa* Volgin, 1966
106. *Acarotalpa* sp. n.: Japan (Okabe, pers. comm.)
35. *Rhizoglyphus* Claparede, 1869
107. *Rhizoglyphus actinidia* Zhang, 1994: China: Hubei (Zhang et al. 1994) (may be *Schwiebea*)
108. *Rhizoglyphus allii* Bu et Wang, 1995: China (Bu and Liu 1997; Bu and Wang 1995)
109. *Rhizoglyphus caladii* Manson, 1972 (= *R. longispinosus* Ho et Chen, 2001): China: Taiwan (Fan and Zhang 2004; Ho and Chen 2001; Tseng 1989)
110. *Rhizoglyphus echinopus* (Fumouze et Robin, 1868) (= *R. callae* Oudemans, 1924; *R. zachvatkini* Volgin, 1952): Japan (Fan and Zhang 2004); China (Bu and Liu 1997; Lin and Ding 1990; Fan et al. 2010), incl. Hong Kong (Manson 1972a) and Taiwan (Tseng 1989)
111. *Rhizoglyphus huainanensis* Zhang, Li et Zhuge, 2000: China (Zhang et al. 2000)
112. *Rhizoglyphus kangdingensis* Wang X.-z., 1983: China (Bu and Liu 1997; Bu and Li 1998; Wang 1983)

113. *Rhizoglyphus narcissi* Lin et Ding, 1990: China (Bu and Liu 1997; Lin and Ding 1990)
114. *Rhizoglyphus robini* Claparède, 1869: Primorskiy Kray (as *echinopus*) (Tareev 1970), Irkutskaya Oblast' (our data), Tomskaya Oblast' (Sorokin 1953); Japan (Fan and Zhang 2004; Kurosa and Nakao 1993; Manson 1972a; Oshima 1980); South Korea (Chu et al. 1967; Lee and Choi 1980; Ree et al. 1997); China (Fan et al. 2010), incl. Taiwan (Tseng 1989)
115. *Rhizoglyphus setosus* Manson, 1972: Japan (*Cycas revolute*) (Fan and Zhang 2004) and imported on rotten rhizomes of *Curcuma roscoeana* from Thailand (Akiyama et al. 1997; Mizoguchi et al. 2005); China: Hong Kong (Manson 1972a) and Taiwan (Fan et al. 2007; Fan and Zhang 2004; Ho and Chen 1987)
116. *Rhizoglyphus singularis* Manson, 1972 (= *R. tsutienensis* Ho et Chen, 2000): China: Taiwan (Fan and Zhang 2004; Ho and Chen 2000)
117. *Rhizoglyphus tardus* Volgin, 1952: Novosibirskaya Oblast' (Volgin 1952)
- 36. *Mezorhizoglyphus*** Kadzhaja, 1966 (= *Boletacarus* Volgin et Mironov, 1980 syn. nov.)
118. *Mezorhizoglyphus bratskensis* Klimov, 1996: Irkutskaya Oblast' (Klimov 1996a)
119. *Mezorhizoglyphus colchicus* Kadzhaja, 1966: Magadanskiy Kray (Volgin 1975), Primorskiy Kray, Irkutskaya Oblast' (Klimov 1996a)
120. *Mezorhizoglyphus sibiricus* (Volgin et Mironov, 1980) (= *Boletacarus sibiricus* Volgin et Mironov, 1980): the Republic of Buryatia (Volgin and Mironov 1980)
121. *Mezorhizoglyphus* sp1. n.: Primorskiy Kray (our data); Japan (Okabe coll.)
122. *Mezorhizoglyphus* sp2. n. aff. *sibiricus*: Primorskiy Kray
- 37. *Kanekobia*** Fain, Yunker, van Goethem, et Johnston, 1982
123. *Kanekobia potamona* (Kaneke et Kadosaka, 1978) (= *Ewingia potamona* Kaneke et Kadosaka, 1978): Japan (Fain et al. 1982; Kaneko and Kadosaka 1978)
- 38. *Boletoglyphus*** Volgin, 1953 (= *Fantovia* Samšičák, 1957; *Ellipsopus* Fain and Ide, 1976; *Lindquistia* Mahunka, 1977)
124. *Boletoglyphus boletophagi* (F. Türk et S. Türk, 1952) (= *Schwiebea boletophagi* F. Türk et S. Türk, 1952; *Boletoglyphus cribrosus* Volgin, 1953): Irkutskaya Oblast' (Klimov 1998b)
125. *Boletoglyphus extremiorientalis* Klimov, 1998: Primorskiy Kray, Khabarovskiy Kray, Sakhalin (Klimov 1998b); Japan (Klimov 1998b) (as *boletophagi*) (Kurosa 2000)
- 39. *Capillaroglyphus*** Klimov, 1998
126. *Capillaroglyphus polypori* Klimov, 1998: Primorskiy Kray (Klimov 1998a); Japan (Klimov 1998a) (Fig. 29) (Kurosa 2000)
- 40. *Terglyphus*** Samšičák, 1965
127. *Terglyphus padrtorum* Samšičák, 1965: China (Samšičák, 1965)
- 41. *Schwiebea*** Oudemans, 1916 (= *Megninietta* Jacot, 1936; *Robinisca* Zachvatkin, 1941; *Troupeauia* Zachvatkin, 1941)
128. *Schwiebea araujoae* Fain, 1977: Japan (Okabe et al. 2008), China: Taiwan (Ho 1993)
129. *Schwiebea callae* Jiang, 1991: China (Jiang 1991b)
130. *Schwiebea chinica* Samšičák, 1965: China (Samšičák 1965)
131. *Schwiebea cuncta* Ho, 1993: China: Taiwan (Ho 1993)
132. *Schwiebea danielopoli* Fain, 1982: Sakha (Yakutia) Republic; Eastern Siberia, Western Sayan Mountains (Bugrov 1995)
133. *Schwiebea elongata* (Banks, 1906) (= *S. athiasae* Fain 1977, *S. barbei* Cooremani 1959, *S. codognoensis* Fain et Pagani 1989, *S. obesa* Fain et Fauvel 1988, *S. receptacula* Manson 1972): Kuril Islands: Kunashir (as *receptacula*) (Bugrov 1995); Japan (Okabe et al. 2008; Okabe and Oconnor 2000)
134. *Schwiebea estradai* Fain et Ferrando, 1991: Japan (Okabe et al. 2008)
135. *Schwiebea falticis* Woodring, 1966: Japan (tentatively identified) (Woodring 1966)
136. *Schwiebea jiangxiensis* Jiang, 1995: China: Jiangxi (Jiang 1995)
137. *Schwiebea kurilensis* Bugrov, 1995: Kuril Islands: Kunashir (Bugrov 1995)
138. *Schwiebea lebruni* Fain, 1977: Japan (Okabe et al. 2008)
139. *Schwiebea longibursata* Fain et Wauthy, 1979: Sakhalin (Bugrov 1995), Primorskiy Kray, Irkutskaya Oblast' (Bugrov 1995, 1997)
140. *Schwiebea meilingensis* Jiang, 1997: China (Jiang 1997)
141. *Schwiebea mertzis* Woodring, 1966: Japan (tentative identification) (Woodring 1966); China: Taiwan (Tseng 1989)
142. *Schwiebea neomycolicha* Klimov, 1998: Primorskiy Kray (Klimov 1998c)
143. *Schwiebea nova* (Oudemans, 1906) (= *Schwiebea rossica* Zachvatkin, 1941): Primorskiy Kray



- (Tareev 1970); Irkutskaya Oblast', Novosibirskaya Oblast' (Klimov 1998c)
144. *Schwiebea parallela* (J. Müller, 1860) (= *Schwiebea tshernyshevi* Zachvatkin, 1941; *Schwiebea eurynymphae* (non Oudemans, 1910) sensu F. Türk, E. Türk 1957): Kamchatka (Klimov 1998c), Kuril Islands: Kunashir (Bugrov 1995), Paramushir (Klimov 1998c), Primorskiy Kray (Tareev 1970), Novosibirskaya Oblast' (Bugrov 1995)
145. *Schwiebea ruienensis* Fain et Wauthy, 1979: Sakhalin, Irkutskaya oblast', Western Sayan Mountains (Bugrov 1995)
146. *Schwiebea sakhalinensis* Bugrov, 1995: Sakhalin (Bugrov 1995)
147. *Schwiebea similis* Manson, 1972: Japan (Okabe et al. 2008); China, incl. Hong Kong (Manson 1972b; Fan et al. 2010)
148. *Schwiebea* sp1. n.: Primorskiy Kray (heavily sclerotized)
149. *Schwiebea* sp2 n.: South Korea, ex *Necrodes asiaticus*
150. *Schwiebea* sp3. n.: Primorskiy Kray
151. *Schwiebea* sp4. n.: Primorskiy Kray
152. *Schwiebea* sp5. n.: Kuril Islands, ex *Sphecodes* sp.
153. *Schwiebea taiwanensis* Ho, 1993: China: Taiwan (Ho 1993)
154. *Schwiebea talpa* Oudemans, 1906 (= *Schwiebea pachyderma* Zachvatkin, 1941): Primorskiy Kray (Tareev 1970), Irkutskaya Oblast' (our data)
155. *Schwiebea xianggangensis* Jiang, 1998: China (Jiang 1998)
156. *Schwiebea zhangzhouensis* Lin et Lin, 2000: China: Fujian (Lin and Lin 2000)
157. *Schwiebea zingiberi* Manson, 1972: China: Hong Kong (Manson 1972b)
- 42. *Lamtoglyphus*** Fain, 1974
158. *Lamtoglyphus longiretinalis* (Klimov, 1998) (= *Michaelopus longiretinalis* Klimov, 1998): South Korea (Klimov 1998 (1999)) (this genus is different from *Schwiebea* only by the presence of eye spots in heteropomorphic deutonymphs, and has been implicitly synonymized with *Schwiebea* (Okabe et al. 2008). In addition, *Lamtoglyphus* aff. *coineaui* (Okabe et al. 2008) and *Lamtoglyphus* sp. (Kurosa 2000) were recorded for Japan.
- 43. *Fagacarus*** Fain et Norton, 1979
159. *Fagacarus* sp. n.: Primorskiy Kray (Klimov 2000 (2001)a)
- 44. *Histiogaster*** Berlese, 1883 (= *Tyroglyphopsis* Vitzthum, 1926)
160. *Histiogaster carpio* (Kramer, 1881): China (Jiang 1994)
161. *Histiogaster cercus* (Zhang, 1994) (= *Thyreophagus cercus* Zhang, 1994): China: Hubei (Zhang et al. 1994). Tentative generic assignment; the posterior lobe resembles that of *Reckiacarus*, however, the incomplete dorsal setation suggest close affinities with *Histiogaster*.
162. *Histiogaster filippovi* Zachvatkin, 1941: Primorskiy Kray (Zachvatkin 1941)
163. *Histiogaster hylecoeti* Cooreman, 1952: Japan (Kurosa 2000)
164. *Histiogaster orientalis* Volgin, 1966: Khabarovskiy Kray (Volgin 1966a)
165. *Histiogaster ornatus* Volgin, 1966: Primorskiy Kray (Volgin 1966a)
166. *Histiogaster rotundus* Woodring, 1966: Japan (Hiroyuki et al. 2003)
167. *Histiogaster* sp1. n.: Primorskiy Kray, decaying wood on beach
168. *Histiogaster* sp2. n.: Primorskiy Kray, oak sap flow
- 45. *Reckiacarus*** Kadzhaja, 1972
169. *Reckiacarus* sp1. n.: Irkutskaya Oblast'
170. *Reckiacarus* sp2. n.: Primorskiy Kray
171. *Reckiacarus* sp3. n.: Sakhalin and Kuril Islands, ex *Rhopalum*. Tentative generic assignment.
- 46. *Calvoliella*** Samšičák, 1961
172. *Calvoliella laphriae* (Samšičák, 1956): Khabarovskiy Kray, Primorskiy Kray, Amurskaya Oblast', Irkutskaya Oblast', Kemerovskaya Oblast' (Klimov 2000b); probably Japan (Fig. 28) (Kurosa 2000)
- 47. *Umakefeq*** Klimov, 2000
173. *Umakefeq macrophthalmus* Klimov, 2000: Primorskiy Kray (Klimov 1999 (2000))
174. *Umakefeq mesoophthalmus* Klimov, 2000: Primorskiy Kray (Klimov 1999 (2000))
175. *Umakefeq microphthalmus* Klimov, 2000: Primorskiy Kray (Klimov, 1999 (2000))
- 48. *Thyreophagus*** Rondani, 1874 (= *Monieziella* Berlese, 1897; *Fumouzeia* Zachvatkin, 1953; *Michaelopus* Fain et Johnston, 1974)
176. *Thyreophagus corticalis* Michael, 1885: Primorskiy Kray
177. *Thyreophagus entomophagus* (Laboulbène, 1852): Novosibirskaya Oblast' (Zachvatkin 1953b); China (Fan et al. 2010), incl. Taiwan (Olsen 1983; Tseng 1989)
178. *Thyreophagus gallegoi* Portus et Gomez, 1982: China (Fan et al. 2010; Jiang 1991b), incl. Hong Kong (Olsen 1983)

**49. gen. n.**

179. gen. n. sp. n.: Primorskiy Kray, ex *Lycoperdina mandarina*

**Incertae sedis**

180. *Tyroglyphus eriocheir* Pearse, 1930: China (Pearse 1930). This taxon has some resemblance to *Pontoppidania*, and probably, like this genus, belongs to the subfamily Tyrophaginae, family Acaridae.

*Sinoglyphus lentinusi* Zou et Wang, 1989: probably nom. nud.; cited in this ref. (Zhaopeng 1992), then in this ref. (Fan et al. 2010) for ref. (Zou and Wang 1989), but not found in ref. (Zou and Wang 1989).

***Sphecacarus Klimov* gen. n.**

Type species *Tyroglyphus psenuli* Vitzthum, 1931 (= *Sphecacarus psenuli* (Vitzthum, 1931), comb. n.)

**Diagnosis.** *Adults.* By short, filiform *ve* (situated at same transverse line with *vi*) and heavily barbed *scx* resembles *Paraforcellinia* Kadzhaja, 1973 and *Aleuroglyphus* Zachvatkin, 1941. From these taxa differs by solenidion  $\sigma'$  shorter than  $\sigma''$  (longer in *Aleuroglyphus*, all *Forcellinia*, unknown in *Paraforcellinia*). We anticipate that in *Paraforcellinia* (briefly described, ex ant nest, very similar to *Forcellinia*) solenidion  $\sigma'$  is shorter than  $\sigma''$ . Setae *vi* 2–2.5 times shorter than *ve* in *Paraforcellinia*, while in *Sphecacarus* *ve* are more than 3 times shorter than *vi*. *Sphecacarus* differs from *Aleuroglyphus* by the following (character states of *Aleuroglyphus* are in parentheses): *si* as long as *se* (*si* distinctly shorter); *scx* not flattened, evenly barbed (flattened, with barbed edges); dorsal hysterosomal setae nearly the same in length, barbed (*c*<sub>1</sub>, *c*<sub>2</sub>, *d*<sub>1</sub>, *d*<sub>2</sub>, *e*<sub>1</sub>, *e*<sub>2</sub>, *h*<sub>1</sub> distinctly shorter than other hysterosomal setae, smooth); *u*, *v*, *s* I–II spiniform, with normal ends (spiniform, with hooked ends).

*Heteromorphic deutonymphs.* Similar to *Forecellinia*, *Mycetoglyphus*, and *Tyrohagus* in having conoids *ps*<sub>2</sub> completely anterior to median sucker *ad*<sub>1+2</sub>. By this character state can be readily distinguished from *Lackerbaueria* (which has *ps*<sub>2</sub> nearly on the same level as *ad*<sub>1+2</sub>). Differs from *Forcellinia* and the two related genera by filiform and pectinated seate *cG* and *mG* I–II, and the absence of solenidion  $\omega_2$  (*cG* and *mG* I–II spiniform and smooth;  $\omega_2$  present in *Forecellinia*, *Mycetoglyphus*, and *Tyrohagus*).

**Species included.** *Sphecacarus americanus* (Baker, 1962) comb. n. (= *Lackerbaueria americana* Baker, 1962); *S. psenuli* (Vitzthum, 1931)

comb. n. (= *Sphecacarus psenuli* Vitzthum, 1931, ?= *Lackerbaueria krombeini* Baker, 1962).

**ACKNOWLEDGEMENTS**

We thank Barry OConnor for discussion, Kimiko Okabe and Sergey Mironov for unpublished data and specimens. This work is supported by grants from the Russian Ministry of Education and Science 02.740.11.5139 to PBK and AVT, 2.2.3.1/13701 to AVT, and the US National Science Foundation to Barry OConnor (NSF DEB-0613769).

**REFERENCES**

- Akiyama, M., Sakata, T., Mori, N., Kato, T., Amano, H. and Kuwahara, Y. 1997. Chemical ecology of astigmatid mites. 46. Neryl formate, the alarm pheromone of *Rhizoglyphus setosus* Manson (Acarina: Acaridae) and the common pheromone component among four *Rhizoglyphus* mites. *Applied Entomology and Zoology*, 32: 75–79.
- Bu, G. and Liu, H. 1997. Study of genus *Rhizoglyphus* in China. *Xinan Nongye Daxue Xuebao*, 19: 78–84.
- Bu, G. and Wang, F. 1995. A new species of the genus *Rhizoglyphus* (Acari: Acaridae) from Shaanxi, China. *Entomotaxonomia*, 17: 309–312.
- Bu, G.-S. and Li, L.-S. 1998. Taxonomic notes on and key to known species of the genus *Rhizoglyphus* (Acari: Acaridae) from China. *Systematic and Applied Acarology*: 179–182.
- Bugrov, S.A. 1995. [New species of the genus *Schwiebia* (Astigmata, Acaridae) from Russia and adjacent countries]. *Zoologicheskyy Zhurnal*, 74: 61–75. [in Russian]
- Bugrov, S.A. 1997. [Free-living Astigmata (Acari-formes) of the Moscow District fauna]. *Zoologicheskyy Zhurnal*, 76: 147–156. [in Russian]
- Channabasavanna, G.P., Krishna Rao, N.S. and Ranganath, H.R. 1981. A new *Caloglyphus* (Astigmata: Acaridae) from poultry litter in India, with taxonomic comments on the genus. *Indian Journal of Acarology*, 6: 57–63.
- Chu, J.K., Song, S.B., Kim, D.K. and Kim, Y.K. 1967. Epidemiological studies on the acaroid mite [sic!]. *Korean Journal of Parasitology*, 5: 71–77.
- Diaz, A., Okabe, K., Eckenrode, C.J., Villani, M.G. and OConnor, B.M. 2000. Biology, ecology, and management of the bulb mites of the genus *Rhizoglyphus* (Acari: Acaridae). *Experimental and Applied Acarology*, 24: 85–113.
- Fain, A., Yunker, C.E., van Goethem, J. and Johnston, D.E. 1982. Notes on the Ewingiidae (Acari, Astigmata) living in association with pagurids and fresh-water crabs (Crustacea), with description of a new genus and a new species. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique Entomologie*, 54: 1–7.



- Fan, Q.-H., Chen, Y. and Wang, Z.-Y. 2010. Acaridia (Acari: Astigmatina) of China: A review of research progress. *Zoosymposia*, 4.
- Fan, Q.-H., Su, X.-X. and Chen, Y. 2007. Species, hosts, distribution and inspection techniques of *Rhizoglyphus* from Taiwan. *Kunchong Zhishi*, 44: 596–602.
- Fan, Q.-H. and Zhang, Z.-Q. 2004. *Revision of Rhizoglyphus Claparède (Acari: Acaridae) of Australasia and Oceania*. Systematic and Applied Acarology Society, London. 374 pp.
- Fan, Q.-H. and Zhang, Z.-Q. 2007. *Tyrophagus (Acari: Astigmata: Acaridae)*. Manaaki Whenua Press, Lincoln, New Zealand.
- Fan, Q.H. and Zhang, Z.Q. 2006. Revision of *Tyrobobus* Oudemans, 1924 (Acari: Astigmata: Acaridae). *Zootaxa*: 1–43.
- Griffiths, D.A. 1970. A further systematic study of the genus *Acarus* L., 1758 (Acaridae, Acarina), with a key to species. *Bulletin of the British Museum (Natural History) Zoology*, 19: 85–118.
- Hiroyuki, H., Naoki, M., Kimiko, O., Ritsuo, N. and Yasumasa, K. 2003. Chemical ecology of astigmatid mites LXIX. Neryl formate [3,7-dimethyl-(Z)-2,6-octadienyl formate] as the alarm pheromone of an acarid mite, *Histiogaster rotundus* Woodring (Acari: Acaridae). *Applied Entomology and Zoology*, 38: 379–385.
- Ho, C.-C. 1993. Two new species and a new record of *Schwiebea* Oudemans from Taiwan (Acari: Acaridae). *International Journal of Acarology*, 19: 45–50.
- Ho, C.-C. and Chen, J.-S. 1987. A new record of bulb mite, *Rhizoglyphus setosus* Manson, (Acarina: Acaridae) from Taiwan. *Journal of Agricultural Research of China*, 36: 237–238.
- Ho, C.-C. and Chen, W.-H. 2000. A new species of *Rhizoglyphus* Claparede (Acari: Acaridae) infesting bulbs from Taiwan. *Chinese Journal of Entomology*, 20: 347–351.
- Ho, C.C. and Chen, W.H. 2001. A new species of *Rhizoglyphus* Claparede (Acari: Acaridae) from Taiwan infesting the taro and giant alocasia. *Plant Protection Bulletin (Taichung)*, 43: 47–49.
- Hughes, A.M. 1976. *The mites of stored food and houses*. Her Majesty's Stationery Office, London.
- Ishikawa, K. and Morikawa, K. 1964. On a new mite, *Acotyledon corporis*, n. sp. found on a dead body. *Reports of Research of Matsuyama Shinonome Girls School*, 1: 41–46.
- Jeong, K.Y., Kim, C. and Yong, T.S. 2010. Enzymatic activities of allergen extracts from three species of dust mites and cockroaches commonly found in Korean home. *Korean Journal of Parasitology*, 48: 151–155.
- Jiang, Z. 1989. Three new records of genus *Tyrophagus* from Jiangxi, China. *Journal of Nanchang University (Natural Science)*, 13: 22–27.
- Jiang, Z. 1992. A new species of Acaridae from China (Acari: Acariformes). *Journal of Nanchang University (Natural Science)*, 16: 239–242, 253.
- Jiang, Z. 1993a. A new species of *Tyrophagus* from China (Acari: Acariformes). *Journal of Nanchang University (Natural Science)*, 17: 83–86.
- Jiang, Z. 1993b. A new species of *Tyrophagus* from Jiangxi (Acari: Acariformes). *Journal of Nanchang University (Natural Science)*, 17: 85–88.
- Jiang, Z. 1995. A new species of Acaridae from China (Acari: Acariformes). *Journal of Nanchang University (Natural Science)*, 19: 30–33.
- Jiang, Z.-t. 1991a. A new genus and species of Acaridae from China (Acari: Acariformes). *Acta Zootaxonomica Sinica*, 16: 61–65.
- Jiang, Z.-T. 1991b. A new species and a genus and three newly recorded species of the Acaridae, Glycyphagidae from China (Acariformes, Acaroidea). *Jiangxi Science*, 9: 240–244.
- Jiang, Z.-T. 1993c. *Tyrophagus ganjiangensis* sp. nov. form China (Acari: Acaridae). *Jiangxi Science*, 11: 37–41.
- Jiang, Z.-T. 1994. A new species and a genus and species newly recorded of the family Acaridae from China (Acari: Acaroidea). *Jiangxi Science*, 12: 188–122.
- Jiang, Z.-T. 1996. A new genus and species of Acaridae from China (Acari: Acariformes). *Acta Zootaxonomica Sinica*, 21: 76–82.
- Jiang, Z.-T. 1997. A new species of *Schwiebea* from China (Acari: Acariformes). *Journal of Nanchang University (Natural Science)*, 22: 299–301.
- Jiang, Z.-T. 1998. A new species of *Schwiebea*: *Schwiebea xianggangensis* (Acari: Acariformes). *Journal of Nanchang University (Natural Science)*, 22: 120–123.
- Jung, J.-A., Cho, M.-R., Kim, H.-H., Kang, T.-J., Lee, J.-H. and Do, K.-R. 2010a. Damages by *Tyrophagus similis* (Acari: Acaridae) in greenhouse spinach in Korea. *Korean Journal of Applied Entomology*, 49: 429–432.
- Jung, Y., Hong, S. and Lee, W. 2010b. New record on *Suidasia medanensis* Oudemans 1924 from stored food in Korea (Astigmata: Acaridae). *Korean Journal of Applied Entomology*, 49: 273–276.
- Kaneko, K. and Kadosaka, T. 1978. A new parasitic mite (Astigmata: Ewingidae) from the gills of fresh-water crab in Japan. *Japanese Journal of Sanitary Zoology*, 29: 313–315.
- Kasuga, S. and Amano, H. 2006. Infestation of *Tyrophagus similis* Volgin (Acari: Acaridae) on spinach during the seed germination period. *Journal of the Acarological Society of Japan*, 15: 69–73.
- Kim, E.H., Kim, H.K., Choi, D.H. and Ahn, Y.J. 2003. Acaricidal activity of clove bud oil compounds against *Tyrophagus putrescentiae* (Acari: Acaridae). *Applied Entomology and Zoology*, 38: 261–266.

- Klimov, P.B. 1996a. [A review of the Palaearctic genus *Mezorhizoglyphus* Kadzhaja, 1966 (Acari, Acaridae)]. *Far Eastern Entomologist*, 25: 1–8. [in Russian]
- Klimov, P.B. 1996b. [A new species of acaroid mites of the genus *Caloglyphus* (Acari, Acaridae) from the Russian Far East]. *Zoologicheskyy Zhurnal*, 75: 613–619. [in Russian]
- Klimov, P.B. 1997. New data on the fauna and biology of acarid mites (Acari, Acaridae) from the Far East. *Acarina*, 5: 69–72.
- Klimov, P.B. 1998a. [A new tribe of acarid mites of the subfamily Rhizoglyphinae (Acariformes, Acaridae)]. *Far Eastern Entomologist*, 59: 1–19. [in Russian]
- Klimov, P.B. 1998b. [Review of the genus *Boletoglyphus* (Acariformes, Acaridae)]. *Zoologicheskyy Zhurnal*, 77: 1121–1127. [in Russian]
- Klimov, P.B. 1998c. [To the knowledge of acarid mites of the genus *Schwiebea* (Acariformes, Acaridae) from the Far East, with notes on systematic of the genus]. *Vestnik Zoologii*, 32: 13–30. [in Russian]
- Klimov, P.B. 1998d. [To the knowledge of mites and ticks (Acari) of Kuril Islands]. *Far Eastern Entomologist*, 36: 1–36. [in Russian]
- Klimov, P.B. 1998e. [*Viedebanttia egorovi* sp. n., a new acarid mite from South Korea, with notes on the genus (Acariformes: Acaridae)]. *Zoosystematica Rossica*, 7: 251–254. [in Russian]
- Klimov, P.B. 1998 (1999). *Michaelopus longiretinalis* sp. nov. (Acari, Acaridae), a new species of acaroid mite associated with beetle *Onthophagus lenzi* (Coleoptera, Scarabaeidae) from South Korea. *Acarina*, 6: 17–19.
- Klimov, P.B. 1999 (2000). A description of a new genus, *Umakefeq* gen. n., including three new species of mycetophagous acarid mites (Acariformes, Acaridae) from eastern Palaearctic. *Acarina*, 7: 93–106.
- Klimov, P.B. 2000a. [A review of acarid mites of the tribe Caloglyphini (Acaridae, Acariformes) with description of a new genus and species from Siberia and Russian Far East]. *Vestnik Zoologii*, 34: 27–35. [in Russian]
- Klimov, P.B. 2000b. [Bionomy of acarid mite *Calvoliella laphriae* (Samšičák, 1956) (Acariformes, Acaridae), phoretic on flies of the genus *Laphria* (Diptera, Asilidae)]. *Chteniya Pamyati Alekseya Ivanovicha Kurentsova*, 9: 91–104. [in Russian]
- Klimov, P.B. 2000 (2001)a. Notes on systematics, distribution and ecology of mites of the genus *Fagacarus* (Acariformes: Acaridae). *Acarina*, 8: 105–115.
- Klimov, P.B. 2000 (2001)b. *Paulacarellus faini* sp. n., a new species of acarid mite (Acariformes: Acaridae) from the Russian Far East. *Acarologia* (Paris), 41: 249–253.
- Klimov, P.B. and OConnor, B.M. 2003. Phylogeny, historical ecology and systematics of some mushroom-associated mites of the genus *Sancassania* (Acari: Acaridae), with new generic synonymies. *Invertebrate Systematics*, 17: 469–514.
- Klimov, P.B. and OConnor, B.M. 2008. Origin and higher-level relationships of psoroptidian mites (Acari: Astigmata: Psoroptidia): evidence from three nuclear genes. *Molecular Phylogenetics and Evolution*, 47: 1135–1156.
- Klimov, P.B. and OConnor, B.M. 2009a. Conservation of the name *Tyrophagus putrescentiae*, a medically and economically important mite species (Acari: Acaridae). *International Journal of Acarology*, 35: 95–114.
- Klimov, P.B. and OConnor, B.M. 2009b. *Tyrophagus putrescentiae* (Schrank, 1781) (Acariformes, Acaridae): proposed conservation of usage by designation of a replacement neotype. *Bulletin of Zoological Nomenclature*, 66: Case 3501.
- Krishna Rao, N.S., Ranganath, H.R. and Channabasavanna, G.P. 1982. *Caloglyphus karnatakaensis* sp. nov. (Acari: Acaridae) from India, with taxonomic comments on the genus *Caloglyphus*. *Indian Journal of Acarology*, 7: 37–43.
- Kugoh, T. 1957. A new genus and species of acarid (tyroglyphid) mite from Tokyo, Japan. *Bulletin of the Tokyo Medical and Dental University*, 4: 73–79.
- Kurosa, K. 2000. Mites associated with insects and small mammals in Oda-chô, Ehime Prefecture, southwest Japan, In: Odamiyama-no-Shizen-Henshû-Inkai, Yamamoto, E. (Eds.). Nature of Odamiyama, Oda-chô, Ehime, pp. 897–946.
- Kurosa, K. and Nakao, H. 1993. Acaridae, In: S. Ehara (Ed.). Plant mites of Japan in colors. Zenkoku Noson Kyoiku Kyokai, pp. 154–161.
- Kuwahara, Y., Leal, W.S., Suzuki, T., Maeda, M. and Masutani, T. 1989. Antifungal activity of *Caloglyphus polyphyllae* sex pheromone and other mite exudates. *Naturwissenschaften*, 76: 578–579.
- Leal, W.S., Kuwahara, Y. and Suzuki, T. 1989.  $\beta$ -acaridial, the sex pheromone of the acarid mite *Caloglyphus polyphyllae*. Pheromone study of acarid mites, 21 (1). *Naturwissenschaften*, 76: 332–333.
- Lee, W.-K. and Choi, W.-Y. 1980. Studies on the mites (order Acarina) in Korea. I. Suborder Sarcopiformes. *Korean Journal of Parasitology*, 18: 119–144.
- Lin, Z. and Ding, T. 1990. A new species of the genus *Rhizoglyphus* (Acarina: Acaridae) from China. *Entomotaxonomia*, 12: 69–72.
- Lin, Z. and Lin, B. 2000. A new species of *Schwiebea* from Fujian, China (Acarina: Acaridae). *Entomological Journal of East China*, 9: 12–14.
- Manson, D.C.M. 1972a. A contribution to the study of the genus *Rhizoglyphus* Claparede, 1869 (Acarina: Acaridae). *Acarologia*, 13: 621–650.
- Manson, D.C.M. 1972b. Three new species, and a redescription of mites of the genus *Schwiebea* (Acarina: Tyroglyphidae). *Acarologia*, 14: 71–80.

- Mizoguchi, A., Murakami, K., Shimizu, N., Mori, N., Nishida, R. and Kuwahara, Y. 2005. S-isorobinal as the female sex pheromone from an alarm pheromone emitting mite, *Rhizoglyphus setosus*. *Experimental and Applied Acarology*, 36: 107–117.
- Mori, N., Fukui, M. and Kuwahara, Y. 1998. Mating behavior of the astigmatid mite, *Caloglyphus rodriguezii* Samsinak (Acarina: Acaridae). *Applied Entomology and Zoology*, 33: 385–390.
- Nakao, H. 1991. Studies on acarid mites (Acari: Astigmata) damaging vegetable plants. 2. Damage to vegetable seedlings. *Japanese Journal of Applied Entomology and Zoology*, 35: 303–309.
- Nakao, H. and Kurosa, K. 1988. Description of four species of acarid mites newly recorded from Japan, with reference to the damage caused to crops (Acari: Astigmata). *Japanese Journal of Applied Entomology and Zoology*, 32: 135–142.
- Noge, K., Mori, N., Tanaka, C., Nishida, R., Tsuda, M. and Kuwahara, Y. 2005. Identification of astigmatid mites using the second internal transcribed spacer (ITS2) region and its application for phylogenetic study. *Experimental and Applied Acarology*, 35: 29–46.
- Noguchi, S., Mori, N., Kurosa, K. and Kuwahara, Y. 1998. Chemical ecology of astigmatid mites XLIX. beta-acaridial (2(E)-(4-methyl-3-pentenylidene)-butanedial), the alarm pheromone of *Tyrophagus longior* Gervais (Acarina: Acaridae). *Applied Entomology and Zoology*, 33: 53–57.
- Okabe, K., Hinomoto, N. and BM, O.C. 2008. Ecological and morphological attributes of parthenogenetic Japanese *Schwiebea* species (Acari: Acaridae). *Experimental and Applied Acarology*, 44: 77–88.
- Okabe, K., Miyazaki, K. and Yamamoto, H. 2001. Population increase in mushroom pest mites on cultivated *Hypozygus marmoreus* and their vectoring of weed fungi between mushroom cultivation media. *Japanese Journal of Applied Entomology and Zoology*, 45: 75–81.
- Okabe, K. and Oconnor, B.M. 2000. Morphometric and systematic analyses of populations of the *Schwiebea barbei* — group (Acari: Acaridae) with particular reference to populations from North America and Japan. *International Journal of Acarology*, 26: 115–126.
- Olsen, A.R. 1982. A new pest of dried fish from the orient, *Lardoglyphus angelinae* new sp. (Acarina, Acaridae). *Journal of Stored Products Research*, 18: 181–188.
- Olsen, A.R. 1983. Food-contaminating mites from imported foods entering the united states through southern California. *International Journal of Acarology*, 9: 189–193.
- Olsen, A.R., Bryce, J.R., Lara, J.R., Madenjian, J.J., Potter, R.W., Reynolds, G.M. and Zimmerman, M.L. 1987. Survey of stored-product and other economic pests in import warehouses in Los-Angeles. *Journal of Economic Entomology*, 80: 455–459.
- Oshima, S. 1980. Family Acaridae, In: S. Ehara (Ed.). Illustrations of the mites and ticks of Japan. Zenkoku Noson Kyoiku Kyokai, Tokio, pp. 380–385.
- Oudemans, A.C. 1900. Further notes on Acari. *Tijdschrift voor Entomologie*, 43: 109–128.
- Pearse, A.S. 1930. Parasites of Fukien crabs. *Proceedings of the Natural History Society of Fukien Christian University, Foochow, China*, 3: 10–18.
- Phillipsen, W.J. and Coppel, H.C. 1977. *Acotyledon formosani* sp. n. associated with the Formosan subterranean termite, *Coptotermes formosanus* Shiraki (Acarina: Acaridae — Isoptera: Rhinotermitidae). *Journal of the Kansas Entomological Society*, 50: 399–409.
- Ree, H.-I., Jeon, S.-H., Lee, I.-Y., Hong, C.-S. and Lee, D.-K. 1997. Fauna and geographical distribution of house dust mites in Korea. *Korean Journal of Parasitology*, 35: 9–17.
- Ree, H.-I. and Lee, I.-Y. 1997. Development of mass rearing technique of *Tyrophagus putrescentiae* (Acari: Acaridae) found in house dust. *Korean Journal of Parasitology*, 35: 149–154.
- Rojas, E.W. and Klimov, B.M. 2007. Mites of the genus *Rhizoglyphus* (Acari: Acaridae) infesting cultivated bulbs in central and southern Chile, with taxonomic notes on *Acarus hyacinthi* Boisduval and *Rhizoglyphus frickorum* Nesbitt. *International Journal of Acarology*, 33: 87–90.
- Ryono, A., Mori, N., Okabe, K. and Kuwahara, Y. 2001. Chemical ecology of astigmatid mites LVIII. 2-Hydroxy-6-methylbenzaldehyde: Female sex pheromone of *Cosmoglyphus hughesi* Samsinak (Acari: Acaridae). *Applied Entomology and Zoology*, 36: 77–81.
- Sakata, T., Okabe, K. and Kuwahara, Y. 2001. Structural elucidation of twelve novel esters composed of five fatty acids and three new branched alcohols together with four monoterpenoids from *Sancasania shanghaiensis* (Acari: Acaridae). *Bioscience Biotechnology and Biochemistry*, 65: 919–927.
- Samšínák, K. 1957. Příspěvek k poznání roztočů čeledi Tyroglyphidae žijících na broucích z čeledi Lamellicornia (Acari). *Časopis České společnosti entomologické*, 54: 180–188.
- Samšínák, K. 1961. Die termitophilen Acari aus China. *Časopis České společnosti entomologické*, 58: 193–207.
- Samšínák, K. 1965. Termitophile Milben aus der VR China. 2. Acaridoidea. *Reichenbachia*, 5: 291–294.
- Samšínák, K. 1966. Die Neuerrichtung der Gattung *Cosmoglyphus* Oudemans, 1932, gleichzeitig ein Beitrag zum Problem der ‘Copra itch’. *Zoologischer Anzeiger*, 176: 27–42.



- Sasa, M. 1952. Two new acaroid (tyroglyphoid) mites of possible sanitary or economic importance in Japan. *Japanese Journal of Experimental Medicine*, 22: 511–521.
- Sasa, M. and Asanuma, K. 1951. A new food infesting acarid mite, *Hoshikadania konoii* n.g., n. sp., with notes on the comparative-morphology of its different sexes and stages. *Japanese Journal of Experimental Medicine*, 21: 209–222.
- Shinkaji, N., Okabe, K. and Amano, H. 1986. Rhizoglypline mites (Acari: Acaridae) infesting rakkyo (*Allium chinense* G. Don) and Chinese chive (*A. tuberosum* Rottler), and their susceptibility to dimethoate and disulfoton. *Japanese Journal of Applied Entomology and Zoology*, 30: 285–289.
- Sorokin, S.V. 1953. [Notes on ecology of bread mites (Acarina, Tyroglyphoidea) in ant nests]. *Entomologicheskoe obozrenie*, 33: 209–213. [in Russian]
- Tareev, V.N. 1970. *Akaroidnye kleshchi (Acaroidea) Primorskogo Kraya. Abstract of Candidate of Science Thesis [=Acaroid mites (Acaroidea) of Primorskiy Kray]*. Avtoreferat dissertatsii kandidata biologicheskikh nauk., Vladivostok. pp. [in Russian]
- Timms, S., Ferro, D.N. and Emberson, R.M. 1981. General biology and nomenclature of *Sancassania berleseii* (Michael). *Acarologia*, 22: 385–390.
- Tomita, A., Shimizu, N., Mori, N., Nishida, R., Nakao, H. and Kuwahara, Y. 2003. Chemical ecology of astigmatid mites. LXXI. Neryl formate (Z)-3,7-dimethyl-1-2,6-octadienyl formate as the alarm pheromone of *Tyroborus lini* Oudemans 1924, and its recovery after forced discharge. *Journal of the Acarological Society of Japan*, 12: 11–19.
- Tseng, Y.H. 1989. Redescription of *Tyrophagus kentinus* Tseng, *Tyrophagus bambusae* Tseng and *Aleuroglyphus formosanus* Tseng and describes a new record species *Rhizoglyphus canadii* Manson. *Chinese Journal of Entomology, Special Publication*, 3: 37–50.
- Tu, D., Zhu, Z.-M., Xia, B., Yu, L.-P. and Gong, Z.-Q. 2003. A new record of the genus *Caloglyphus* from China (Acari: Acaridae). *Acta Arachnologica Sinica*, 12: 24–26.
- Türk, E. and Türk, F. 1957. Systematik und Ökologie der Tyroglyphiden Mitteleuropas, In: H.-J. Stammer (Ed.). Beiträge zur Systematik und Ökologie mitteleuropäischer Acarina. Band 1. Tyroglyphidae und Tarsonemini. Akademische Verlagsgesellschaft Geest and Portig K.-G., Leipzig, pp. 4–384.
- Volgin, V.I. 1952. [New species of the genus *Rhizoglyphus* Clap. (Acarina, Tyroglyphidae)]. *Zoologicheskyy Zhurnal*, 31: 249–252. [in Russian]
- Volgin, V.I. 1966a. [New genus and new species of mites of the family (Acarina, Acariiformes)]. *Trudy Zoologicheskogo instituta akademii nauk SSSR*, 37: 286–296. [in Russian]
- Volgin, V.I. 1975. [Family Acaridae, Glycyphagidae and Saprogllyphidae]. In: M.S. Gilyarov (Ed.). Opredelitel' obitayushchikh v pochve kleshchey. Sarcopiformes. [Identification key to soil inhabiting mites. Sarcopiformes]. Nauka Publ., Moscow, pp. 416–476. [in Russian]
- Volgin, V.I., Kadzhaya, G. Sch. 1966b. [On geographical distribution of stored food mites in the USSR]. Nauka Publ., Moscow. 54–55 pp. [in Russian]
- Volgin, V.I. and Mironov, S.V. 1980. [New genus and new species of mites of the family Acaridae (Acariiformes, Acaroidea)]. *Parazitologicheskyy sbornik Zoologicheskogo Instituta akademii nauk SSSR*, 29: 169–179. [in Russian]
- Wang, C.L., Powell, J.E. and O'Connor, B.M. 2002. Mites and nematodes associated with three subterranean termite species (Isoptera: Rhinotermitidae). *Florida Entomologist*, 85: 499–506.
- Wang, S.-C. 1964. Five new records to Chinese Acariidae. *Acta Entomologica Sinica*, 13: 900–902.
- Wang, X.-Z. 1982. Acarina: Acariidae — Acarus, In: Z. K.Q.G.Z.K.K.D.T.s.o.t.C.S.E.t.t.Q.-X. Plateau] (Ed.), Xizang Kunchong. [Insects of Xizang]. Science Press, Peking, pp. 471–473.
- Wang, X.-Z. 1983. A new species of roots mite from Western of Sichuan Plateau, China (Acari, Acariidae). *Contributions from Shanghai Institute of Entomology*, 3: 243–246.
- Wang, X.-Z. 1985. A survey of the Acaroidea occurring in the east China. *Contributions from Shanghai Institute of Entomology*, 5: 351–357.
- Woodring, J.P. 1966. North American Tyroglyphidae (Acari): II. The genus *Schwiebea*, with descriptions of four new species. *Proceedings of the Louisiana Academy of Sciences*, 29: 85–112.
- Zachvatkin, A.A. 1937. [Study on systematics of mites of the family Tyroglyphidae]. *Uchenye zapiski Moskovskogo gosudarstvennogo universiteta*, 13: 169–202. [in Russian]
- Zachvatkin, A.A. 1940. [Key to mites damaging stored agricultural products in the USSR]. *Uchenye zapiski Moskovskogo gosudarstvennogo universiteta*, 42: 7–68. [in Russian]
- Zachvatkin, A.A. 1941. [Tyroglyphoid mites Tyroglyphoidea]. In: S.A. Zernov (Ed.). Fauna SSSR: Paukoobraznuyye. Akademiya Nauk SSSR, Moscow-Leningrad, p. 475. [in Russian]
- Zachvatkin, A.A. 1953a. [Investigations on the morphology and postembryonic development of tyroglyphids (Sarcopiformes, Tyroglyphoidea)]. In: V.B. Dubinin, Lange, A.B. (Eds.). Zachvatkin A.A. Sbornik nauchnykh rabot [=Collection of Scientific Works]. Moscow State University, Moscow, pp. 19–120. [in Russian]
- Zachvatkin, A.A. 1953b. [Some results of study of the bread mite fauna in the USSR], Sbornik nauchnykh rabot [=Collection of Scientific Works]. Moscow University Press, Moscow, pp. 169–176. [in Russian]

- Zhang, A.-H. and Li, Y.-R. 2002. A new species of the genus *Sinosuidasia* from China (Acari: Acaridae). *Acta Zootaxonomica Sinica*, 27: 483–485.
- Zhang, H., Li, C.-P. and Zhuge, H.-P. 2000. A new species of genus *Rhizoglyphus* from China (Acari-formes: Acaridida: Acaridae). *Acta Arachnologica Sinica*, 9: 72–74.
- Zhang, J.-z., Liu, J.-y., Xu, W.-d. and Zhao, Z.-s. 1997. Studies on the biology of *Caloglyphus fujianensis* Zou, Wang et Zhang (Acarina: Acaridae). *Wuyi Science Journal*, 13: 221–228.
- Zhang, Z.-f., Jiang, J.-g. and Zeng, H.-w. 1994. Two new species of Acaridae (Acari) from China. *Acta Entomologica Sinica*, 37: 374–377.
- Zhaopeng, S. 1992. Acarid mites, In: X. Chen, Ma, E. (Eds.). Research of Acarology in China. Chongqing Publishing House, Chongqing, pp. 37–40.
- Zou, P., Wang, H.z. and Chang, J.z. 1987. A new species of *Caloglyphus* parasitizing white grubs (Acari: Acaridae). *Wuyi Science Journal*, 7: 115–120.
- Zou, P. and Wang, X. 1989. A new species and two new records of Acaridae associated with edible fungi from China (Acarina: Acaroidea). *Acta Agriculturae Shanghai*, 5: 21–24.
- Zou, P. and Wang, X. 1991. A new species and two new records of *Caloglyphus* associated with mushroom from China. *Journal of Shanghai Agricultural College*, 9: 297–306.