

A new dermestid beetle species,
Anthrenus (Anthrenops) fugong sp.n.
(Coleoptera: Dermestidae: Megatominae), from Yunnan China

Новый вид *Anthrenus (Anthrenops) fugong* sp.n.
(Coleoptera: Dermestidae: Megatominae) из Китая (Юньнань)

J. Háva

И. Гава

Daugavpils University, Institute of Life Sciences and Technology, Department of Biosystematics, Vienības Str. 13, Daugavpils LV-5401 Latvia; Private Entomological Laboratory and Collection, Rýznerova 37, 62 Unětice u Prahy, Praha-západ CZ-252 Czech Republic. E-mail: jh.dermestidae@volny.cz.

Даугавпилсский университет, Институт естественных наук и технологий, отдел биосистематики, Даугавпилс, Латвия;
Частная энтомологическая лаборатория и коллекция, Прага, Чехия.

Key words: Taxonomy, new species, Coleoptera, Dermestidae, *Anthrenus*, *Anthrenops*, Yunnan, China.

Ключевые слова: таксономия, новый вид, Coleoptera, Dermestidae, *Anthrenus*, *Anthrenops*, Юньнань, Китай.

Abstract. *Anthrenus (Anthrenops) fugong* sp.n. from China (Yunnan) is described, illustrated and compared with related Chinese species. A list of the species so far known from China is provided.

Резюме. В статье описывается новый вид *Anthrenus (Anthrenops) fugong* sp.n. из Юньнаня (Китай). Приводится список видов рода *Anthrenus*, известных из Китая.

Introduction

When identifying some dermestids collected by the Czech entomologist in Yunnan, China, an undescribed species of the genus *Anthrenus* Geoffroy, 1762 belonging to the subgenus *Anthrenops* Reitter, 1881 was found. The genus *Anthrenus* currently includes somewhat more than 250 valid species or subspecies worldwide [Háva, 2015]. Up to now, 26 different species of the genus *Anthrenus* are known from China [Háva, 2001, 2004, 2005, 2015; Háva, Kadej, 2006; Kadej, Háva 2015; Hermann, Háva, 2019].

Material and Methods

Nomenclature and systematic in this paper follow Háva [2015].

The size of the beetles and their body parts can be useful in species recognition, so following measurements were made: total length (TL) — linear distance from anterior margin of pronotum to apex of elytra; elytral width (EW) — maximal linear transverse distance.

The type specimens of the described species are provided with a red, printed label showing the following text: «HOLOTYPE [respectively PARATYPE], *Anthrenus (Anthrenops) fugong* sp.n. Jiří Háva det. 2019».

Following abbreviation refer to the collections, in which the examined type materials are deposited: JHAC — Jiří Háva, Private Entomological Laboratory & Collection, Prague-west, Czech Republic.

Results

Anthrenus (Anthrenops) fugong Háva, sp.n.
Figs 1–6.

Type material. Holotype (♂): SW China, Yunnan prov., Gaoligong Shan Mts., 1600 m, 26°52' N 98°51' E, 25–27.V.2016, Guquancun SW of Fugong, P. Kabátek lgt., (JHAC). Paratypes (2♂♂): the same data as holotype (JHAC).

Description. Body TL 2.4–2.5 mm, EW 1.6–1.7 mm; body brown-black, small, oval. Dorsal surface covered by brown and white scales. Individual scales small, broad, subtriangular. Head covered by intermixed brown and white scales. Antennae 9 antennomered, antennomeres I–VII brown, VIII–IX black, antennal club 2 antennomered, compact (Fig. 3). Forehead with median ocellus. Eyes with entire median margin. Palpi brownish-black. Pronotum covered by brown scales discally (intermixed individual white scales) and with white scales on lateral margins (Figs 1–2). Scutellum small, triangular without scales. Elytra with brown and white scales; white scales forming spots on each elytron, other parts covered by brown scales. Epipleuron with brown scales. Ventral surface covered with white and brown scales. Prosternum only with white scales. Metasternum only with white scales, with a small patch at lateral margins from brown scales. Abdominal ventrites I–V without spots in the middle and with large black spots at antero-lateral margins (Fig. 4). 9th ventrite with large dark brown spots medially (Fig. 6). Legs brown with white scales and white setae. Male aedeagus (Fig. 5).

Female. Unknown.

Differential diagnosis. Within the genus *Anthrenus* the subgenus *Anthrenops* is distinctly characterised by the 9-segmented antenna. From other *Anthrenops* species from eastern Palaearctic Region, the



Figs 1–6. *Anthrenus fugong* sp.n.: 1 — habitus, dorsal aspect, holotype, 2 — habitus, dorsal aspect, paratype, 3 — antenna of male, 4 — abdomen, ventral aspect, 5 — genitalia, ventral aspect, 6 — 9th ventrite.

Рис. 1–6. *Anthrenus fugong* sp.n.: 1 — внешний вид, сверху, голотип, 2 — внешний вид, сверху, параптип, 3 — усиок самца, 4 — брюшко, 5 — гениталии самца, вентрально, 6 — 9-й вентрит.

new species differs by the form of antennae as well as shape of the genitalia and arrangement of scales on elytra.

Etymology. Toponymy, named according to Fugong County in China (Fig. 7).

Supplemented list of Chinese *Anthrenus* species

Anthrenus Geoffroy, 1762

- Anthrenus (Anthrenodes)* Chobaut, 1898
- Anthrenus debilis* Háva, 2005
- Anthrenus maculifer* Reitter, 1881
- Anthrenus (Anthrenops)* Reitter, 1881
- Anthrenus fugong* sp.n. (Fig. 7)
- Anthrenus (Anthrenus)* Geoffroy, 1762
- Anthrenus flavipes* LeConte, 1854
- Anthrenus latefasciatus* Reitter, 1892
- Anthrenus nipponensis* Kalík et Ohbayashi, 1985
- Anthrenus oceanicus* Fauvel, 1903
- Anthrenus pimpinellae* (Fabricius, 1775)
- Anthrenus scrophulariae* (Linnaeus, 1758)
- Anthrenus (Florilinus)* Mulsant et Rey, 1868
- Anthrenus cimrmani* Háva, 2005
- Anthrenus emili* Herrmann et Háva, 2019
- Anthrenus flavidus* Solsky, 1876
- Anthrenus museorum* (Linnaeus, 1761)
- Anthrenus qinlingensis* Háva, 2004
- Anthrenus (Nathrenus)* Casey, 1900
- Anthrenus bomiensis* Háva, 2004
- Anthrenus knizeki* Háva, 2004
- Anthrenus kucerae* Háva, 2005

Anthrenus longisetosus Kadej et Háva, 2015

Anthrenus pilosus Pic, 1923

Anthrenus propinquus Háva, 2005

Anthrenus schawalleri Háva et Kadej, 2006

Anthrenus sichuanicus Háva, 2004

Anthrenus tryznai Háva, 2004

Anthrenus turnai Háva, 2004

Anthrenus verbasci (Linnaeus, 1767)

Anthrenus (Solskinus) Mroczkowski, 1962

Anthrenus (Solskinus) becvari Háva, 2004

Anthrenus (Solskinus) sinensis Arrow, 1915




Fig. 7. Location of Fugong County (*) and Nujiang Prefecture (**) within Yunnan province.

Рис. 7. Расположение округа Фугонг (розовый) и префектуры Нуцзян (желтый) в провинции Юньнань.

Acknowledgements

I am very indebted to colleague Petr Kabátek (Prague, Czech Republic) for providing me the interesting material.

References

- Háva J. 2001. New and little known species of the genus *Anthrenus* O. F. Müller, 1764 (Coleoptera: Dermestidae) from World // Entomologische Zeitschrift, Stuttgart. Vol.111. P.52–61.
- Háva J. 2004. New interesting Dermestidae (Coleoptera) from China // Entomologische Zeitschrift, Stuttgart. Vol.114. P.225–232.
- Háva J. 2005. New Dermestidae (Coleoptera) from China. Part 2. Descriptions of four new species of *Anthrenus* Müller 1764 // Entomologische Zeitschrift, Stuttgart. Vol.115. P.155–157.
- Háva J. 2015. World Catalogue of Insects. Vol.13. Dermestidae (Coleoptera). Leiden/Boston: Brill, xxvi + 419 p.
- Háva J., Kadej M. 2006. Three new species of *Anthrenus* O.F. Müller, 1764 from Botswana, India and China (Coleoptera: Dermestidae) // Mitteilungen des Internationalen Entomologischen Vereins. Vol.31. P.119–130.
- Herrmann A., Háva J. 2019. Description of *Anthrenus (Florilinus) emili* sp. nov. from China province Sichuan (Coleoptera: Dermestidae: Megatominae) // Studies and Reports, Taxonomical Series. Vol.15. No.1. P.63–67.
- Kadej M., Háva J. 2015. A new species of *Anthrenus* Geoffroy, 1762 from China, with revised checklist of the Chinese species // The Coleopterists Bulletin. Vol.69. No.3. P.459–462.

Поступила в редакцию 19.9.2019